

## AXIOMATIZATION OF QUASIGROUPS

JONATHAN D.H. SMITH

*Department of Mathematics, Iowa State University*  
*Ames, Iowa 50011–2064, U.S.A.*

**e-mail:** jdsmith@math.iastate.edu  
<http://orion.math.iastate.edu/jdsmith/>

### Abstract

Quasigroups were originally described combinatorially, in terms of existence and uniqueness conditions on the solutions to certain equations. Evans introduced a universal-algebraic characterization, as algebras with three binary operations satisfying four identities. Now, quasigroups are redefined as heterogeneous algebras, satisfying just two conditions respectively known as hypercommutativity and hypercancellativity.

**Keywords:** quasigroup, heterogeneous algebra, hyperidentity.

**2000 Mathematics Subject Classification:** Primary: 20N05;  
Secondary: 08A68.

### REFERENCES

- [1] L. Euler, *Recherches sur une nouvelle espèce de carrés magiques*, Mém. de la Société de Vlislingue **9** (1779), pp. 85 ff.
- [2] T. Evans, *Homomorphisms of non-associative systems*, J. London Math. Soc. **24** (1949), 254–260.
- [3] I.M. James, *Quasigroups and topology*, Math. Zeitschr. **84** (1964), 329–342.
- [4] H.O. Pflugfelder, *Quasigroups and Loops: Introduction*, Heldermann, Berlin, 1990.
- [5] A. Sade, *Quasigroupes obéissant à certaines lois*, Rev. Fac. Sci. Univ. Istanbul, Ser. A **22** (1957), 151–184.

- [6] J.D.H. Smith and A.B. Romanowska, *Post-Modern Algebra*, Wiley, New York, NY, 1999.
- [7] W. Taylor, *Hyperidentities and hypervarieties*, Aequationes Math. **23** (1981), 30–49.

Received 27 February 2006  
Revised 31 March 2006