

## SUBLATTICES CORRESPONDING TO VERY TRUE OPERATORS IN COMMUTATIVE BASIC ALGEBRAS

IVAN CHAJDA

*Department of Algebra and Geometry*  
*Palacký University Olomouc, 17. listopadu 12*  
*771 46 Olomouc, Czech Republic*  
**e-mail:** ivan.chajda@upol.cz

AND

FILIP ŠVRČEK

*Department of Algebra and Geometry*  
*Palacký University Olomouc, 17. listopadu 12*  
*771 46 Olomouc, Czech Republic*  
**e-mail:** filip.svrcek@upol.cz

### Abstract

We introduce the concept of very true operator on a commutative basic algebra in a way analogous to that for fuzzy logics. We are motivated by the fact that commutative basic algebras form an algebraic axiomatization of certain non-associative fuzzy logics. We prove that every such operator is fully determined by a certain relatively complete sublattice provided its idempotency is assumed.<sup>1</sup>

**Keywords:** commutative basic algebra, very true operator, idempotent operator, relatively complete sublattice.

**2010 Mathematics Subject Classification:** 06B23, 03G25, 03B45.

### REFERENCES

- [1] M. Botur and F. Švrček, *Very true on CBA fuzzy logic*, Math. Slovaca **60** (2010) 435–446. doi:10.2478/s12175-010-0023-9
- [2] M. Botur, I. Chajda and R. Halaš, *Are basic algebras residuated lattices?*, Soft Comp. **14** (2010) 251–255. doi:10.1007/s00500-009-0399-z

---

<sup>1</sup>This work is supported by the project CZ.1.07/2.3.00/20.0051 Algebraic Methods in Quantum Logic.

- [3] I. Chajda, R. Halaš and J. Kühr, *Distributive lattices with sectionally antitone involutions*, Acta Sci. Math. (Szeged) **71** (2005) 19–33.
- [4] I. Chajda, R. Halaš and J. Kühr, *Semilattice Structures* (Heldermann Verlag (Lemgo, Germany), 2007).
- [5] P. Hájek, *On very true*, Fuzzy Sets and Systems **124** (2001) 329–333.
- [6] L.A. Zadeh, *A fuzzy-set-theoretical interpretation of linguistic hedges*, J. Cybern. **2** (1972) 4–34.

Received 31 March 2014

Revised 17 July 2014