

ON THE SUBSEMIGROUP GENERATED BY ORDERED IDEMPOTENTS OF A REGULAR SEMIGROUP

ANJAN KUMAR BHUNIYA AND KALYAN HANSDA

Department of Mathematics, Visva-Bharati
Santiniketan – 731235, India

e-mail: anjankbhuniya@gmail.com, kalyanh4@gmail.com

Abstract

An element e of an ordered semigroup S is called an ordered idempotent if $e \leq e^2$. Here we characterize the subsemigroup $\langle E_{\leq}(S) \rangle$ generated by the set of all ordered idempotents of a regular ordered semigroup S . If S is a regular ordered semigroup then $\langle E_{\leq}(S) \rangle$ is also regular. If S is a regular ordered semigroup generated by its ordered idempotents then every ideal of S is generated as a subsemigroup by ordered idempotents.

Keywords: ordered regular, ordered inverse, ordered idempotent, downward closed, completely regular.

2010 Mathematics Subject Classification: 20M10, 06F05.

REFERENCES

- [1] A.K. Bhuniya and K. Hansda, *Complete semilattice of ordered semigroups*, communicated.
- [2] C. Eberhart, W. Williams and I. Kinch, *Idempotent-generated regular semigroups*, *J. Austral. Math. Soc.* **15** (1) (1973) 35–41. doi:10.1017/S1446788700012726
- [3] T.E. Hall, *On regular semigroups whose idempotents form a subsemigroup*, *Bull. Austral. Math. Soc.* **1** (1969) 195–208. doi:10.1017/S0004972700045950
- [4] T.E. Hall, *Orthodox semigroups*, *Pacific J. Math.* **1** (1971) 677–686. doi:10.2140/pjm.1971.39.677
- [5] T.E. Hall, *On regular semigroups*, *J. Algebra* **24** (1973) 1–24. doi:10.1016/0021-8693(73)90150-6
- [6] K. Hansda, *Bi-ideals in Clifford ordered semigroup*, *Discuss. Math. Gen. Alg. and Appl.* **33** (2013) 73–84. doi:10.7151/dmgaa.1195

- [7] K. Hansda, *Regularity of subsemigroups generated by ordered idempotents*, Quasigroups and Related Systems **22** (2014) 217–222. quasigroups.eu/contents/download/2014/22-21.pdf
- [8] N. Kehayopulu, *On completely regular poe-semigroups*, Math. Japonica **37** (1) (1992) 123–130.
- [9] D.B. McAlister, *A note on congruences on orthodox semigroups*, Glasgow J. Math. **26** (1985) 25–30. doi:10.1017/S0017089500005735
- [10] J.C. Meakin, *Congruences on orthodox semigroups*, J. Austral. Math. Soc. **XII** (3) (1971) 222–341. doi:10.1017/S1446788700009794
- [11] J.C. Meakin, *Congruences on orthodox semigroups II*, J. Austral. Math. Soc. **XIII** (3) (1972) 259–266. doi:10.1017/S1446788700013665
- [12] J.E. Milles, *Certain congruences on orthodox semigroups*, Pacific J. Math. **64** (1) (1976) 217–226. doi:10.2140/pjm.1976.64.217
- [13] M. Yamada, *Orthodox semigroups whose idempotents satisfy a certain identity*, Semigroup Forum **6** (1973) 113–128. doi:10.1007/BF02389116

Received 16 June 2015
Revised 18 September 2015