Discussiones Mathematicae General Algebra and Applications 39 (2019) 277–278 doi:10.7151/dmgaa.1319



THE CLONE OF $K^*(n, r)$ -FULL TERMS

KHWANCHEEWA WATTANATRIPOP

Department of Mathematics, Faculty of Science Khon Kaen University, Khon Kaen 40002, Thailand

e-mail: khwancheewa12@gmail.com

AND

THAWHAT CHANGPHAS

Department of Mathematics, Faculty of Science Khon Kaen University, Khon Kaen 40002, Thailand Centre of Excellence in Mathematics, CHE, Si Ayuttaya Rd. Bangkok 10400, Thailand

e-mail: thacha@kku.ac.th

Dedicated to Professor Klaus Denecke on the occasion of 75th birthday

Abstract

Let τ_n be a type of algebras in which all operation symbols have arity n, for a fixed $n \geq 1$. For $0 < r \leq n$, this paper introduces a special kind of n-ary terms of type τ_n called $K^*(n,r)$ -full terms. The set of all $K^*(n,r)$ -full terms of type τ_n is closed under the superposition operation S^n ; hence forms a clone denoted by $clone_{K^*(n,r)}(\tau_n)$. We prove that $clone_{K^*(n,r)}(\tau_n)$ is a Menger algebra of rank n. We study $K^*(n,r)$ -full hypersubstitutions and the related $K^*(n,r)$ -full closed identities and $K^*(n,r)$ -full closed varieties. A connection between identities in $clone_{K^*(n,r)}(\tau_n)$ and $K^*(n,r)$ -full closed identities is established. The results obtained generalize the results of Denecke and Jampachon [K. Denecke and P. Jampachon, *Clones of full terms*, Algebra and Discrete Math. 4 (2004) 1–11].

Keywords: restricted range transformation, full term, clone, Menger algebra, hypersubstitution, hyperidentity, solid variety.

2010 Mathematics Subject Classification: 08B15, 08A62, 08B05.

References

- K. Denecke, Menger algebras and clones of terms, East-West J. Math. 5 (2003) 179–193.
- [2] K. Denecke and P. Jampaclon, Clones of full terms, Algebra and Discrete Math. 4 (2004) 1–11.
- [3] K. Denecke and L. Freiberg, The algebra of strongly full terms, Novi Sad J. Math. 34 (2004) 87–98.
- B.M. Schein and V.S. Trochimenko, Algebras of multiplace functions, Semigroup Forum 17 (1979) 1–64. doi:10.1007/BF02194309
- [5] K. Denecke and S.L. Wismath, Hyperidentities and Clones (Gordon and Breach Science Publisher, 2000). doi:10.1201/9781482287516
- J.M. Howie and R.B. McFadden, Idempotent rank in finite full transformation semigroups, Proc. Roy. Soc. Edinb. Sect. A 114 (1990) 161–167. doi:10.1017/S0308210500024355
- S. Puapong and S. Leeratanavalee, The algebra of generalized full terms, Int. J. Open Problems Compt. Math. 4 (2011) 54–65. doi:10.1155/2013/396464
- [8] P. Sarawut, Some algebraic properties of generalized clone automorphisms, Acta Universitatis Apulensis 41 (2015) 165–175. doi:10.17114/j.aua.2015.41.13

Received 11 June 2019 Revised 18 July 2019 Accepted 18 July 2019