

RELATIVELY COMPLEMENTED ORDERED SETS

IVAN CHAJDA AND ZUZANA MORÁVKOVÁ

Department of Algebra and Geometry, Palacký University of Olomouc
Tomkova 40, 779 00 Olomouc, Czech Republic
e-mail: chajda@risc.upol.cz

Abstract

We investigate conditions for the existence of relative complements in ordered sets. For relatively complemented ordered sets with 0 we show that each element $b \neq 0$ is the least one of the set of all upper bounds of all atoms contained in b .

Key words and phrases: modular ordered set, complemented, relatively complemented ordered set, atom.

1991 Mathematics Subject Classification: 06A99.

Let (A, \leq) be an ordered set and $B \subseteq A$. Denote by

$$L(B) = \{x \in A; x \leq b \text{ for all } b \in B\},$$
$$U(B) = \{x \in A; b \leq x \text{ for all } b \in B\}.$$

If $B = \{b_1, \dots, b_n\}$, we shall write briefly $L(b_1, \dots, b_n)$ or $U(b_1, \dots, b_n)$ instead of $L(B)$ or $U(B)$, respectively. Moreover, for $B, C \subseteq A$ we write $L(B, C)$ for $L(B \cup C)$ and $U(B, C)$ for $U(B \cup C)$. Following [5], an ordered set (A, \leq) is *modular* if for every $a, b, c \in A$ it holds:

$$a \leq c \Rightarrow L(c, U(a, b)) = L(U(a, L(b, c))).$$

Modular ordered sets were treated in [2], a special sort of them, the so called distributive ordered sets were investigated in [2] and [4].

Let (A, \leq) be an ordered set and $a \in A$. An element $b \in A$ is called a *complement of a* if

$$L(U(a, b)) = A \quad \text{and} \quad U(L(a, b)) = A.$$

Complemented ordered sets were studied in [1]. A generalization of the complement called a pseudocomplement in an ordered set was introduced in [5].

It is well known that if L is a complemented modular lattice, then L is also relatively complemented. The aim of our paper is to find a generalization of this result for ordered sets. However, there are several possibilities how to introduce the concept of a relative complement in an ordered set. We can pick up the following two:

Definition. Let (A, \leq) be an ordered set, $a, b \in A$ and $a \leq b$. Let $x \in [a, b] = \{z \in A : a \leq z \leq b\}$. An element $y \in [a, b]$ is called a *weak relative complement of x in $[a, b]$* if

$$\begin{aligned} U(x, y) \cap [a, b] &= \{b\} \quad \text{and} \\ L(x, y) \cap [a, b] &= \{a\}. \end{aligned}$$

An element $y \in [a, b]$ is called a *strong relative complement of x in $[a, b]$* if

$$U(x, y) = U(b) \quad \text{and} \quad L(x, y) = L(a).$$

An ordered set (S, \leq) is *strongly relatively complemented* if for every interval $[a, b]$ of S , each $x \in [a, b]$ has a strong relative complement in $[a, b]$.

Of course, every strong relative complement of $x \in [a, b]$ is also a weak relative complement of x in $[a, b]$ but not vice versa.

For the sake of brevity, we will write $U_{[a,b]}(x, y)$ or $L_{[a,b]}(x, y)$ instead of $U(x, y) \cap [a, b]$ or $L(x, y) \cap [a, b]$, respectively.

Theorem 1. Let (S, \leq) be a modular ordered set. Let $a, b \in S$, $a \leq b$, and $x \in [a, b]$. Suppose $y \in S$ is a complement of x . The set $U(a, L(y, b))$ has the least element p if and only if the set $L(U(a, y), b)$ has the greatest element p ; in such a case, p is a strong relative complement of x in $[a, b]$.

Proof. Denote by $A = U(a, L(y, b))$ and $B = L(U(a, y), b)$. Since (S, \leq) is modular, we have

$$\begin{aligned} A &= U(a, L(y, b)) = U(L(U(a, y), b)) = U(B), \\ B &= L(U(a, y), b) = L(U(a, L(y, b))) = L(A). \end{aligned}$$

(Let us note that the second line follows by an application of the dual of modular law since modularity is selfdual, see [2], [3].) Hence, if p is the least

element of A , then $A = U(p)$ and $B = L(A) = L(U(p)) = L(p)$, thus p is the greatest element of B . Dually we can show the converse implication. Moreover, the modularity of (S, \leq) yields

$$\begin{aligned}
 U(x, p) &= U(x, L(p)) = U(x, B) = U(x) \cap U(B) = U(x) \cap A = \\
 &= U(x) \cap U(a, L(y, b)) = U(x, a, L(y, b)) = U(x, L(y, b)) = \\
 &= U(L(U(x, y), b)) = U(L(b)) = U(b), \\
 L(x, p) &= L(x, U(p)) = L(x, A) = L(x) \cap L(A) = L(x) \cap B = \\
 &= L(x) \cap L(U(a, y), b) = L(x, U(a, y), b) = L(x, U(a, y)) = \\
 &= L(U(a, L(x, y))) = L(U(a)) = L(a). \quad \blacksquare
 \end{aligned}$$

Example 1. Applying methods of [2], we can check that the set (S, \leq) in Figure 1 is modular.




Figure 1

Of course, $L(U(x, y)) = L(\emptyset) = S$ and $U(L(x, y)) = U(\emptyset) = S$; thus y is a complement of x in (S, \leq) . Further, $U(a, L(y, b)) = U(a, c) = U(p)$. Thus (S, \leq) , a, b, x, y satisfy the assumption of Theorem 1, and hence p is a strong relative complement of x in $[a, b]$.

Example 2. Let (S, \leq) be the ordered set depicted in Figure 2. S is modular and the element y is a complement of x . The set $A = U(a, L(y, b)) = U(a, c) = \{b, d\}$ has not a least element. The set $B = L(U(a, y), b) = L(d, b) = \{a, c\}$ has not a greatest element. It is easy to see that the element x has not a weak relative complement in $[a, b]$.




Figure 2

Although Theorem 1 is a generalization of the well known lattice statement, we can remove the assumption of modularity of (S, \leq) and the complementarity of x to obtain a bit more general result:

Theorem 2. *Let (S, \leq) be an ordered set, let $a, b \in S$, $a \leq b$, and $x \in [a, b]$. If there exists an element $y \in S$ such that:*

- (i) *the set $L(U(a, y), b)$ has the greatest element e and the set $U(a, L(y, b))$ has the least element f ,*
- (ii) *the set $L(U(a, y), x)$ has the greatest element a and the set $U(x, L(y, b))$ has the least element b ,*

then e and f are strong relative complements of x in $[a, b]$.

Proof. Set $A = U(a, L(y, b))$ and $B = L(U(a, y), b)$. By (i), there exist $e, f \in S$ with $A = U(f)$, $B = L(e)$. Prove $f \leq e$:

Since $c \leq b$ for each $c \in L(y, b)$, we have $U(b) \subseteq U(L(y, b))$. However, $a \leq b$ yields $U(b) \subseteq U(a)$, thus $U(b) \subseteq L(a, L(y, b))$, whence

$$(*) \quad L(b) \supseteq L(U(a, L(y, b))).$$

Analogously, $c \leq y$ for each $c \in L(y, b)$ yields $U(y) \subseteq U(L(y, b))$, clearly $U(a, y) \subseteq U(a, L(y, b))$, whence

$$(**) \quad L(U(a, y)) \supseteq L(U(a, L(y, b))).$$

Applying $(*)$ and $(**)$, we conclude

$$L(e) = B = L(U(a, y), b) \supseteq L(U(a, L(y, b))) = L(A) = L(U(f)) = L(f),$$

i.e. $L(e) \supseteq L(f)$ proving $f \leq e$.

Moreover, (i) and (ii) imply

$$\begin{aligned} L(e, x) &= L(e) \cap L(x) = B \cap L(x) = L(U(a, y), b) \cap L(x) = \\ &= L(U(a, y), b, x) = L(U(a, y), x) = L(a), \\ U(f, x) &= U(f) \cap U(x) = A \cap U(x) = U(a, L(y, b)) \cap U(x) = \\ &= U(a, L(y, b), x) = U(L(y, b), x) = U(b). \end{aligned}$$

Further, we obtain

$$\begin{aligned} U(b) &= U(b, L(U(a, y), b)) = U(b, B) = U(b, e) \subseteq U(x, e) \subseteq U(x, f) = U(b), \\ L(a) &= L(a, U(a, L(y, b))) = L(a, A) = L(a, f) \subseteq L(x, f) \subseteq L(x, e) = L(a), \end{aligned}$$

proving $U(x, e) = U(b)$ and $L(x, f) = L(a)$. Thus e and f are strong relative complements of x in $[a, b]$. ■

Example 3. It is easy to see that the ordered set (S, \leq) in Figure 3 is not modular and for x, y, a, b we have $x \in [a, b]$ and

$$\begin{aligned} L(U(a, y), b) &= L(e), \\ U(a, L(y, b)) &= U(f), \\ L(U(a, y), x) &= L(a), \\ U(x, L(y, b)) &= U(b). \end{aligned}$$




Figure 3

Hence, by Theorem 2, e and f are strong relative complements of x in $[a, b]$.

An analogous result is valid also for weak relative complements:

Theorem 3. Let (S, \leq) be an ordered set, let $a, b \in S$, $a \leq b$, and $x \in [a, b]$. If there exists an element $y \in S$ such that:

- (i) the set $L(U(a, y), b)$ has the greatest element e and the set $U(a, L(y, b))$ has the least element f ,
- (ii)* the set $L(U(a, y), x)$ has a maximal element a and the set $U(x, L(y, b))$ has a minimal element b ,

then e and f are weak relative complements of x in $[a, b]$.

Proof. The proof of $f \leq e$ is the same as in that of Theorem 2. Applying (i) and (ii)* we obtain

$$\begin{aligned} L_{[a,b]}(e, x) &= L(e) \cap L(x) \cap [a, b] = B \cap L(x) \cap [a, b] = \\ &= L(U(a, y), b) \cap L(x) \cap [a, b] = L(U(a, y), b, x) \cap [a, b] = \\ &= L(U(a, y), x) \cap [a, b] = L_{[a,b]}(U(a, y), x) = L_{[a,b]}(a) = \{a\} \end{aligned}$$

and dually

$$\begin{aligned} U_{[a,b]}(f, x) &= U(f) \cap U(x) \cap [a, b] = A \cap U(x) \cap [a, b] = \\ &= U(a, L(y, b)) \cap U(x) \cap [a, b] = U(a, L(y, b), x) \cap [a, b] = \\ &= U(L(y, b), x) \cap [a, b] = U_{[a,b]}(L(y, b), x) = U_{[a,b]}(b) = \{b\}. \end{aligned}$$

Since $f \leq e$, we conclude

$$\begin{aligned} U_{[a,b]}(b) &= U(b) \cap [a, b] = U(b, L(U(a, y), b)) \cap [a, b] = U(b, B) \cap [a, b] = \\ &= U(b, e) \cap [a, b] \subseteq U(x, e) \cap [a, b] \subseteq U(x, f) \cap [a, b] = U(b) \cap [a, b] = U_{[a,b]}(b), \end{aligned}$$

whence $U_{[a,b]}(x, e) = U_{[a,b]}(b)$. Dually, it can be shown that $L_{[a,b]}(x, f) = L_{[a,b]}(a)$. We have proved that e and f are weak relative complements of x in $[a, b]$. ■

Example 4. Consider the ordered set (S, \leq) depicted in Figure 4. Although (S, \leq) is not modular, the elements a, b, x, y satisfy (i) and (ii)* of Theorem 3, thus e and f are weak relative complements of x in $[a, b]$. Moreover, the element x has no strong relative complement in $[a, b]$, since $[a, b] = \{a, b, x, e, f\}$ and for the only possible candidates e and f we have

$$\begin{aligned} L(e, x) &= \{a, y\} \neq \{a\} = L(a), \\ U(e, x) &= \{b, h\} \neq \{b\} = U(b), \end{aligned}$$

analogously also for f .




Figure 4

Now, we turn our attention to some aspects of atomicity in relatively complemented ordered sets. In the case of lattices, it is well known that if L is a relatively complemented lattice of finite length and $b \in L$, $b \neq 0$, then $b = \vee A(b)$, where $A(b)$ is the set of all atoms of L less or equal to b (see e.g. [6]). In the case of ordered sets with 0 we investigate whether $U(A(b)) = U(b)$.

An element a of an ordered set (S, \leq) is called an *atom* if either a covers 0 whenever 0 is the least element of (S, \leq) or a is a minimal element of (S, \leq) in the opposite case. For $b \in S$, denote by $A(b)$ the set of all atoms of S below b . (S, \leq) is called *atomic* if for any $b \in S$, $b \neq 0$ (whenever 0 in S exists) there exists an atom $a \in S$ with $a \leq b$. It is almost evident that if (S, \leq) is of a finite length, then (S, \leq) is atomic.

Theorem 4. *Let (S, \leq) be a strongly relatively complemented ordered set of a finite length with 0. If $b \in S$ and $b \neq 0$, then $U(A(b)) = U(b)$.*

Proof. If b is an atom in S , then $A(b) = \{b\}$, and hence $U(A(b)) = U(b)$. Suppose b is not an atom in S and $b \neq 0$. Since (S, \leq) is of a finite length and hence atomic, there exists $p_1 \in A(b)$. Since (S, \leq) is strongly relatively complemented, there exists $c_1 \in [0, b]$ with $U(p_1, c_1) = U(b)$.

(a) If c_1 is an atom of (S, \leq) then $c_1 \leq b$ implies $c_1 \in A(b)$. Denote by $D = A(b) \setminus \{p_1, c_1\}$. Clearly $U(D) \supseteq U(b)$ (since $U(D) = S$ if $D = \emptyset$ and, for $D \neq \emptyset$, $d \leq b$ for each $d \in D$). Then

$$U(A(b)) = U(p_1, c_1) \cap U(D) = U(b) \cap U(D) = U(b).$$

(b) Suppose c_1 is not an atom of (S, \leq) . We can repeat the same consideration for the element c_1 (instead of the element b), i.e. there exists

$p_2 \in A(c_1)$ and $c_2 \in [0, c_1]$ such that $U(p_2, c_2) = U(c_1)$. Since (S, \leq) is of a finite length, we will finish after n steps of this procedure to obtain an element $c_n \in S$ such that $U(p_n, c_n) = U(c_{n-1})$ and $c_n \in A(c_{n-1})$. Denote by $D_n = A(c_{n-1}) \setminus \{p_n, c_n\}$. Evidently, $U(D_n) \supseteq U(c_{n-1})$ and

$$U(A(c_{n-1})) = U(p_n, c_n) \cap U(D_n) = U(c_{n-1}) \cap U(D_n) = U(c_{n-1}).$$

Further, let $D_{n-1} = A(c_{n-2}) \setminus \{p_{n-1}, A(c_{n-1})\}$. Clearly $U(D_{n-1}) \supseteq U(c_{n-2})$ and again

$$\begin{aligned} U(A(c_{n-2})) &= U(p_{n-1}) \cap U(A(c_{n-1})) \cap U(D_{n-1}) = \\ &= U(p_{n-1}, c_{n-1}) \cap U(D_{n-1}) = U(c_{n-2}) \cap U(D_{n-1}) = U(c_{n-2}). \end{aligned}$$

Analogously we proceed to prove $U(A(c_k)) = U(c_k)$ for $k = 1, \dots, n$. For $D_1 = A(b) \setminus \{p_1, A(c_1)\}$ we have $U(D_1) \supseteq U(b)$, thus also

$$\begin{aligned} U(A(b)) &= U(p_1) \cap U(A(c_1)) \cap U(D_1) = \\ &= U(p_1) \cap U(c_1) \cap U(D_1) = U(b) \cap U(D_1) = U(b). \end{aligned}$$

■

Example 5. Let (S, \leq) be the ordered set depicted in Figure 5. Then (S, \leq) is strongly relatively complemented and of a finite length. For the element $b \in S$ we really have $A(b) = \{a, c, d\}$ and $U(A(b)) = U(a, c, d) = U(b)$.




Figure 5

Remark. If (S, \leq) is a strongly relatively complemented ordered set of a finite length without 0, the assertion of Theorem 4 does not hold in general, see, e.g., Figure 6, where $A(b) = \{a, c\}$ but $U(A(b)) = U(a, c) = U(d) = \{d, b\} \neq \{b\} = U(b)$.




Figure 6

Moreover, b is not even a minimal element of $U(A(b))$. On the other hand, we can state the following

Theorem 5. Let (S, \leq) be an ordered set, let $b \in S$ and $b \neq 0$ whenever 0 in S exists. If b covers an atom a and $\text{card}(A(b)) \geq 2$, then b is a minimal element of $U(A(b))$.

Proof. Let $b \neq 0$ covers an atom a and let $\text{card}(A(b)) \geq 2$. Suppose b is not minimal in $U(A(b))$. Then there exists $m \in U(A(b))$ with $m < b$. Since $a \in A(b)$, we conclude $a \leq m < b$. But b covers a , i.e. $a = m$, hence $a \in U(A(b))$. Since $\text{card}(A(b)) \geq 2$, there exists $c \in A(b)$ with $c \neq a$. It is easy to check that c and a are not comparable. However $a \in U(A(b))$ implies $c \leq a$, a contradiction. ■

Example 6. Let (S, \leq) be an ordered set with the diagram depicted in Figure 7.

Then (S, \leq) has not 0 and it is not of a finite length. $A(b) = \{a, c\}$, i.e. $\text{card}(A(b)) = 2$. In accordance with Theorem 5, b is a minimal element in the set $U(A(b)) = U(a, c) = \{b, d_1, d_2, \dots, e_1, e_2, \dots\}$. On the other hand $U(A(b)) \neq \{b\} = U(b)$.




Figure 7

Remark. The condition “ b covers an atom a ” in Theorem 5 is not necessary. For the set (S, \leq) in Figure 8 we have that b is the unique and hence minimal element of $U(A(b))$ but b covers no atom of S .




Figure 8

On the other hand, if $b \neq 0$ (whenever 0 in S exists) and b is not an atom of S , then, if b is minimal in $U(A(b))$, $\text{card}(A(b)) \geq 2$. Namely, if $\text{card}(A(b)) = 0$, then $A(b) = \emptyset$ and $U(A(b)) = U(\emptyset) = S$; thus b is a minimal element of S . Since $b \neq 0$, S has not 0. However, b is not an atom, a contradiction. If $\text{card}(A(b)) = 1$, then $A(b) = \{a\}$ and $a \neq b$, i.e. $U(A(b)) = U(a)$ and b cannot be the minimal element of $U(A(b))$, a contradiction again.

References

- [1] I. Chajda, *Complemented ordered sets*, Arch. Math. (Brno) **28** (1992), 25–34.
- [2] I. Chajda and J. Rachůnek, *Forbidden configurations for distributive and modular ordered sets*, Order **5** (1989), 407–423.
- [3] R. Halaš, *Pseudocomplemented ordered sets*, Arch. Math. (Brno) **29** (1993), 153–160.
- [4] J. Niederle, *Boolean and distributive ordered sets*, Order **12** (1995), 189–210.
- [5] J. Rachůnek and J. Larmerová, *Translations of modular and distributive ordered sets*, Acta Univ. Palacký Olomouc, Fac. Rerum Nat., Math., **31** (1988), 13–23.
- [6] V.N. Salij, *Lattices with Unique Complementations* (Russian), Nauka, Moskva 1984.

Received 21 September 1998

Revised 7 June 1999