

## BALANCED CONGRUENCES\*

IVAN CHAJDA

*Department of Algebra and Geometry*  
*Palacký University of Olomouc*  
*Tomkova 40, CZ-77900 Olomouc, Czech Republic*  
**e-mail:**chajda@risc.upol.cz

AND

GÜNTHER EIGENTHALER

*Institut für Algebra und Computermathematik*  
*Technische Universität Wien*  
*Wiedner Hauptstraße 8–10, A-1040 Wien, Austria*  
**e-mail:**g.eigenthaler@tuwien.ac.at

### Abstract

Let  $\mathcal{V}$  be a variety with two distinct nullary operations 0 and 1. An algebra  $\mathfrak{A} \in \mathcal{V}$  is called balanced if for each  $\Phi, \Psi \in \text{Con}(\mathfrak{A})$ , we have  $[0]\Phi = [0]\Psi$  if and only if  $[1]\Phi = [1]\Psi$ . The variety  $\mathcal{V}$  is called balanced if every  $\mathfrak{A} \in \mathcal{V}$  is balanced. In this paper, balanced varieties are characterized by a Mal'cev condition (Theorem 3). Furthermore, some special results are given for varieties of bounded lattices.

**Keywords:** balanced congruence, balanced algebra, balanced variety, Mal'cev condition.

**2000 Mathematics Subject Classification:** Primary 08A30; Secondary 08B05.

---

\*This paper is a result of the collaboration of the authors within the framework of the “Aktion Österreich-Tschechische Republik” (grant No. 26p2 “Local and global congruence properties”).

## 1. BALANCED CONGRUENCES ON BOUNDED LATTICES

Let  $\mathfrak{L} = (L; \vee, \wedge, 0, 1)$  be a bounded lattice with least element 0 and greatest element 1. We study congruences on  $\mathfrak{L}$  such that the 0-class determines the 1-class and vice versa.

Let  $M \subseteq L$  and  $a, b \in L$ . Denote by  $\Theta(M)$  the least congruence on  $\mathfrak{L}$  containing the relation  $M \times M$  and by  $\Theta(a, b)$  the principal congruence on  $\mathfrak{L}$  generated by  $(a, b)$ , i.e.  $\Theta(a, b) = \Theta(M)$  for  $M = \{a, b\}$ . We will use the following result of G. Grätzer and E. T. Schmidt, see [3]:

**Proposition 1.** *Let  $\mathfrak{L}$  be a distributive lattice and  $a, b, c, d \in L$  with  $c \leq d$ . Then  $a\Theta(c, d)b$  if and only if  $a \wedge c = b \wedge c$  and  $a \vee d = b \vee d$ .*

Here and in the following, we write  $a\Phi b$  instead of  $(a, b) \in \Phi$ , for any  $\Phi \in \text{Con}(\mathfrak{L})$  and  $a, b \in L$ .

From now on, every lattice under consideration is bounded, i.e. it has 0 and 1.

**Definition 1.** Let  $\mathfrak{L}$  be a lattice with 0 and 1 and  $\Phi \in \text{Con}(\mathfrak{L})$ . Put  $I = [0]\Phi$  and  $F = [1]\Phi$ . We say that  $\Phi$  is *balanced* if

$$[0]\Phi = [0]\Theta(F) \text{ and } [1]\Phi = [1]\Theta(I).$$

**Example 1.** By Corollary 2.1 in [2], every complemented lattice is locally regular at 0 (in the sense of [1]), i.e. for every  $\Phi, \Psi \in \text{Con}(\mathfrak{L})$ , if  $[a]\Phi = [a]\Psi$  for some  $a \in L$ , then  $[0]\Phi = [0]\Psi$ . Dually, it is locally regular at 1, i.e.  $[a]\Phi = [a]\Psi$  implies  $[1]\Phi = [1]\Psi$ . Thus every congruence on every complemented lattice is balanced.

**Definition 2.** Let  $\mathfrak{L}$  be a lattice with 0 and 1. We say that  $\mathfrak{L}$  is a *d-lattice* if for each  $a, b, c, d \in L$  the following holds:

$$\begin{aligned} a\Theta(c, 1)0 &\Rightarrow a \wedge c = 0 \\ b\Theta(d, 0)1 &\Rightarrow b \vee d = 1. \end{aligned}$$

**Example 2.** The lattice  $\mathfrak{N}_5$  is a d-lattice. Of course,  $\mathfrak{N}_5$  has just five congruences: the identical one  $\omega$ , the full square  $N_5 \times N_5$  and three nontrivial ones  $\Theta_1, \Theta_2, \Theta_3$  defined by the following partitions (see Figure 1):

$$\begin{aligned} \Theta_1 &\dots \{0, b, c\}, \{a, 1\} \\ \Theta_2 &\dots \{0\}, \{a\}, \{b, c\}, \{1\} \\ \Theta_3 &\dots \{0, a\}, \{b, c, 1\}. \end{aligned}$$




Figure 1

It is an easy exercise to verify our condition for  $\Theta_1, \Theta_2, \Theta_3$ ; for  $\omega$  and  $N_5 \times N_5$  it is trivial (since  $\omega = \Theta(0, 0) = \Theta(1, 1)$  and  $N_5 \times N_5 = \Theta(0, 1)$ ).

**Lemma 1.** *Every bounded distributive lattice is a d-lattice.*

The proof follows immediately by Proposition 1 putting  $b = 0$  and  $d = 1$  in the first case and  $a = 1$  and  $c = 0$  in the second one. ■

**Lemma 2.** *Let  $\mathfrak{L}$  be a bounded lattice and  $\Phi \in \text{Con}(\mathfrak{L})$ . Take  $S = \{a \in L : c \wedge a = 0 \text{ for some } c \in [1]\Phi\}$ . Then  $S \subseteq [0]\Phi$ .*

**Proof.** Let  $a \in S$ . Then  $a \wedge c = 0$  for some  $c \in [1]\Phi$ . Hence,  $c\Phi 1$  thus also  $(c \vee a)\Phi(1 \vee a) = 1$ , i.e.  $c \vee a \in [1]\Phi$ . It yields  $c\Phi(c \vee a)$  and  $0 = (c \wedge a)\Phi((c \vee a) \wedge a) = a$  proving  $a \in [0]\Phi$ . ■

**Theorem 1.** *Let  $\mathfrak{L}$  be a d-lattice and  $\Phi \in \text{Con}(\mathfrak{L})$ . Then  $\Phi$  is balanced if and only if*

$$\begin{aligned} [0]\Phi &= \{a \in L : c \wedge a = 0 \text{ for some } c \in [1]\Phi\} \text{ and} \\ [1]\Phi &= \{b \in L : d \vee b = 1 \text{ for some } d \in [0]\Phi\}. \end{aligned}$$

**Proof.** Let  $\mathfrak{L}$  be a d-lattice and let  $\Phi \in \text{Con}(\mathfrak{L})$  be balanced. Define a set  $S = \{a \in L : c \wedge a = 0 \text{ for some } c \in [1]\Phi\}$ . By Lemma 2 we have  $S \subseteq [0]\Phi$ . Conversely, let  $a \in [0]\Phi$ . Since  $\Phi$  is balanced, we have  $0\Theta(F)a$  for  $F = [1]\Phi$ . Applying the fact that  $\text{Con}(\mathfrak{L})$  is a compactly generated lattice, there exist elements  $c_1, \dots, c_k \in F$  such that  $0[\Theta(1, c_1) \vee \dots \vee \Theta(1, c_k)]a$ . Set  $c = c_1 \wedge \dots \wedge c_k$ . Then  $c \leq c_i \leq 1$  whence  $\Theta(1, c_i) \subseteq \Theta(1, c)$ , i.e.  $0\Theta(1, c)a$ . Since  $\mathfrak{L}$  is a d-lattice, it implies  $0 = c \wedge a$ . Since  $c \in F$ , we have  $a \in S$ , i.e.  $[0]\Phi \subseteq S$ . We have shown  $[0]\Phi = S$ . Dually it can be shown that  $[1]\Phi = \{b \in L : d \vee b = 1 \text{ for some } d \in [0]\Phi\}$ .

Conversely, let  $\Phi \in \text{Con}(\mathfrak{L})$ , and suppose that  $[0]\Phi$ ,  $[1]\Phi$  are described as in Theorem 1. Set  $I = [0]\Phi$ ,  $F = [1]\Phi$  and  $\Psi_1 = \Theta(F)$ ,  $\Psi_2 = \Theta(I)$ . Evidently,  $[1]\Psi_1 = F$  and  $[0]\Psi_2 = I$ .

It gives immediately

$$\begin{aligned} [0]\Phi &= \{a \in L : c \wedge a = 0 \text{ for some } c \in F\} = \\ &= \{a \in L : c \wedge a = 0 \text{ for some } c \in [1]\Psi_1\} \end{aligned}$$

thus  $I \subseteq [0]\Psi_1$  by Lemma 2. However, both  $\Phi$  and  $\Psi_1$  have the 1-class  $F$  and  $\Psi_1$  is the least congruence with this property, i.e.  $\Psi_1 \subseteq \Phi$  whence  $[0]\Psi_1 \subseteq I$ . We have shown that  $[0]\Phi = I = [0]\Theta(F)$ .

Analogously one can prove  $[1]\Phi = [1]\Theta(I)$ , i.e.  $\Phi$  is balanced.  $\blacksquare$

**Example 3.** Consider the distributive lattice  $\mathfrak{L}$  visualized in Figure 2. Consider the congruences  $\Phi_1$  and  $\Phi_2$  determined by the following partitions:

$$\begin{aligned} \Phi_1 \quad & \dots \quad \{0, a\}, \{p, q\}, \{c, 1\} \\ \Phi_2 \quad & \dots \quad \{0, p\}, \{a, q\}, \{c\}, \{1\}. \end{aligned}$$

Then  $\Phi_1$  is balanced since

$$\begin{aligned} [0]\Phi_1 &= \{0, a\} = \{x \in L : c \wedge x = 0\}, \\ [1]\Phi_1 &= \{c, 1\} = \{y \in L : a \vee y = 1\}. \end{aligned}$$

On the contrary,  $\Phi_2$  is not balanced since  $[1]\Phi_2 = \{1\} = [1]\omega$ , whereas  $[0]\Phi_2 \neq [0]\omega$ .




Figure 2

## 2. BALANCED ALGEBRAS WITH TWO NULLARY OPERATIONS

**Definition 3.** Let  $\tau$  be a type containing two distinct nullary operations denoted by 0 and 1 and let  $\mathfrak{A}$  be an algebra of type  $\tau$ . A congruence  $\Phi \in \text{Con}(\mathfrak{A})$  is called *balanced* if

$$[0]\Phi = [0]\Theta(F) \text{ and } [1]\Phi = [1]\Theta(I)$$

where, as in Definition 1,  $I = [0]\Phi, F = [1]\Phi$  and for  $M \subseteq A$ ,  $\Theta(M)$  denotes the least congruence on  $\mathfrak{A}$  containing  $M \times M$ . The algebra  $\mathfrak{A}$  is called *balanced* if each  $\Phi \in \text{Con}(\mathfrak{A})$  is balanced. A variety  $\mathcal{V}$  of type  $\tau$  is called *balanced* if every  $\mathfrak{A} \in \mathcal{V}$  has this property.

**Lemma 3.** *An algebra  $\mathfrak{A}$  with nullary operations 0 and 1 is balanced if and only if for each  $\Phi, \Psi \in \text{Con}(\mathfrak{A})$  the following condition holds:*

$$[0]\Phi = [0]\Psi \text{ if and only if } [1]\Phi = [1]\Psi.$$

**Proof.** Of course, if  $\Phi$  and  $\Psi$  are balanced and  $[0]\Phi = [0]\Psi = I$ , then  $[1]\Phi = [1]\Theta(I) = [1]\Psi$  and vice versa. Conversely, suppose that  $[0]\Phi = [0]\Psi$  if and only if  $[1]\Phi = [1]\Psi$  for each  $\Phi, \Psi \in \text{Con}(\mathfrak{A})$ . Then  $[0]\Phi = I = [0]\Theta(I)$ , thus also  $[1]\Phi = [1]\Theta(I)$  and similarly,  $[0]\Phi = [0]\Theta(F)$  with  $F = [1]\Phi$ , i.e. every congruence on  $\mathfrak{A}$  is balanced. So  $\mathfrak{A}$  is balanced. ■

**Example 4.** The lattice  $\mathfrak{N}_5$  (see Figure 1) is balanced, since for  $\Theta_2$  (of Example 2), we have  $[0]\Theta_2 = [0]\omega$  and  $[1]\Theta_2 = [1]\omega$ . We have  $\Theta_2 \neq \omega$ , so it is a nontrivial example.

The consequence of Example 1 is that every complemented lattice is balanced (this yields another proof that  $\mathfrak{N}_5$  is balanced). However, this condition is not necessary. For example, every simple lattice is balanced but it need not be complemented, see e.g. Figure 3.




Figure 3

**Problem.** If  $\mathfrak{L}$  is a bounded distributive lattice (or a d-lattice), is it true that  $\mathfrak{L}$  is balanced if and only if  $\mathfrak{L}$  is complemented?

**Lemma 4.** *Let  $\mathfrak{A}$  be an algebra with nullary operations 0 and 1. If  $\mathfrak{A}$  is balanced then for each  $\Phi \in \text{Con}(\mathfrak{A})$  the following property holds:*

$$(S) \quad [0]\Phi \text{ is a singleton if and only if } [1]\Phi \text{ is a singleton.}$$

**Proof.** Suppose, e.g., that  $[1]\Phi = \{1\}$ . Then  $[1]\Phi = [1]\omega$  for the identical congruence  $\omega \in \text{Con}(\mathfrak{A})$ , thus, by Lemma 3, also  $[0]\Phi = [0]\omega = \{0\}$ . The converse can be shown analogously. ■

For varieties, the converse statement is also valid:

**Theorem 2.** *Let  $\mathcal{V}$  be a variety of type  $\tau$  containing two distinct nullary operations 0 and 1.  $\mathcal{V}$  is balanced if and only if for each  $\mathfrak{A} \in \mathcal{V}$  and every  $\Phi \in \text{Con}(\mathfrak{A})$  property (S) holds.*

**Proof.** Let  $\mathcal{V}$  satisfy condition (S), let  $\mathfrak{A} \in \mathcal{V}$  and  $\Phi, \Psi \in \text{Con}(\mathfrak{A})$ . Suppose  $[1]\Phi = [1]\Psi$ . Then clearly also  $[1]\Phi = [1]\Phi \wedge \Psi$ , so we can assume  $\Psi \subseteq \Phi$  without loss of generality. Consider the factor algebra  $\mathfrak{A}/\Psi$  and the factor congruence  $\Phi/\Psi \in \text{Con}(\mathfrak{A}/\Psi)$ . Since  $[1]\Psi = [1]\Phi$ , the class of  $\Phi/\Psi$  containing the element  $[1]\Psi \in \mathfrak{A}/\Psi$  is a singleton. Thus, by (S), also the class  $[0]\Psi/\Phi/\Psi$  is a singleton whence  $[0]\Psi = [0]\Phi$ .

Analogously we can show that  $[0]\Phi = [0]\Psi \Rightarrow [1]\Phi = [1]\Psi$ , thus  $\mathfrak{A}$  and hence also  $\mathcal{V}$  is balanced, by Lemma 3. The converse assertion follows directly by Lemma 4. ■

### 3. A CHARACTERIZATION OF BALANCED VARIETIES

The following Theorem characterizes balanced varieties by a Mal'cev condition.

**Theorem 3.** *Let  $\mathcal{V}$  be a variety of type  $\tau$  which contains two distinct nullary operations 0 and 1. The following conditions are equivalent:*

1.  $\mathcal{V}$  is balanced.
2. *There exist – for some  $m, n, k$  and  $h$  – unary terms  $p_1, \dots, p_m, q_1, \dots, q_n$ ,  $(2m+1)$ -ary terms  $r_1, \dots, r_k$  and  $(2n+1)$ -ary terms  $s_1, \dots, s_h$  such that the following identities hold in  $\mathcal{V}$ :*

$$\begin{aligned}
& p_1(0) = \dots = p_m(0) = 1, \quad q_1(1) = \dots = q_n(1) = 0, \text{ and} \\
& x = r_1(p_1(x), \dots, p_m(x), 1, \dots, 1, x), \\
& r_i(1, \dots, 1, p_1(x), \dots, p_m(x), x) = r_{i+1}(p_1(x), \dots, p_m(x), 1, \dots, 1, x), \\
& i = 1, \dots, k-1, \\
& 0 = r_k(1, \dots, 1, p_1(x), \dots, p_m(x), x), \text{ and} \\
& x = s_1(q_1(x), \dots, q_n(x), 0, \dots, 0, x), \\
& s_j(0, \dots, 0, q_1(x), \dots, q_n(x), x) = s_{j+1}(q_1(x), \dots, q_n(x), 0, \dots, 0, x), \\
& j = 1, \dots, h-1, \\
& 1 = s_h(0, \dots, 0, q_1(x), \dots, q_n(x), x).
\end{aligned}$$

3. *There exist – for some  $m, n$  – unary terms  $p_1, \dots, p_m, q_1, \dots, q_n$  and 3-ary terms  $r_1, \dots, r_m, s_1, \dots, s_n$  such that the following identities hold in  $\mathcal{V}$ :*

$$\begin{aligned}
& p_1(0) = \dots = p_m(0) = 1, \quad q_1(1) = \dots = q_n(1) = 0, \text{ and} \\
& x = r_1(p_1(x), 1, x), \\
& r_i(1, p_i(x), x) = r_{i+1}(p_{i+1}(x), 1, x), \quad i = 1, \dots, m-1, \\
& 0 = r_m(1, p_m(x), x), \text{ and} \\
& x = s_1(q_1(x), 0, x), \\
& s_j(0, q_j(x), x) = s_{j+1}(q_{j+1}(x), 0, x), \quad j = 1, \dots, n-1, \\
& 1 = s_n(0, q_n(x), x).
\end{aligned}$$

4. *There exist – for some  $m, n$  – unary terms  $p_1, \dots, p_m, q_1, \dots, q_n$  such that*

$$\begin{aligned}
& [p_1(x) = 1 \& \dots \& p_m(x) = 1] \Leftrightarrow x = 0, \\
& [q_1(x) = 0 \& \dots \& q_n(x) = 0] \Leftrightarrow x = 1.
\end{aligned}$$

**Proof.** (1)  $\Rightarrow$  (2): Let  $\mathfrak{A} = \mathfrak{F}_{\mathcal{V}}(x)$  be a free algebra of  $\mathcal{V}$  with one free generator  $x$ , let  $\Psi = \Theta(x, 0) \in \text{Con}(\mathfrak{A})$  and  $B = [1]\Psi$ . Set  $\Phi = \Theta(B)$ , i.e.,  $\Phi$  is the least congruence on  $\mathfrak{A}$  having the 1-class equal to  $B$ . Then  $[1]\Psi = [1]\Phi$  thus, by (1), also  $[0]\Psi = [0]\Phi$ , i.e.,  $0\Phi x$ . As the lattice  $\text{Con}(\mathfrak{A})$  is compactly generated, there exist  $b_1, \dots, b_m \in B$  such that

$$(*) \quad x[\Theta(1, b_1) \vee \dots \vee \Theta(1, b_m)]0.$$

Since  $b_i \in \mathfrak{F}_{\mathcal{V}}(x)$ , there exist unary terms  $p_i(x)$  such that  $b_i = p_i(x)$  for  $i = 1, \dots, m$ . Since  $b_i \in [1]\Theta(x, 0)$ , it implies immediately

$$p_i(0) = 1 \text{ for } i = 1, \dots, m.$$

Applying the well-known Mal'cev scheme (see [4]) on (\*), we obtain

$$\begin{aligned}
x &= r_1(p_1(x), \dots, p_m(x), 1, \dots, 1, x), \\
r_j(1, \dots, 1, p_1(x), \dots, p_m(x), x) &= r_{j+1}(p_1(x), \dots, p_m(x), 1, \dots, 1, x), \\
0 &= r_k(1, \dots, 1, p_1(x), \dots, p_m(x), x)
\end{aligned}$$

for some  $(2m+1)$ -ary terms  $r_1, \dots, r_k$  and  $j = 1, \dots, k-1$ .

If we set  $\Psi = \Theta(x, 1)$  and  $B = [0]\Psi$ , then the constants 0 and 1 only interchange their roles and we obtain the remaining identities of (2) analogously.

(2)  $\Rightarrow$  (3): For  $i = 2, \dots, m-1$  and  $j = 1, \dots, k$  put

$$\begin{aligned}
r_{j1}(u, v, x) &= r_j(u, p_2(x), \dots, p_m(x), v, 1, \dots, 1, x) \\
r_{ji}(u, v, x) &= r_j(1, \dots, 1, u, p_{i+1}(x), \dots, p_m(x), p_1(x), \dots, p_{i-1}(x), v, 1, \dots, 1, x),
\end{aligned}$$

where there are  $i-1$  nullary operations 1 at the beginning and  $m-i$  nullary operations 1 after  $v$ , and

$$r_{jm}(u, v, x) = r_j(1, \dots, 1, u, p_1(x), \dots, p_{m-1}(x), v, x).$$

Furthermore, put  $p_{ji}(x) = p_i(x)$  for  $i = 1, \dots, m$  and  $j = 1, \dots, k$ .

Then we have for  $i = 1, \dots, m-1$  and  $j = 1, \dots, k$ :

$$\begin{aligned}
&r_{ji}(1, p_{ji}(x), x) = \\
&= r_j(1, \dots, 1, 1, p_{i+1}(x), \dots, p_m(x), p_1(x), \dots, p_{i-1}(x), p_i(x), 1, \dots, 1, x) = \\
&= r_{j,i+1}(p_{j,i+1}(x), 1, x)
\end{aligned}$$

and for  $j = 1, \dots, k-1$ :

$$\begin{aligned}
r_{jm}(1, p_{jm}(x), x) &= r_j(1, \dots, 1, 1, p_1(x), \dots, p_{m-1}(x), p_m(x), x) = \\
&= r_{j+1}(p_1(x), \dots, p_m(x), 1, \dots, 1, x) = \\
&= r_{j+1,1}(p_{j+1,1}(x), 1, x),
\end{aligned}$$

where the second identity follows from (2).

Furthermore we have, again by (2):

$$\begin{aligned}
r_{11}(p_{11}(x), 1, x) &= r_1(p_1(x), p_2(x), \dots, p_m(x), 1, 1, \dots, 1, x) = x, \\
r_{km}(1, p_{km}(x), x) &= r_k(1, \dots, 1, 1, p_1(x), \dots, p_{m-1}(x), p_m(x), x) = 0.
\end{aligned}$$

By writing  $m$  instead of  $km$ ,  $r_1, \dots, r_m$  instead of  $r_{11}, \dots, r_{km}$  and  $p_1, \dots, p_m$  instead of  $p_{11}, \dots, p_{km}$  (both in lexicographic order), we obtain the first part of (3). The second part of (3) can be shown analogously.



(3)  $\Rightarrow$  (4): Take the same terms  $p_i, q_j$  as in (3). Then  $p_i(0) = 1, q_j(1) = 0$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$ .

If we suppose  $p_i(x) = 1$  for  $i = 1, \dots, m$ , then, by (3), we obtain

$$\begin{aligned} x &= r_1(p_1(x), 1, x) = r_1(1, 1, x) = \\ &= r_2(1, 1, x) = \dots = r_k(1, 1, x) = \\ &= r_m(1, p_m(x), x) = 0, \end{aligned}$$

i.e.,  $x = 0$ , thus we have the first implication of (4). The second one can be shown analogously.

(4)  $\Rightarrow$  (1): Let  $\mathfrak{A} \in \mathcal{V}$  and  $\Phi \in \text{Con}(\mathfrak{A})$ . Suppose, e.g.,  $[1]\Phi = \{1\}$  and take  $b \in [0]\Phi$ . Then  $[b]\Phi = [0]\Phi$  and, by (4), we have in the factor algebra  $\mathfrak{A}/\Phi$

$$[p_i(b)]\Phi = p_i([b]\Phi) = p_i([0]\Phi) = [1]\Phi = \{1\}$$

whence  $p_i(b) = 1$  for  $i = 1, \dots, m$ . Applying (4) again, we conclude  $b = 0$ , thus  $[0]\Phi = \{0\}$ . Analogously we can show

$$[0]\Phi = \{0\} \Rightarrow [1]\Phi = \{1\}.$$

By Theorem 2,  $\mathcal{V}$  is balanced. ■

**Remark.** The equivalence of (2) and (3) can also be proved directly since (3)  $\Rightarrow$  (2) can be seen as follows:

Take the  $r_i$  in (3) and define the  $(2m+1)$ -ary terms  $\bar{r}_i$  by

$$\bar{r}_i(x_1, \dots, x_m, y_1, \dots, y_m, x) = r_i(x_i, y_i, x), i = 1, \dots, m,$$

then we have

$$\begin{aligned} \bar{r}_i(p_1(x), \dots, p_m(x), 1, \dots, 1, x) &= r_i(p_i(x), 1, x) \text{ and} \\ \bar{r}_i(1, \dots, 1, p_1(x), \dots, p_m(x), x) &= r_i(1, p_i(x), x), \end{aligned}$$

and we obtain the identities of (2) with  $m = k$  and  $\bar{r}_i$  instead of  $r_i$ . Similarly, we obtain the identities of (2) with  $n = h$  and some  $\bar{s}_i$  instead of  $s_i$ .

**Example 5.** The variety of all ortholattices  $(L; \vee, \wedge, ^\perp, 0, 1)$  is balanced. We can take  $m = 1 = n$  and  $p_1(x) = q_1(x) = x^\perp$ . Then  $p_1(x) = 1$  if and only if  $x = 0$  and  $q_1(x) = 0$  if and only if  $x = 1$ .

**Example 6.** Every variety of double  $p$ -algebras is balanced (a double  $p$ -algebra is an algebra  $(L; \vee, \wedge, *, ^+, 0, 1)$  of type  $(2, 2, 1, 1, 0, 0)$ , where  $(L; \vee, \wedge, 0, 1)$  is a bounded lattice,  $a^*$  is a pseudocomplement of  $a \in L$  and  $a^+$  is a dual pseudocomplement). We can take  $m = 1 = n$  and  $p_1(x) = x^*$ ,  $q_1(x) = x^+$ . Of course,  $0^* = 1$ ,  $1^+ = 0$ , and  $x^* = 1 \Rightarrow x = 0$ ,  $x^+ = 0 \Rightarrow x = 1$ .

## REFERENCES

- [1] I. Chajda, *Locally regular varieties*, Acta Sci. Math. (Szeged) **64** (1998), 431–435.
- [2] I. Chajda and G. Eigenthaler, *A remark on congruence kernels in complemented lattices and pseudocomplemented semilattices*, Contributions to General Algebra **11** (1999), 55–58.
- [3] G. Grätzer and E.T. Schmidt, *Ideals and congruence relations in lattices*, Acta Math. Sci. Hungar. **9** (1958), 137–175.
- [4] A.I. Mal'cev, *On the general theory of algebraic systems* (Russian), Mat. Sb. **35** (1954), 3–20.

Received 27 March 2000

Revised 2 August 2000