

ON ABSOLUTE RETRACTS AND ABSOLUTE CONVEX RETRACTS IN SOME CLASSES OF ℓ -GROUPS

JÁN JAKUBÍK

Matematický ústav SAV,
Grešáková 6, 040 01 Košice, Slovakia
email: kstefan@saske.sk

Abstract

By dealing with absolute retracts of ℓ -groups we use a definition analogous to that applied by Halmos for the case of Boolean algebras. The main results of the present paper concern absolute convex retracts in the class of all archimedean ℓ -groups and in the class of all complete ℓ -groups.

Keywords: ℓ -group, absolute retract, absolute convex retract, archimedean ℓ -group, complete ℓ -group, orthogonal completeness.

2000 Mathematics Subject Classification: 06F20, 06F15.

1. INTRODUCTION

Retracts of abelian ℓ -groups and of abelian cyclically ordered groups were investigated in [6], [7], [8].

Suppose that \mathcal{C} is a class of algebras. An algebra $A \in \mathcal{C}$ is called an absolute retract in \mathcal{C} if, whenever $B \in \mathcal{C}$ and A is a subalgebra of B , then A is a retract of B (i.e., there is a homomorphism h of B onto A such that $h(a) = a$ for each $a \in A$). Cf., e.g., Halmos [3].

Further, let \mathcal{C} be a class of ℓ -groups. An element $A \in \mathcal{C}$ will be called an absolute convex retract in \mathcal{C} if, whenever $B \in \mathcal{C}$ and A is a convex ℓ -subgroup of B , then A is a retract of B .

Let \mathcal{G} and Arch be the class of all ℓ -groups, or the class of all archimedean ℓ -groups, respectively.

It is easy to verify (cf. Section 2 below) that for $A \in \mathcal{G}$ the following conditions are equivalent:

- (i) A is an absolute retract in \mathcal{G} ;
- (ii) A is an absolute convex retract in \mathcal{G} ;
- (iii) $A = \{0\}$.

In this note we prove

- (α) Let A be an absolute retract in the class Arch. Then the ℓ -group A is divisible, complete and orthogonally complete.

By applying a result of [5] we obtain

- (β) Let $A \in \text{Arch}$ and suppose that the ℓ -group A is complete and orthogonally complete. Then A is an absolute convex retract in the class Arch.

The question whether the implication in (α) (or in (β), respectively) can be reversed remains open.

Let us denote by

Compl - the class of all complete ℓ -groups;

Compl* - the class of all ℓ -groups which are complete and orthogonally complete.

- (γ) Let $A \in \text{Compl}$. Then the following conditions are equivalent:

- (i) A is orthogonally complete.
- (ii) A is an absolute convex retract in the class Compl.

As a corollary we obtain that each ℓ -group belonging to Compl* is an absolute convex retract in the class Compl*.

We prove that if the class $\mathcal{C} \subseteq \mathcal{G}$ is closed with respect to direct products and if A_i ($i \in I$) are absolute (convex) retracts in \mathcal{C} , then their direct product $\prod_{i \in I} A_i$ is also an absolute (convex) retract in \mathcal{C} .

2. PRELIMINARIES

For ℓ -groups we apply the notation as in Conrad [1]. Hence, in particular, the group operation in an ℓ -group is written additively.

We recall some relevant notions. Let G be an ℓ -group. G is *divisible* if for each $a \in G$ and each positive integer n there is $x \in G$ with $nx = a$. A system $\emptyset \neq \{x_i\}_{i \in I} \subseteq G^+$ is called *orthogonal* (or *disjoint*) if $x_{i(1)} \wedge x_{i(2)} = 0$ whenever $i(1)$ and $i(2)$ are distinct elements of I . If each orthogonal subset of G possesses the supremum in G then G is said to be *orthogonally complete*. G is *complete* if each nonempty bounded subset of G has the supremum and the infimum in G .

G is *archimedean* if, whenever $0 < x \in G$ and $y \in G$, then there is a positive integer n such that $nx \not\leq y$. For each archimedean ℓ -group G there exists a complete ℓ -group $D(G)$ (the Dedekind completion of G) such that

- (i) G is a closed ℓ -subgroup of $D(G)$;
- (ii) for each $x \in D(G)$ there are subsets $\{y_i\}_{i \in I}$ and $\{z_j\}_{j \in J}$ of G such that the relations

$$\sup\{x_i\}_{i \in I} = x = \inf\{z_j\}_{j \in J}$$

are valid in $D(G)$.

Let G_1 be a linearly ordered group and let G_2 be an ℓ -group. The symbol $G_1 \circ G_2$ denotes the lexicographic product of G_1 and G_2 . The elements of $G_1 \circ G_2$ are pairs (g_1, g_2) with $g_1 \in G_1$ and $g_2 \in G_2$. For each $g_2 \in G_2$, the pair $(0, g_2)$ will be identified with the element g_2 of G_2 . Then G_2 is a convex ℓ -subgroup of $G_1 \circ G_2$.

Lemma 2.1. *Let A be an ℓ -group, $A \neq \{0\}$, and let G_1 be a linearly ordered group, $G_1 \neq \{0\}$. Put $B = G_1 \circ A$. Then A fails to be a retract of B .*

Proof. By way of contradiction, suppose that A is a retract of B . Let h be the corresponding retract homomorphism of B onto A ; i.e., $h(a) = a$ for each $a \in A$. There exists $g_1 \in G_1$ with $g_1 > 0$. Denote $(g_1, 0) = b$, $h(b) = a$. Further, there exists $a_1 \in A$ with $a_1 > a$. We have $a_1 < b$, whence $h(a_1) \leq h(b)$, thus $a_1 \leq a$, which is a contradiction. ■

Let us denote by \mathcal{A} the class of all abelian lattice ordered groups. If A, G_1 and B are as in Lemma 2.1 and $A, G_1 \in \mathcal{A}$, then also B belongs to \mathcal{A} . Thus Lemma 2.1 yields

Proposition 2.2. *Let $\mathcal{C} \in \{\mathcal{G}, \mathcal{A}\}$ and let A be an absolute retract (or an absolute convex retract, respectively) in the class \mathcal{C} . Then $A = \{0\}$. ■*

It is obvious that $\{0\}$ is an absolute (convex) retract in both the classes \mathcal{G} and \mathcal{A} .

Let us remark that if $G_1, B \in \mathcal{G}$ and if G_1 is a retract of B , then G_1 need not be a convex ℓ -subgroup of B . This is verified by the following example:

Let G_1 be a linearly ordered group, $G_1 \neq \{0\}$. Further, let $G_2 \in \mathcal{G}$, $G_2 \neq \{0\}$. Put $B = G_1 \circ G_2$. If $g_1 \in G_1$, then the element $(g_1, 0)$ of B will be identified with the element g_1 of G_1 . Thus G_1 turns out to be an ℓ -subgroup of B which is not a convex subset of B . For each $(g_1, g_2) \in B$ we put $h((g_1, g_2)) = g_1$. Then h is a homomorphism of B onto G_1 such that $h(g_1) = g_1$ for each $g_1 \in G_1$. Hence G_1 is a retract of B .

3. PROOFS OF (α) , (β) AND (γ)

In this section we assume that A is an archimedean ℓ -group. Hence A is abelian.

It is well-known that there exists the divisible hull A^d of A . Thus

- (i) A^d is a divisible ℓ -group;
- (ii) A is an ℓ -subgroup of A^d ;
- (iii) if $g \in A^d$, then there are $a \in A$, a positive integer n and an integer m such that $ng = ma$.

Lemma 3.1. *Assume that A is an absolute retract in the class Arch. Then the ℓ -group A is divisible.*

Proof. By way of contradiction, suppose that A fails to be divisible. Thus there are $a_1 \in A$ and $n \in \mathbb{N}$ such that there is no x in A with $nx = a_1$.

Put $B = A^d$. In view of the assumption, A is a retract of B ; let h be the corresponding retract homomorphism.

There exists $b \in B$ with $nb = a_1$. Then $b \notin A$. Denote $h(b) = a$. We have

$$a_1 = h(a_1) = h(nb) = nh(b) = na,$$

which is a contradiction. ■

Lemma 3.2. *Assume that A is an absolute retract in the class Arch. Then A is a complete ℓ -group.*

Proof. By way of contradiction, suppose that A fails to be complete. Put $B = D(A)$. Then A is an ℓ -subgroup of B and $A \neq B$. Thus there is $b \in B$ such that b does not belong to A .

In view of the assumption, A is a retract of B ; let h be the corresponding retract homomorphism. Put $h(b) = a$.

There exists a subset $\{a_i\}_{i \in I}$ of A such that the relation

$$b = \bigvee_{i \in I} a_i$$

is valid in B . Hence $a_i \leq b$ for each $i \in I$. This yields

$$a_i = h(a_i) \leq h(b) = a$$

for each $i \in I$. Thus $b \leq a$.

At the same time, there exists a subset $\{a'_j\}_{j \in J}$ of A such that the relation

$$b = \bigwedge_{j \in J} a'_j$$

holds in B . Hence $b \leq a'_j$ for each $j \in J$, thus by applying the homomorphism h we obtain that $a \leq a'_j$ for each $j \in J$. Therefore $a \leq b$. Summarizing, $a = b$ and we arrived at a contradiction. ■

Lemma 3.3. *Suppose that H is a complete ℓ -group. Then there exists an ℓ -group K such that*

- (i) H is a convex ℓ -subgroup of K ;
- (ii) K is complete and orthogonally complete;
- (iii) for each $0 < k \in K$ there exists a disjoint subset $\{x_i\}_{i \in I}$ of H such that the relation

$$k = \bigvee_{i \in I} x_i$$

is valid in K .

Proof. This is a consequence of results of [5]. ■

Lemma 3.4. *Assume that A is an absolute retract in the class Arch. Then the ℓ -group A is orthogonally complete.*

Proof. In view of Lemma 3.2, A is complete. Put $A = H$ and let K be as in Lemma 3.3. According to the assumption, A is a retract of K . Let h be the corresponding retract homomorphism.

Let $0 < k \in K$ and let $\{x_i\}_{i \in I}$ be as in Lemma 3.3. Put $h(k) = a$. Then $a \geq h(x_i) = x_i$ for each $i \in I$, whence $k \leq a$. Thus the condition (i) of Lemma 3.3 yields that $k \in A$. Hence $K^+ \subseteq A$ and then $K \subseteq A$. Therefore $K = A$ and so A is orthogonally complete. ■

From Lemmas 3.1, 3.2 and 3.4 we conclude that (α) is valid.

Let $G_1, G_2 \in \mathcal{G}$; their direct product is denoted by $G_1 \times G_2$. If $g_1 \in G_1$, then the element $(g_1, 0)$ of $G_1 \times G_2$ will be identified with g_1 . Similarly, for $g_2 \in G_2$, the element $(0, g_2)$ of $G_1 \times G_2$ will be identified with g_2 . Under this identification, both G_1 and G_2 are convex ℓ -subgroups of $G_1 \times G_2$.

Definition 3.5. (Cf. [2].) Let $G_1 \in \text{Arch}$. We say that G_1 has the *splitting property* if, whenever $H \in \text{Arch}$ and G_1 is a convex ℓ -subgroup of H , then G_1 is a direct factor of H .

Proposition 3.6. (Cf. [4].) Let $G_1 \in \text{Arch}$. Then the following conditions are equivalent:

- (i) G_1 has the splitting property.
- (ii) The ℓ -group G_1 is complete and orthogonally complete.

Lemma 3.7. *Let $H \in \mathcal{G}$ and let G_1 be a direct factor of H . Then G_1 is a retract of H .*

Proof. There exists $G_2 \in \mathcal{G}$ such that $H = G_1 \times G_2$. For $(g_1, g_2) \in H$ we put $h((g_1, g_2)) = g_1$. Then h is a retract homomorphism of H onto G_1 . ■

Proof of (β) . Let $A, B \in \text{Arch}$ and suppose that A is a convex ℓ -subgroup of B . Further, suppose that A is complete and orthogonally complete. In view of Proposition 3.6, A is a direct factor of B . Hence according to Lemma 3.7, A is a retract of B . Therefore A is an absolute convex retract in the class Arch. ■

Lemma 3.8. *Let $A \in \text{Compl}$. Suppose that A is an absolute convex retract in the class Compl . Then A is orthogonally complete.*

Proof. Put $H = A$ and let K be as in Lemma 3.3. In view of Lemma 3.3 (i), A is a convex ℓ -subgroup of K . Hence according to the assumption, A is a retract of K . Now it suffices to apply the same method as in the proof of Lemma 3.4. ■

Lemma 3.9. *Let $A \in \text{Compl}$. Suppose that A is orthogonally complete. Then A is an absolute convex retract in the class Compl .*

Proof. In view of (β) , A is an absolute convex retract in the class Arch . It is well-known that the class Compl is a subclass of Arch . Hence A is an absolute convex retract in the class Compl . ■

From Lemmas 3.8 and 3.9 we conclude that (γ) holds.

Corollary 3.10. *Let $A \in \text{Compl}^*$. Then A is an absolute convex retract in the class Compl^* .* ■

4. DIRECT PRODUCTS

Let A_i ($i \in I$) be ℓ -groups; consider their direct product

$$(1) \quad A = \prod_{i \in I} A_i.$$

Without loss of generality we can suppose that $A_{i(1)} \cap A_{i(2)} = \{0\}$ whenever $i(1)$ and $i(2)$ are distinct elements of I . For $a \in A$ and $i \in I$, we denote by a_i or by $a(A_i)$ the component of a in the direct factor A_i . Let $i \in I$. Put

$$A'_i = \{a \in A : a_i = 0\}.$$

Then we have

$$(2) \quad A = A_i \times A'_i,$$

$$A'_i = \prod_{j \in I \setminus \{i\}} A_j.$$

Let $i(0) \in I$ and $a^{i(0)} \in A_{i(1)}$. There exists $a \in A$ such that

$$a_i = \begin{cases} a^{i(0)} & \text{if } i = i(0), \\ 0 & \text{otherwise.} \end{cases}$$

Then the element a of A will be identified with the element $a^{i(0)}$ of $A_{i(0)}$. Under this identification, each A_i turns out to be a convex ℓ -subgroup of A .

Lemma 4.1. *Let B be an ℓ -group and let A be an ℓ -subgroup of B . Suppose that (1) is valid. Let i be a fixed element of I and assume that A_i is a retract of B ; the corresponding retract homomorphism will be denoted by h_i . Then for each $a \in A$ the relation*

$$h_i(a) = a_i$$

is valid.

Proof. a) At first let $0 \leq a' \in A'_i$ and $0 \leq a^i \in A_i$. Then $a' \wedge a^i = 0$, thus

$$0 = h_i(a') \wedge h_i(a^i) = h_i(a') \wedge a^i.$$

Since this is valid for each $a^i \in A_i$ and $h_i(a') \in A_i$ we conclude that $h_i(a') = 0$. Then $h_i(-a') = 0$ as well and this yields that $h_i(a'') = 0$ for each $a'' \in A'_i$.

b) Let $a \in A$. In view of (2) we have

$$a = a_i + a(A'_i).$$

Thus

$$h_i(a) = h_i(a_i) + h_i(a(A'_i)).$$

According to a), $h_i(a(A'_i)) = 0$. Thus $h_i(a) = a_i$. ■

Lemma 4.2. *Let B be an ℓ -group and let A be an ℓ -subgroup of B . Suppose that (1) is valid and that for each $i \in I$, A_i is a retract of B ; the corresponding retract homomorphism will be denoted by h_i . For $b \in B$ we put*

$$h(b) = b^1 \in A,$$

where $b_i^1 = h_i(b)$ for each $i \in I$. Then

- (i) h is a homomorphism of B into A ;
- (ii) $h(a) = a$ for each $a \in A$.

Proof. The definition of h and the relation (1) immediately yield that (i) is valid. Let $a \in A$ and $i \in I$. Put $h(a) = a^1$. We have

$$a = a_i + a(A'_i),$$

thus by applying (i),

$$h(a) = h(a_i) + h(a(A'_i)),$$

$$a_i^1 = h_i(a_i) + h_i(a(A'_i)).$$

Since $h_i(a_i) = a_i$ and because $(a(A'_i))_i = 0$, according to Lemma 4.1, we obtain

$$a_i^1 = a_i \quad \text{for each } i \in I,$$

thus $a^1 = a$. ■

Corollary 4.3. *Let the assumptions of Lemma 4.2 be valid. Then A is a retract of B .* ■

From Corollary 4.3 we immediately conclude

Proposition 4.4. *Assume that \mathcal{C} is a class of ℓ -groups which is closed with respect to direct products. Let A_i ($i \in I$) be absolute retracts in \mathcal{C} and let (1) be valid. Then A is an absolute retract in \mathcal{C} .* ■

Proposition 4.5. *Assume that \mathcal{C} is a class of ℓ -groups which is closed with respect to direct products. Let A_i ($i \in I$) be absolute convex retracts in \mathcal{C} and let (1) be valid. Then A is an absolute convex retract in \mathcal{C} .*

Proof. Let $B \in \mathcal{C}$ and suppose that A is a convex ℓ -subgroup of B . Then all A_i are convex ℓ -subgroups of B . Hence in view of the assumption, all A_i are retracts of B . Thus according to Corollary 4.3, A is a retract of B . Therefore A is an absolute convex retract in the class \mathcal{C} . ■

5. AN EXAMPLE

The assertions of the following two lemmas are easy to verify; the proofs will be omitted.

Lemma 5.1. *Let A be an ℓ -group which is complete and divisible. Then*

- (i) *we can define (in a unique way) a multiplication of elements of A with reals such that A turns out to be a vector lattice;*
- (ii) *if $r > 0$ is a real, $0 < a \in A$, $X = \{q_1 \in Q : 0 < q_1 \leq r\}$, $Y = \{q_2 \in R : r \leq q_2\}$, then the relations*

$$\sup(q_1 a) = ra = \inf(q_2 a)$$

are valid in A ;

- (iii) *if A_1 is an ℓ -subgroup of A such that A_1 is complete and divisible, and $a_1 \in A$, then for each real r the multiplication ra_1 in A_1 gives the same result as the multiplication ra_1 in A . ■*

Lemma 5.2. *Let A be as in Lemma 5.1 and suppose that $A = \prod_{i \in I} A_i$. Then all A_i are complete and divisible; moreover, for each real r , each $a \in A$ and each $i \in I$ we have*

$$(ra)_i = ra_i. \quad \blacksquare$$

Let R be the additive group of all reals with the natural linear order. We denote by $\mathcal{C}_{\mathcal{R}}$ the class of all lattice ordered groups which can be expressed as direct products of ℓ -groups isomorphic to R .

We remark that if $B \in \mathcal{C}_{\mathcal{R}}$ and if A is an ℓ -subgroup of B which is isomorphic to R , then A need not be a convex ℓ -subgroup of B . In fact, suppose that

$$B = \prod_{i \in I} B_i,$$

where each B_i is isomorphic to R ; let φ_i be an isomorphism of R onto B_i . For each $r \in R$ put

$$\varphi(r) = (\dots, \varphi_i(r), \dots)_{i \in I},$$

$$A = \varphi(R).$$

A is an ℓ -subgroup of B ; if I has more than one element, then A fails to be convex in B .

Let B be as above; suppose that A is an ℓ -group isomorphic to R and that A is an ℓ -subgroup of B . Let $0 < a \in A$. Then $a_i = a(B_i) \geq 0$ for each $i \in I$ and there exists $i(0) \in I$ with $a_{i(0)} > 0$. Thus, in view of Lemma 5.1, we have $(ra)_{i(0)} > 0$ for each $r \in R$ with $r \neq 0$. Further, for each $a_1 \in A$ there exists a uniquely determined element $r \in R$ with $a_1 = ra$. This yields that the mapping

$$\varphi_{i(0)} : a_1 \mapsto (a_1)_{i(0)}$$

is an isomorphism of A into $B_{i(0)}$.

Let $b \in B_{i(0)}$. There exists a unique $r \in R$ such that

$$b = ra_{i(0)}.$$

Then, in view of Lemma 5.2, $b = (ra)_{i(0)}$ and hence the mapping $\varphi_{i(0)}$ is an isomorphism of A onto $B_{i(0)}$.

For each $b \in B$ we put

$$h(b) = \varphi_{i(0)}^{-1} (b_{i(0)}).$$

Then h is a homomorphism of B into A . For $a_1 \in A$ the definition of $\varphi_{i(0)}$ yields

$$h(a_1) = a_1.$$

Thus we obtain

Lemma 5.3. *Let $B \in \mathcal{C}_{\mathcal{R}}$ and let A be an ℓ -subgroup of B such that A is isomorphic to R . Then A is a retract of B .* ■

Corollary 5.4. *Let A be an ℓ -group isomorphic to R . Then A is an absolute retract in the class $\mathcal{C}_{\mathcal{R}}$.* ■

From Lemma 5.4 and Corollary 4.5 we conclude

Proposition 5.5. *Each element of $\mathcal{C}_{\mathcal{R}}$ is an absolute retract in the class $\mathcal{C}_{\mathcal{R}}$.* ■

REFERENCES

- [1] P. Conrad, *Lattice Ordered Groups*, Tulane University, New Orleans, LA, 1970.
- [2] P. Conrad, *The essential closure of an archimedean lattice-ordered group*, Duke Math. J. **38** (1971), 151–160.
- [3] P. Halmos, *Injective and projective Boolean algebras*, p. 114–122 in: “*Lattice Theory*” (R.P. Dilworth, editor), Amer. Math. Soc., Providence, RI, 1961.
- [4] J. Jakubík, *Splitting property of lattice ordered groups*, Czechoslovak Math. J. **24** (1974), 257–269.
- [5] J. Jakubík, *Orthogonal hull of a strongly projectable lattice ordered group*, Czechoslovak Math. J. **28** (1978), 484–504.
- [6] J. Jakubík, *Retracts of abelian cyclically ordered groups*, Archivum Math. (Brno) **25** (1989), 13–18.
- [7] J. Jakubík, *Retracts of abelian lattice ordered groups*, Czechoslovak Math. J. **39** (1989), 477–489.
- [8] J. Jakubík, *Retract varieties of lattice ordered groups*, Czechoslovak Math. J. **40** (1990), 104–112.

Received 21 October 2002