

EXTENSION OF CLASSICAL SEQUENCES TO NEGATIVE INTEGERS

BENALI BENZAGHOU

Faculté de Mathématiques
USTHB BP 32 El-Alia,
16111 Bab-Ezzouar, Alger

e-mail: benrect@wissal.dz

AND

DANIEL BARSKY

Université Paris 13, Institut Galilée,
LAGA UMR 7539, 99 Av. J.-B.
Clément 93430 Villetaneuse, France

e-mail: barsky@math.univ-paris13.fr

Abstract

We give a method to extend Bell exponential polynomials to negative indices. This generalizes many results of this type such as the extension to negative indices of Stirling numbers or of Bernoulli numbers.

Keywords: Stirling numbers, Bell polynomials.

2000 Mathematics Subject Classification: 11B73.

1. INTRODUCTION

Several classical sequences have a "natural" extension to negative indices which preserves algebraic relations. For example, the binomial polynomials

$$\binom{x}{k} = \frac{x(x-1)\dots(x-k+1)}{k!}$$

allows to define the binomial coefficients $\binom{n}{k}$ for $n \in \mathbb{Z}$, $k \in \mathbb{N}$.

The sequence $(x)_n = x(x-1)\dots(x-n+1)$, $n \in \mathbb{N}$, is extended to negative integers by

$$(x)_{-n} = \frac{1}{(x+1)\dots(x+n)}$$

so that the relation

$$(x)_n(x-n)_m = (x)_{n+m}$$

remains valid for $n \in \mathbb{Z}$ and $m \in \mathbb{Z}$.

The factorial sequence $\gamma(n) = n!$ is classically extended in [3] by

$$\gamma(-n) = \frac{(-1)^{n-1}}{(n-1)!}, \quad n > 0.$$

In [1], extensions of the Stirling numbers of the second kind, $S(n, k)$, and of the first kind, $s(n, k)$, are obtained for negative n . We remark that Stirling numbers are values of Bell exponential polynomials, $B_{n,k}(a_1, a_2, \dots)$, $n, k \in \mathbb{N}$, on particular sequences. We give an extension of the Bell polynomials for $n, k \in \mathbb{Z}$. This allows us to recover Branson's result and much more. We thank the referee for his remarks.

2. NOTATIONS AND DEFINITIONS

C is a commutative field, of characteristic zero. For a sequence $u : \mathbb{Z} \rightarrow C$, let us note:

$$\text{supp } u = \{n, u(n) \neq 0\},$$

$$\text{ord } u = \inf \text{supp } u,$$

$$s(C) = \{u, \text{ord } u > -\infty\},$$

$$s_0(C) = \{u, \text{ord } u \geq 0\},$$

e_k the sequence defined by $e_k(n) = \delta_{n,k}$, $k \in \mathbb{Z}$.

For $u \in s_0(C)$, let us denote:

$$(1) \quad g_u(X) = \sum_{n=0}^{\infty} u(n) \frac{X^n}{n!}$$

the associated Hurwitz series (or exponential) to u .

For $u \in s_0(C)$ and $v \in s_0(C)$, the product $g_u(X) \cdot g_v(X) = g_\omega(X)$ defines the Hurwitz product $\omega = u \mathbin{\boxtimes} v$ of sequences u and v , and

$$(2) \quad (u \mathbin{\boxtimes} v)(n) = \sum_{j=0}^n \binom{n}{j} u(j) v(n-j).$$

Let us denote by $\mathcal{A} = \mathcal{A}(C)$ the Hurwitz algebra of sequences of $s_0(C)$ provided with the usual addition and Hurwitz product. The order, ord , is a valuation on \mathcal{A} .

Let us denote T the shift operator on \mathcal{A} :

$$(3) \quad (Tu)(n) = u(n+1)$$

and q the operator of multiplication by n :

$$(4) \quad (qu)(n) = nu(n).$$

Then

$$(5) \quad g_{Tu}(X) = \frac{d}{dX} g_u(X),$$

$$(6) \quad g_{qu}(X) = X \frac{d}{dX} g_u(X)$$

where $\frac{d}{dX}$ stands for the operator of formal differentiation.

Let us define for $k \in \mathbb{Z}$, $g_{e_k}(X) = \frac{X^k}{\gamma(k)}$. If we impose the validity of (5) and $\gamma(-1) = 1$, we obtain

$$(7) \quad \gamma(n) = \begin{cases} n! & \text{for } n \geq 0 \\ \frac{(-1)^{-n-1}}{(-n-1)!} & \text{for } n < 0 \end{cases}$$

what allows us to define the Hurwitz series $\sum_n u(n) \frac{X^n}{\gamma(n)}$ of a sequence u of finite order (positive or negative), and to define the Hurwitz product of two sequences u and v of $s(C)$

$$(8) \quad (u \text{ III } v)(n) = \sum_{i+j=n} \frac{\gamma(n)}{\gamma(i)\gamma(j)} u(i)v(j),$$

actually $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right] = \frac{\gamma(n)}{\gamma(k)\gamma(n-k)}$, $n \in \mathbb{Z}$, $k \in \mathbb{Z}$ is the Roman coefficient [3].

$s(C)$, provided with the generalized Hurwitz product (8) is the fraction fields of the ring \mathcal{A} .

Let u be a sequence of strictly positive order; the composition of series $(g_v \circ g_u)(X) = g_{\bar{w}}(X)$ allows to define the composition of sequences, $\bar{w} = v \circ u$.

For $k \in \mathbb{N}$

$$(9) \quad (e_k \circ u)(n) = \mathcal{B}_{n,k}(u)$$

is Bell partial exponential polynomial [2]. It is a polynomial in $u(1), u(2), \dots, u(n), \dots$ with coefficients in \mathbb{Z} .

For $v \in s_0(C)$,

$$(v \circ u)(n) = \sum_{k=1}^n \mathcal{B}_{n,k}(u)v(k).$$

Proposition 2.1. *The set Ω of sequences of order one is a group for the composition. The inverse \bar{u} of u corresponds to the series $g_{\bar{u}}(X)$ reciprocal of the series $g_u(X)$.*

Examples 2.2. Let “ a ” be the sequence defined by $g_a(X) = e^X - 1$; then $g_{\bar{a}}(X) = \log(1 + X)$ then

$$\begin{cases} \mathcal{B}_{n,k}(a) &= S(n, k) \\ \mathcal{B}_{n,k}(\bar{a}) &= s(n, k) \end{cases}$$

are the Stirling numbers.

Let $(t)_q$ be the sequence $(t)_q(n) = t(t-1)\dots(t-n+1)$ and t^q be the sequence $t^q(n) = t^n$. Then

$$t^q = (t)_q \circ a,$$

$$(t)_q = t^q \circ \bar{a}$$

$Y_q(u, t) = t^q \circ u$ is the sequence of Bell exponential polynomials [2] and

$$Y_n(a, t) = \sum_{k=1}^n S(n, k)t^k = P_n(t)$$

is the n th Bell polynomial.

Remark 2.3. By application of the operators T and q (they are derivations in the Hurwitz algebra $\mathcal{A}(C)$), we can obtain various classical relations on the Bell exponential polynomials and the Stirling numbers.

3. EXTENSION OF BELL PARTIAL EXPONENTIAL POLYNOMIALS

Let u be a sequence of order one and k a rational integer; let us define for $k \in \mathbb{N}$

$$g_{(e_k \circ u)}(X) = \begin{cases} \frac{g_u^k(X)}{\gamma(k)} \\ \frac{1}{\gamma(-k)X^k} \left[\frac{X}{g_u(X)} \right]^k \end{cases}$$

and so

$$g_{(e_{-k} \circ u)}(X) = \sum_n \mathcal{B}_{n,-k}(u) \frac{X^n}{\gamma(n)}.$$

Let us recall the definition of the generalized Bernoulli numbers:

$$\left[\frac{X}{g_u(X)} \right]^k = \sum_{n=0}^{\infty} b_n^{(k)}(u) \frac{X^n}{n!}$$

then:

- for $n < -k$, $\mathcal{B}_{n,-k}(u) = 0$
- for $n \geq 0$, $\mathcal{B}_{n,-k}(u) = \frac{(-1)^{k-1}(k-1)!n!}{(n+k)!} b_{n+k}^{(k)}(u)$
- for $0 < n \leq k$, $\mathcal{B}_{-n,-k}(u) = \frac{(-1)^{n+k}(n-1)!(k-1)!}{(k-n)!} b_{k-n}^{(k)}(u)$

Theorem 3.1. *Let u be a sequence of order one and $0 < n \leq k$, then*

$$\mathcal{B}_{-n,-k}(u) = (-1)^{n+k} \mathcal{B}_{k,n}(\bar{u})$$

where \bar{u} is the inverse (for composition) of u .

Proof. The $\mathcal{B}_{n,k}(u)$ are rational functions in the variables u_1, u_2, \dots with coefficients in \mathbb{Q} . One can always suppose the u_n algebraically free on \mathbb{Q} and $\mathbb{Q}(u_1, u_2, \dots)$ embedded in the field of complex numbers \mathbb{C} . One can also suppose that $\overline{\lim} |u_n|^{1/n} < +\infty$, so that one can represent the $b_n^{(k)}(u)$ by Cauchy's formula:

$$\frac{b_{k-n}^{(k)}}{(k-n)!} = \frac{1}{2i\pi} \int_C \left(\frac{z}{g_u(z)} \right)^k \frac{dz}{z^{k-n+1}}, \quad 0 < n \leq k$$

C circle with radius $\varepsilon > 0$ and center 0.

By the change of variable $z = g_{\bar{u}}(t)$, and after integration, we obtain

$$\begin{aligned} \frac{b_{k-n}^{(k)}}{(k-n)!} &= \frac{k}{2i\pi n} \int_{C'} g_{\bar{u}}^n(t) \frac{dt}{t^{k+1}} \\ &= \frac{k}{n!} \frac{\mathcal{B}_{k,n}(\bar{u})}{k!} \end{aligned}$$

from which we get the relation of the theorem. ■

Corollary 3.2. For $0 < n \leq k$,

$$S(-n, -k) = (-1)^{n+k} s(k, n)$$

$$s(-n, -k) = (-1)^{n+k} S(k, n).$$

Remark 3.3. We check that:

$$\begin{aligned} (X)_{-n} &= \frac{1}{(X+1)\dots(X+n)} \\ &= \sum_{k \geq 0} s(-n, -k) X^{-k} = \sum_{k \geq 0} S(-n, -k) (X)_{-k}. \end{aligned}$$

More generally, if one considers the Bell exponential polynomials associated with a sequence u of order one:

$$Y_n(u, t) = \sum_{k \geq 0} \mathcal{B}_{n,k}(u) t^k$$

one can extend them to negative integers by the series:

$$Y_{-n}(u, t) = \sum_{k \geq 0} \mathcal{B}_{-n,-k}(u) t^{-k} = \sum_{k \geq n} (-1)^{n+k} \mathcal{B}_{k,n}(\bar{u}) t^{-k}.$$

For Bernoulli polynomials, with $D = \frac{d}{dX}$ and B_q the sequence of the classical Bernoulli numbers:

$$B_n(X) = g_{B_q}(D) \cdot X^n = (B_q \mathbin{\boxtimes} X^q)(n),$$

$$B_{-n}(X) = g_{B_q}(D) \cdot X^{-n} = \sum_{k \geq 0} \binom{-n}{k} B_k X^{-n-k}$$

since $g_{B_q}(D) = \frac{D}{e^D - 1}$, $B_{-n}(X)$ satisfies:

$$B_{-n}(X+1) - B_{-n}(X) = -nX^{-n-1}.$$

Remark 3.4. For k, n strictly positive integers

$$S(q, k) = \frac{1}{k!} a^{\boxtimes k} = \frac{1}{k!} (1^q - e_0)^{\boxtimes k}$$

where $a^{\boxtimes k}$ denote powers calculated in the Hurwitz algebra. From this we get:

$$S(n, k) = \frac{1}{k!} \sum_{j=1}^k \binom{k}{j} (-1)^{k-j} j^n$$

which allows to define an extension to the negative integers n and positive integers k .

To obtain an analogue for the Stirling numbers of second kind $s(n, k)$, we can consider

$$(X)_{-n} = \frac{1}{(X+1)\dots(X+n)} = \sum_{k \geq 0} s(-n, k) X^k$$

and hence

$$s(-n, k) = (-1)^{n+k-1} S(-k, n).$$

REFERENCES

- [1] D. Branson, *An extension of Stirling numbers*, Fib. Quat. **34** (3) (1996), 213–223.
- [2] L. Comtet, *Analyse combinatoire*, Vol. I and II, Presses Universitaires de France, Paris 1970.
- [3] S. Roman, *The harmonic logarithms and the binomial formula*, J. Combin. Theory, Serie A, **63** (1993), 143–163.

Received 2004
Revised December 2005