

## ON UNIFORMLY STRONGLY PRIME $\Gamma$ -SEMIRINGS (II)

TAPAN KUMAR DUTTA AND SUHRID DHARA

*Department of Pure Mathematics, University of Calcutta*  
*35, Ballygunge Circular Road, Kolkata-700019, India*

**e-mail:** [duttatapankumar@yahoo.co.in](mailto:duttatapankumar@yahoo.co.in)

**e-mail:** [suhridd@yahoo.com](mailto:suhridd@yahoo.com)

### Abstract

The Uniformly strongly prime  $k$ -radical of a  $\Gamma$ -semiring is a special class which we study via its operator semiring.

**Keywords:**  $\Gamma$ -semiring, uniformly right strongly prime  $\Gamma$ -semiring, annihilators, essential ideal, essential extension, uniformly strongly prime  $k$ -radical, matrix  $\Gamma$ -semiring, special class, super  $t$ -system.

**2000 Mathematics Subject Classification:** 16Y60.

### 1. INTRODUCTION

In 1987, D.M. Olson [7] introduced the notion of uniformly strongly prime radical in rings. In 1988, G.L. Booth and N.J. Groenwald [1] extended this notion of uniformly strongly prime radical to  $\Gamma$ -ring. In 1977, G.A.P. Heyman and C. Roos [6] introduced the notion of essential extension of rings.

In this paper we introduce the notions of uniformly right strongly prime ideal and uniformly left strongly prime ideal of a  $\Gamma$ -semiring and show that these two notions are equivalent. Also we study uniformly strongly prime  $k$ -radical of a  $\Gamma$ -semiring via its operators semirings as G.L. Booth and N.J. Groenwald did in case of  $\Gamma$ -ring. Some earlier works on the operator semiring of a  $\Gamma$ -semiring may be found in [4]. We obtain a relation between uniformly strongly prime  $k$ -radical of a  $\Gamma$ -semiring and with that of its matrix  $\Gamma$ -semiring via operator semiring. Lastly, we introduce the notion of super  $t$ -system in a  $\Gamma$ -semiring and obtain the relation between the uniformly strongly prime  $k$ -radical and super  $t$ -system in a  $\Gamma$ -semiring.

2. UNIFORMLY STRONGLY PRIME  $\Gamma$ -SEMIRINGS

**Definition 2.1** ([4]). Let  $S$  and  $\Gamma$  be two additive commutative semigroups. Then  $S$  is called a  $\Gamma$ -semiring if there exists a mapping  $S \times \Gamma \times S \longrightarrow S$  (image to be denoted by  $a\alpha b$ , for  $a, b \in S$  and  $\alpha \in \Gamma$ ) satisfying the following conditions:

- (i)  $a\alpha(b + c) = a\alpha b + a\alpha c$
- (ii)  $(a + b)\alpha c = a\alpha c + b\alpha c$
- (iii)  $a(\alpha + \beta)c = a\alpha c + a\beta c$
- (iv)  $a\alpha(b\beta c) = (a\alpha b)\beta c$

for all  $a, b, c \in S$  and for all  $\alpha, \beta \in \Gamma$ .

Every semiring  $S$  is a  $\Gamma$ -semiring with  $\Gamma = S$  where  $a\alpha b$  denotes the product of elements  $a, \alpha, b \in S$ .

If  $S$  contains an element  $0$  such that  $0+x = x = x+0$  and  $0\alpha x = x\alpha 0 = 0$  for all  $x \in S$ , for all  $\alpha \in \Gamma$ , then  $0$  is called the zero of  $S$ .

Throughout this paper we assume that a  $\Gamma$ -semiring always contains a zero element.

**Definition 2.2** ([4]). Let  $S$  be a  $\Gamma$ -semiring and  $L$  be the left operator semiring and  $R$  be the right operator semiring. If there exists an element

$$\sum_{i=1}^m [e_i, \delta_i] \in L \left( \text{respectively } \sum_{j=1}^n [\nu_j, f_j] \in R \right)$$

such that

$$\sum_{i=1}^m e_i \delta_i a = a \left( \text{respectively } \sum_{j=1}^n a \nu_j f_j = a \right) \text{ for all } a \in S$$

then  $S$  is said to have the left unity

$$\sum_{i=1}^m [e_i, \delta_i] \left( \text{respectively the right unity } \sum_{j=1}^n [\nu_j, f_j] \right).$$

**Definition 2.3** ([4]). A nonempty subset  $I$  of a  $\Gamma$ -semiring  $S$  is called an ideal of  $S$  if  $I + I \subseteq I$ ,  $I \Gamma S \subseteq I$ ,  $S \Gamma I \subseteq I$ , where for subsets  $U, V$  of  $S$  and  $\Delta$  of  $\Gamma$ ,

$$U\Delta V = \left\{ \sum_{i=1}^n u_i \gamma_i v_i : u_i \in U, v_i \in V, \gamma_i \in \Delta \text{ and } n \text{ is a positive integer} \right\}$$

**Definition 2.4.** A  $\Gamma$ -semiring  $S$  is called uniformly right strongly prime if  $S$  and  $\Gamma$  contain finite subsets  $F$  and  $\Delta$  respectively such that for any non zero  $x (\neq 0) \in S$ ,  $x\Delta F\Delta y = \{0\}$  implies that  $y = 0$  for all  $y \in S$ . The pair  $(F, \Delta)$  is called a uniform right insulator for  $S$ .

Analogously we can define uniformly left strongly prime  $\Gamma$ -semiring.

**Theorem 2.5.** A  $\Gamma$ -semiring  $S$  is uniformly right strongly prime if and only if there exist finite subsets  $F$  of  $S$  and  $\Delta$  of  $\Gamma$  such that for any two nonzero elements  $x$  and  $y$  of  $S$  there exist  $f \in F$  and  $\alpha, \beta \in \Delta$  such that  $x\alpha f\beta y \neq 0$ .

**Proof.** Let  $S$  be a uniformly right strongly prime  $\Gamma$ -semiring and  $(F, \Delta)$  be a uniform right insulator for  $S$ . Let  $x, y$  be any two nonzero elements of  $S$ . Suppose that  $x\Delta F\Delta y = \{0\}$ . Then  $y = 0$ , a contradiction. So there exist  $f \in S$  and  $\alpha, \beta \in \Delta$  such that  $x\alpha f\beta y \neq 0$ . ■

The converse follows by reversing the above argument.

**Corollary 2.6.** A  $\Gamma$ -semiring  $S$  is uniformly right strongly prime if and only if  $S$  is uniformly left strongly prime.

So we ignore the word right from uniformly right strongly prime.

**Definition 2.7.** A nonzero ideal  $I$  of a  $\Gamma$ -semiring  $S$  is called an essential ideal of  $S$  if for any nonzero ideal  $J$  of  $S$ ,  $I \cap J \neq (0)$ .

**Example 2.8.** Let  $S = \{r\omega : r \in \mathbf{Z}\}$  and  $\Gamma = \{r\omega^2 : r \in \mathbf{Z}\}$ , where  $\omega$  be a cube root of unity and  $\mathbf{Z}$  be the set of all integers. Then  $S$  is a  $\Gamma$ -semiring with usual addition and multiplication. Let  $I = \{2r\omega : r \in \mathbf{Z}\}$ . Then  $I$  is a nonzero ideal of  $S$ . Let  $J$  be any nonzero ideal of  $S$ . Then  $I \cap J \neq (0)$ . Hence  $I$  is an essential ideal of  $S$ .

**Definition 2.9.** A  $\Gamma$ -semiring  $T$  is said to be an essential extension of a  $\Gamma$ -semiring  $S$  if  $S$  is an essential ideal of  $T$ .

**Definition 2.10.** Let  $A$  be a nonempty subset of a  $\Gamma$ -semiring  $S$ . Right annihilator of  $A$  in  $S$ , denoted by  $\text{ann}_R(A)$ , is defined by  $\text{ann}_R(A) = \{s \in S : A\Gamma s = \{0\}\}$ .

Analogously we can define left annihilator  $\text{ann}_L(A)$  of  $A$  in  $S$ . Annihilator of a nonempty subset  $A$  is denoted by  $\text{ann}(A)$  which is a left as well as a right annihilator of  $A$ .

**Remark 2.11.** If  $S$  is a  $\Gamma$ -semiring then  $\text{ann}_R(A)$  is a right ideal of  $S$  and  $\text{ann}_L(A)$  is a left ideal of  $S$ . If  $A$  is an ideal of a  $\Gamma$ -semiring  $S$  then both annihilators are ideals of  $S$ .

**Lemma 2.12.** Let  $S$  be a  $\Gamma$ -semiring and  $T$  be its essential extension. If  $S$  is a uniformly strongly prime  $\Gamma$ -semiring then for each nonzero  $x$  of  $T$ ,  $x\Delta F = \{0\}$  implies that  $x \in \text{ann}_R(S)$  and  $F\Delta x = \{0\}$  implies that  $x \in \text{ann}_L(S)$ , where  $(F, \Delta)$  is a uniform insulator for  $S$ .

**Proof.** Let  $x\Delta F = \{0\}$ . Then  $s\alpha x\Delta F\Delta s\alpha x = \{0\}$  for all  $s \in S$  and for all  $\alpha \in \Gamma$ . Since  $S$  is an ideal of  $T$  and  $s \in S$ ,  $s\alpha x \in S$ . Again since  $S$  is uniformly strongly prime and  $(F, \Delta)$  is a uniform insulator for  $S$ ,  $s\alpha x = 0$  for all  $s \in S$  and for all  $\alpha \in \Gamma$  i.e.  $S\Gamma x = \{0\}$  i.e.  $x \in \text{ann}_R(S)$  (By Definition 2.10). ■

Similarly we can prove that  $F\Delta x = \{0\}$  implies that  $x \in \text{ann}_L(S)$ .

**Lemma 2.13.** Let  $S$  be a uniformly strongly prime  $\Gamma$ -semiring and  $T$  be its essential extension. Then both annihilators of  $S$  in  $T$  are zero.

**Proof.** Let  $(F, \Delta)$  be a uniform insulator for  $S$ . If possible let  $\text{ann}_R(S) \neq (0)$ . Then  $\text{ann}_R(S)$  is a nonzero ideal of  $T$ . Since  $S$  is an essential ideal of  $T$ ,  $\text{ann}_R(S) \cap S \neq (0)$ . Let  $x(\neq 0) \in \text{ann}_R(S) \cap S$ . Then  $S\Gamma x = \{0\}$ . Now  $x\Delta F\Delta x \subseteq x\Gamma S\Gamma x = \{0\}$ . Since  $S$  is a uniformly strongly prime  $\Gamma$ -semiring,  $x = 0$ , a contradiction. Therefore  $\text{ann}_R(S) = (0)$ . ■

Similarly we can prove that  $\text{ann}_L(S) = (0)$ .

**Theorem 2.14.** Any essential extension of a uniformly strongly prime  $\Gamma$ -semiring  $S$  is a uniformly strongly prime  $\Gamma$ -semiring.

**Proof.** Let  $(F, \Delta)$  be a uniform insulator for  $S$  and  $T$  be an essential extension of  $S$ . Let  $x$  be a nonzero element of  $T$ . Then  $x\Delta F$  and  $F\Delta x$  both are nonzero. Suppose  $x\Delta F = \{0\}$  then by Lemma 2.12,  $x \in \text{ann}_R(S)$ . Also

by Lemma 2.13,  $\text{ann}_R(S) = (0)$ , which implies that  $x = 0$ , a contradiction. Therefore  $x\Delta F \neq \{0\}$ . Similarly  $F\Delta x \neq \{0\}$ . Let  $y, z$  be two nonzero elements of  $T$ . Then there exist  $f_1, f_2 \in F$  and  $\alpha_1, \alpha_2 \in \Delta$  such that  $y\alpha_1 f_1 \neq 0$  and  $f_2\alpha_2 z \neq 0$ . Since  $S$  is an ideal of  $T$ , so  $y\alpha_1 f_1, f_2\alpha_2 z \in S$ . Again since  $S$  is uniformly strongly prime and  $(F, \Delta)$  is a uniform insulator for  $S$ , there exist  $\alpha, \beta \in \Delta$  and  $f \in F$  such that  $y\alpha_1 f_1 \alpha f \beta f_2 \alpha_2 z \neq 0$ . Let  $F' = \{f_1 \alpha f \beta f_2 : y\alpha_1 f_1 \alpha f \beta f_2 \alpha_2 z \neq 0; f_1, f, f_2 \in F; \alpha, \beta, \alpha_1, \alpha_2 \in \Delta; y, z \in T\}$ . Then  $F' \subseteq S \subseteq T$  is a finite subset, since  $F$  and  $\Delta$  are finite subsets. Hence by Theorem 2.5,  $T$  is uniformly strongly prime  $\Gamma$ -semiring with insulator  $(F', \Delta)$ . ■

**Definition 2.15.** Let  $S$  be a  $\Gamma$ -semiring and  $T$  be a nonempty subset of  $S$ . Then  $T$  is said to be a  $\Gamma$ -subsemiring of  $S$  if for  $t_1, t_2 \in T$  and  $\alpha \in \Gamma$ ,  $t_1 + t_2, t_1 \alpha t_2 \in T$ .

**Remark 2.16.** Every ideal of a  $\Gamma$ -semiring  $S$  is a  $\Gamma$ -subsemiring of  $S$ .

**Lemma 2.17.** If  $S$  is a uniformly strongly prime  $\Gamma$ -semiring and  $I$  is an ideal of  $S$ , then  $I$  is also a uniformly strongly prime  $\Gamma$ -subsemiring.

**Proof.** Let  $S$  be a uniformly strongly prime  $\Gamma$ -semiring and  $(F, \Delta)$  be a uniform insulator for  $S$ . If  $I = (0)$  then obviously  $I$  is a uniformly strongly prime  $\Gamma$ -subsemiring. Suppose  $I \neq (0)$  and  $r$  be a fixed nonzero element of  $I$ . Let  $F' = \{f_1 \alpha r \beta f_2 : f_1, f_2 \in F; \alpha, \beta \in \Delta\}$ . Since  $I$  is an ideal of  $S$  and  $F, \Delta$  are finite subsets,  $F'$  is a finite subset of  $I$ . Let  $x (\neq 0) \in I$  and  $y \in I$ . Now  $x\Delta F'\Delta y = \{0\}$  implies that  $x\gamma f_1 \alpha r \beta f_2 \delta y = 0$  for all  $f_1, f_2 \in F$  and for all  $\alpha, \beta, \gamma, \delta \in \Delta$  i.e.  $x\Delta F\Delta(r\beta f_2 \delta y) = \{0\}$  for all  $f_2 \in F$  and for all  $\beta, \delta \in \Delta$ . Since  $r\beta f_2 \delta y \in S$  for all  $f_2 \in F$  and for all  $\beta, \delta \in \Delta$  and  $S$  is a usp  $\Gamma$ -semiring with  $x \neq 0$ ,  $r\beta f_2 \delta y = 0$  for all  $f_2 \in F$  and for all  $\beta, \delta \in \Delta$  i.e.  $r\Delta F\Delta y = \{0\}$ . By previous argument  $y = 0$  as  $r \neq 0$ . Hence  $(F', \Delta)$  is a uniform insulator for  $I$ . Thus  $I$  is a uniformly strongly prime  $\Gamma$ -subsemiring. ■

### 3. UNIFORMLY STRONGLY PRIME $k$ -RADICALS

**Definition 3.1** ([3]). An ideal  $P$  of a  $\Gamma$ -semiring  $S$  is called a uniformly strongly prime ideal (usp ideal) if  $S$  and  $\Gamma$  contain finite subsets  $F$  and  $\Delta$  respectively such that  $x\Delta F\Delta y \subseteq P$  implies that  $x \in P$  or  $y \in P$  for all  $x, y \in S$ .

**Definition 3.2.** Let  $S$  be a  $\Gamma$ -semiring. The uniformly strongly prime  $k$ -radical (usp  $k$ -radical) of a  $\Gamma$ -semiring  $S$ , denoted by  $\tau(S)$  is defined by

$$\tau(S) = \bigcap_{P \in \Lambda_S} P,$$

where  $\Lambda_S$  denote the set of all usp  $k$ -ideals of the  $\Gamma$ -semiring  $S$ .

**Definition 3.3.** Let  $S$  be a  $\Gamma$ -semiring and  $L$  (respectively  $R$ ) be its left (respectively right) operator semiring, the usp  $k$ -radical of  $L$  (respectively  $R$ ), denoted by  $\tau(L)$  (respectively  $\tau(R)$ ) is defined by

$$\tau(L) = \bigcap_{A \in \Lambda_L} A,$$

where  $\Lambda_L$  denote the set of all usp  $k$ -ideals of the left operator semiring  $L$  (respectively  $\tau(R) = \bigcap_{B \in \Lambda_R} B$ , where  $\Lambda_R$  denote the set of all usp  $k$ -ideals of the right operator semiring  $R$ ).

**Theorem 3.4.** Let  $S$  be a  $\Gamma$ -semiring with left and right unities and  $L$  be its left operator semiring then  $\tau(L)^+ = \tau(S)$  and  $\tau(L) = \tau(S)^{+'}$ .

**Proof.** Let  $\Lambda_L$  and  $\Lambda_S$  denote the set of all usp  $k$ -ideals of  $L$  and  $S$  respectively, then

$$\tau(L) = \bigcap_{A \in \Lambda_L} A \quad \text{and} \quad \tau(S) = \bigcap_{P \in \Lambda_S} P.$$

Hence

$$(\tau(L))^+ = \left( \bigcap_{A \in \Lambda_L} A \right)^+ = \bigcap_{A \in \Lambda_L} A^+.$$

Since for every  $A \in \Lambda_L$ ,  $A^+ \in \Lambda_S$  (Cf. Theorem 2.23 of [3]),

$$\bigcap_{A \in \Lambda_S} A \subseteq \bigcap_{A \in \Lambda_L} A^+.$$

Hence

$$(1) \quad \tau(S) \subseteq (\tau(L))^+.$$

Again

$$\tau(S) = \bigcap_{P \in \Lambda_S} P = \bigcap_{P \in \Lambda_S} (P^{+'})^+ = \left( \bigcap_{P \in \Lambda_S} P^{+'} \right)^+.$$

Since for each  $P \in \Lambda_S$ ,  $P^{+'} \in \Lambda_L$  (Cf. Theorem 2.23 of [3] ),

$$\bigcap_{P \in \Lambda_L} P \subseteq \bigcap_{P \in \Lambda_S} P^{+'}.$$

Hence

$$\left( \bigcap_{P \in \Lambda_L} P \right)^+ \subseteq \left( \bigcap_{P \in \Lambda_S} P^{+'} \right)^+,$$

which implies that

$$(2) \quad (\tau(L))^+ \subseteq \tau(S).$$

Thus from (1) and (2) we get  $\tau(S) = (\tau(L))^+$ . ■

Similarly we can prove that  $\tau(L) = \tau(S)^{+'}$ .

**Proposition 3.5.** *Let  $S$  be a  $\Gamma$ -semiring with left and right unities and  $R$  be its right operator semiring then  $\tau(R)^* = \tau(S)$  and  $\tau(R) = \tau(S)^{+'}$ .*

**Corollary 3.6.** *Let  $S$  be a  $\Gamma$ -semiring with left and right unities  $L$  and  $R$  be its left and right operator semirings then  $\tau(L)^+ = \tau(S) = \tau(R)^*$ .*

**Theorem 3.7.** *Let  $S$  be a semiring with identity 1 and  $\tau'(S)$ ,  $\tau(S)$  denote respectively the usp  $k$ -radical of the semiring  $S$  and the usp  $k$ -radical of the  $\Gamma$ -semiring  $S$ , where  $\Gamma = S$ , then  $\tau'(S) = \tau(S)$ .*

**Proof.** By Theorem 2.26 of [3] every usp  $k$ -ideal of a semiring  $S$  is a usp  $k$ -ideal of the  $\Gamma$ -semiring  $S$ , where  $\Gamma = S$ . This follows that  $\tau'(S) = \tau(S)$ . ■

#### 4. USP $k$ -RADICALS IN MATRIX $\Gamma$ -SEMIRING

**Definition 4.1** ([5]). Let  $S$  be a  $\Gamma$ -semiring and  $m, n$  be positive integers. We denote by  $S_{mn}$  and  $\Gamma_{nm}$  respectively the sets of  $m \times n$  matrices with entries from  $S$  and  $n \times m$  matrices with entries from  $\Gamma$ . Let  $A, B \in S_{mn}$

and  $\Delta \in \Gamma_{nm}$ . Then  $A\Delta B \in S_{mn}$  and  $A + B \in S_{mn}$ . Clearly,  $S_{mn}$  forms a  $\Gamma_{nm}$ -semiring with these operations. We call it the matrix  $\Gamma$ -semiring  $S$  or the matrix  $\Gamma_{nm}$ -semiring  $S_{mn}$  or simply the  $\Gamma_{nm}$ -semiring  $S_{mn}$ .

We denote the right operator semiring of the matrix  $\Gamma_{nm}$ -semiring  $S_{mn}$  by  $[\Gamma_{nm}, S_{mn}]$  and the left one by  $[S_{mn}, \Gamma_{nm}]$ . If  $x \in S$ , the notation  $xE_{ij}$  will be used to denote a matrix in  $S_{mn}$  with  $x$  in the  $(i, j)$ -th entry and zeros elsewhere. The notation  $\alpha E_{ij}$ , where  $\alpha \in \Gamma$  will have a similar meaning. If  $P \subseteq S$ ,  $P_{mn}$  will denote the set of all  $m \times n$  matrices with entries from  $P$ . If  $\Delta \subseteq \Gamma$ ,  $\Delta_{nm}$  is similarly defined.

**Theorem 4.2.** *Let  $S$  be a  $\Gamma$ -semiring with left and right unities. Then  $S$  is a usp  $\Gamma$ -semiring if and only if  $S_{mn}$  is a usp  $\Gamma_{nm}$ -semiring for all positive integers  $m$  and  $n$ .*

**Proof.**  $S$  is a usp  $\Gamma$ -semiring if and only if  $L$  is a usp semiring if and only if  $L_m$  is a usp semiring if and only if  $S_{mn}$  is a usp  $\Gamma_{nm}$ -semiring (By Theorem 2.18 of [3] and by Lemma 3.13 of [2]). ■

**Lemma 4.3.** *Let  $S$  be a  $\Gamma$ -semiring and  $m, n$  be positive integers; then a nonempty subset  $P$  of  $S_{mn}$  is a usp  $k$ -ideal of  $S_{mn}$  if and only if  $P = Q_{mn}$ , for some usp  $k$ -ideal  $Q$  of  $S$ .*

**Proof.** Let  $L$  be the left operator semiring of  $S$  and  $L_m$  be the left operator semiring of the  $\Gamma_{nm}$ -semiring  $S_{mn}$ . Let  $P$  be a usp  $k$ -ideal of  $S_{mn}$ . Then by Theorem 2.23 of [3] (applied to  $\Gamma_{nm}$ -semiring  $S_{mn}$ )  $P^{+'}$  is a usp  $k$ -ideal of the left operator semiring  $L_m$  of the  $\Gamma_{nm}$ -semiring  $S_{mn}$ . So  $P^{+'} = Q_m$  for some usp  $k$ -ideal  $Q$  of  $L$ . This implies that  $(P^{+'})^+ = (Q_m)^+$  i.e.  $P = (Q^+)_{mn}$  (By Theorem 2.10 and Proposition 3.4 of [5]). Again by Theorem 2.23 of [3],  $Q^+$  is an usp  $k$ -ideal of  $S$ . So the direct implication follows.

Conversely, suppose  $Q$  is a usp  $k$ -ideal of  $S$ . Then  $Q^{+'}$  is a usp  $k$ -ideal of  $L$  (By Theorem 2.23 of [3]). So  $(Q^{+'})_m$  is a usp  $k$ -ideal of  $L_m$  i.e.  $(Q_{mn})^{+'}$  is a usp  $k$ -ideal of  $L_m$  by previous arguments. Hence  $Q_{mn} = ((Q_{mn})^{+'})^+$  is an usp  $k$ -ideal of the  $\Gamma_{nm}$ -semiring  $S_{mn}$ . ■

**Lemma 4.4.** *Let  $S$  be a  $\Gamma$ -semiring and  $I$  be an ideal of  $S$ . Then  $(S/I)_{mn}$  is isomorphic to  $S_{mn}/I_{mn}$  for all positive integers  $m, n$ .*

**Proof.** Let  $\theta$  be a mapping of the  $\Gamma_{nm}$ -semiring  $(S/I)_{mn}$  to the  $\Gamma_{nm}$ -semiring  $S_{mn}/I_{mn}$  defined by  $\theta((x_{ij}/I)_{mn}) = (x_{ij})_{mn}/I_{mn}$ . Let  $(x_{ij}/I)_{mn}$ ,



$(y_{ij}/I)_{mn} \in (S/I)_{mn}$ . Now  $(x_{ij}/I)_{mn} = (y_{ij}/I)_{mn}$  if and only if  $x_{ij}/I = y_{ij}/I$ , for all  $i, j$  if and only if  $x_{ij} + a_{ij} = y_{ij} + b_{ij}$ , for some  $a_{ij}, b_{ij} \in I$  and for all  $i, j$  if and only if  $(x_{ij} + a_{ij})_{mn} = (y_{ij} + b_{ij})_{mn}$  if and only if  $(x_{ij})_{mn} + (a_{ij})_{mn} = (y_{ij})_{mn} + (b_{ij})_{mn}$  for some matrices  $(a_{ij})_{mn}, (b_{ij})_{mn} \in I_{mn}$  if and only if  $(x_{ij})_{mn}/I_{mn} = (y_{ij})_{mn}/I_{mn}$  if and only if  $\theta((x_{ij}/I)_{mn}) = \theta((y_{ij}/I)_{mn})$ . Therefore  $\theta$  is well defined as well as injective.

Clearly  $\theta$  is onto and semigroup homomorphism under addition.

Now

$$\begin{aligned}
 & \theta\left((x_{ij}/I)_{mn}(\alpha_{ij})_{nm}(y_{ij}/I)_{mn}\right) \\
 &= \theta\left(\left(\sum_{k=1}^n \sum_{l=1}^m (x_{ik}\alpha_{kl}y_{lj})/I\right)_{mn}\right) \\
 &= \left(\sum_{k=1}^n \sum_{l=1}^m (x_{ik}\alpha_{kl}y_{lj})\right)_{mn} / I_{mn} \\
 &= ((x_{ij})_{mn}(\alpha_{ij})_{nm}(y_{ij})_{mn})/I_{mn} \\
 &= ((x_{ij})_{mn}/I)(\alpha_{ij})_{nm}((y_{ij})_{mn}/I) \\
 &= \theta((x_{ij}/I)_{mn})\iota((\alpha_{ij})_{nm})\theta((y_{ij}/I)_{mn}).
 \end{aligned}$$

This shows that  $(\theta, \iota)$  is an isomorphism of  $(S/I)_{mn}$  onto  $S_{mn}/I_{mn}$ , where  $\iota$  is the identity mapping from  $\Gamma_{nm}$  onto  $\Gamma_{nm}$ . ■

**Proposition 4.5.** *Let  $A, B$  be two ideals of a  $\Gamma$ -semiring  $S$ . Then  $(A \cap B)_{mn} = A_{mn} \cap B_{mn}$ , where  $A_{mn}, B_{mn}$  are two ideals of  $\Gamma_{nm}$ -semiring  $S_{mn}$ .*

**Theorem 4.6.** *Let  $S$  be a  $\Gamma$ -semiring. Then  $\tau(S_{mn}) = (\tau(S))_{mn}$  for all positive integers  $m, n$ .*

**Proof.** By Definition 3.2,

$$\tau(S_{mn}) = \bigcap_{P \in \Lambda_{S_{mn}}} P,$$

where  $\Lambda_{S_{mn}}$  denote the set of all usp  $k$ -ideals of the  $\Gamma$ -semiring  $S_{mn}$  and

$$\tau(S) = \bigcap_{Q \in \Lambda_S} Q,$$

where  $\Lambda_S$  denote the set of all usp  $k$ -ideals of the  $\Gamma$ -semiring  $S$ .  
Therefore

$$(\tau(S))_{mn} = \left( \bigcap_{Q \in \Lambda_S} Q \right)_{mn} = \bigcap_{Q \in \Lambda_S} Q_{mn}$$

(By Proposition 4.5). Hence by Lemma 4.3 we get the result. ■

## 5. SPECIAL CLASSES

**Definition 5.1.** A class  $\wp$  of  $\Gamma$ -semirings is called hereditary if  $I$  is an ideal of a  $\Gamma$ -semiring  $S$  and  $S \in \wp$  implies that  $I \in \wp$ .

**Definition 5.2.** A class  $\wp$  of  $\Gamma$ -semirings is called closed under essential extension if  $I$  is an essential ideal of a  $\Gamma$ -semiring  $S$  and  $I \in \wp$  implies that  $S \in \wp$ .

**Definition 5.3.** The class  $\wp$  of  $\Gamma$ -semirings is called a special class if

- (i)  $\wp$  consists of prime  $\Gamma$ -semirings,
- (ii)  $\wp$  is hereditary and
- (iii)  $\wp$  is closed under essential extension.

**Theorem 5.4.** A class  $\mathcal{L}$  of uniformly strongly prime  $\Gamma$ -semirings is a special class.

**Proof.** Since uniformly strongly prime implies prime, so  $\mathcal{L}$  consists of all prime  $\Gamma$ -semirings. By Lemma 2.17, we get  $\mathcal{L}$  is hereditary class. Again by Theorem 2.14,  $\mathcal{L}$  is closed under essential extension. Hence  $\mathcal{L}$  is a special class. ■

**Proposition 5.5.** *The following conditions are equivalent for any class  $\rho$  of prime  $\Gamma$ -semirings:*

- (i) *if  $I$  is an ideal of  $S$ ,  $I \in \rho$  and  $\text{ann}(I) = (0)$ , then  $S \in \rho$*
- (ii)  *$\rho$  is closed under essential extension.*

**Proof.** (i) $\Rightarrow$ (ii) Let  $I$  be an essential ideal of  $S$  and  $I \in \rho$ . Let  $x \in I \cap \text{ann}(I)$  then  $x \in I$  and  $x \in \text{ann}(I)$  which implies that  $x \in I$  and  $I \Gamma x = \{0\} = x \Gamma I$ . So  $x \Gamma I \Gamma x = \{0\}$  which implies that  $\langle x \rangle \Gamma \langle x \rangle = \{0\}$ . Since  $(0)$  is a prime ideal of  $S$  so it follows that  $x = 0$ . Therefore  $\text{ann}(I) = (0)$ . Hence by (i),  $S \in \rho$  which imply (ii).

(ii) $\Rightarrow$ (i) Let  $\rho$  be closed under essential extension. Let  $I \in \rho$  be an ideal of  $S$  with  $\text{ann}(I) = (0)$ . Let  $L$  be an ideal of  $S$  such that  $I \cap L = (0)$ . Now  $I \Gamma L \subseteq I \cap L = (0)$  and  $L \Gamma I \subseteq I \cap L = (0)$ . So  $I \Gamma L = (0) = L \Gamma I$  which implies that  $L \subseteq \text{ann}(I) = (0)$ , so  $L = (0)$ . Therefore  $I$  is an essential ideal of  $S$ . Also  $I \in \rho$ . Hence by (ii)  $S \in \rho$ . ■

A  $\Gamma$ -semiring  $S$  is called a us(1) prime if it has an insulator of the form  $(\{x\}, \{\gamma\})$  where  $x \in S$  and  $\gamma \in \Gamma$ .

As in Theorem 5.4 we can show that the class  $\mathcal{L}_1$  of all us(1) prime  $\Gamma$ -semirings is a special class.

**Definition 5.6.** A pair of subsets  $(T, I)$  of a  $\Gamma$ -semiring  $S$  is called a super  $t$ -system of  $S$  if

- (i)  $I$  is an ideal of  $S$
- (ii)  $T \cap I \subseteq (0)$  and
- (iii) there exist finite subsets  $F$  of  $S$  and  $\Delta$  of  $\Gamma$  such that for all  $a, b \in S \setminus I$ ,  $a \Delta F \Delta b \cap T \neq \phi$ .

The pair  $(F, \Delta)$  will be called an insulator of the super  $t$ -system. Therefore  $I$  is a uniformly strongly prime ideal of  $S$  if and only if  $(S \setminus I, I)$  is a super  $t$ -system.

**Theorem 5.7** *For a  $\Gamma$ -semiring  $S$ ,  $\tau(S) = \{a \in S: \text{whenever } (T, I) \text{ is a super } t\text{-system in } S \text{ with } a \in T \text{ then } 0 \in T\}$ .*

**Proof.** Let  $H = \{a \in S: \text{whenever } (T, I) \text{ is a super } t\text{-system in } S \text{ with } a \in T \text{ then } 0 \in T\}$ . Let  $a \in \tau(S)$ . Suppose  $(T, I)$  is a super  $t$ -system in  $S$  with  $a \in T$  and  $0 \notin T$ . So  $T \cap I = \phi$ . Let  $(F, \Delta)$  be an insulator of the super  $t$ -system. By Zorn's lemma there exists a maximal ideal  $P$  such that  $I \subseteq P$  and  $T \cap P = \phi$ . We now prove that  $P$  is a uniformly strongly prime ideal of  $S$  with  $(F, \Delta)$  be its uniform insulator. If possible let there exist  $x, y \notin P$  such that  $x\Delta F\Delta y \subseteq P$ . Since  $x, y \notin P$ , then  $x, y \notin I$  which implies that  $x, y \in S \setminus I$  and  $(x\Delta F\Delta y) \cap T \subseteq P \cap T = \phi$ , a contradiction, since  $(T, I)$  is a super  $t$ -system in  $S$ . So  $x\Delta F\Delta y \subseteq P$  implies that  $x \in P$  or  $y \in P$  which implies that  $P$  is a uniformly strongly prime ideal of  $S$ . Hence  $a \in \tau(S)$  implies that  $a \in P$ . Again  $a \in T \Rightarrow a \notin P$  as  $T \cap P = \phi$ , which is a contradiction. So  $0 \in T$ . Then  $a \in H \Rightarrow \tau(S) \subseteq H$ .

Conversely, let  $a \notin \tau(S)$ . Then there exists a uniformly strongly prime  $k$ -ideal and hence an ideal say  $Q$  of  $S$  such that  $a \notin Q$ . Then  $a \in S \setminus Q$ , so  $(S \setminus Q, Q)$  is a super  $t$ -system with  $a \in S \setminus Q$  but  $0 \notin S \setminus Q$ , so  $a \notin H$ . Hence  $H \subseteq \tau(S)$ . This completes the proof. ■

### Acknowledgement

The authors are thankful to the learned referee for his kind suggestions.

### REFERENCES

- [1] G.L. Booth and N.J. Groenwald, *On Uniformly Strongly Prime  $\Gamma$ -rings*, Bull. Austral Math. Soc. **37** (1988), 437–445.
- [2] T.K. Dutta and M.L. Das, *On Uniformly Strongly Prime semirings* (communicated).
- [3] T.K. Dutta and S. Dhara, *On Uniformly Strongly Prime  $\Gamma$ -semirings*, Southeast Asian Bull. of Mathematics **30** (2006), 32–39.
- [4] T.K. Dutta and S.K. Sardar, *On The Operator Semirings of a  $\Gamma$ -semiring*, Southeast Asian Bull. of Mathematics, Springer-Verleg, **26** (2002), 203–213.
- [5] T.K. Dutta and S.K. Sardar, *On Matrix  $\Gamma$ -semirings*, Far East J. Math. Soci. (FJMS) **7** (1) (2002), 17–31.
- [6] G.A.P. Heyman and C. Roos, *Essential extensions in radical theory for rings* J. Austral. Math. Soc. Ser A, **23** (1977), 340–347.

- [7] D.M. Olson, *A Uniformly Strongly Prime Radicals* J. Austral Math. Soc. Ser A **43** (1) (1987), 95–102.
- [8] D.M. Olson and S. Veldsman, *Some remarks on Uniformly Strongly prime Radicals*, Math. Japon **33** (3) (1988), 445–455.

Received 10 July 2006

Revised 13 September 2006