THE LATTICE OF VARIETIES OF FIBERED AUTOMATA

Anna Mućka

Faculty of Mathematics and Information Sciences, Warsaw University of Technology 00-661 Warsaw, Poland

e-mail: aniam@b4.pl

Abstract

The class of all fibered automata is a variety of two-sorted algebras. This paper provides a full description of the lattice of varieties of fibred automata.

Keywords: fibered automata, many-sorted algebras, varieties of many-sorted algebras, lattice of varieties.

2000 Mathematics Subject Classification: 08B15, 08A68, 68Q70.

References

- [1] G. Birkhoff and J.D. Lipson, *Heterogenous algebras*, J. Comb. Th. 8 (1970), 115–133
- [2] J.A. Goguen and J. Meseguer, Completeness of many-sorted equational logic, Houston J. Math. 11 (1985), 307–334.
- [3] H. Lugowski, Grundzüge der Universalen Algebra, Teuber, Leipzig 1976.
- [4] I. Rosenberg, A classification of universal algebras by infinitary relations, Algebra Universalis 1 (1972), 350–353.
- [5] J.D.H. Smith, Continued fractions, fibered automata, and a theorem of Rosenberg, Multiple-Valued Logic 8 (2002), 503–515.

Received 6 April 2006 Revised 1 July 2006