Discussiones Mathematicae General Algebra and Applications 28 (2008) 77–89

# COMMUTATIVE DIRECTOIDS WITH SECTIONALLY ANTITONE BIJECTIONS\*

Ivan Chajda, Miroslav Kolařík

Department of Algebra and Geometry Palacký University Olomouc Tomkova 40, 779 00 Olomouc, Czech Republic e-mail: chajda@inf.upol.cz e-mail: kolarik@inf.upol.cz

AND

## Sándor Radeleczki

Institute of Mathematics University of Miskolc 3515 Miskolc-Egyetemváros, Hungary e-mail: matradi@gold.uni-miskolc.hu

#### Abstract

We study commutative directoids with a greatest element, which can be equipped with antitone bijections in every principal filter. These can be axiomatized as algebras with two binary operations satisfying four identities. A minimal subvariety of this variety is described.

**Keywords:** directoid, section antitone bijection, implication algebra, double implication algebra.

2000 Mathematics Subject Classification: 06A12, 03G10, 03G25.

Join-semilattices whose principal filters are Boolean lattices were used by J.C. Abbott [1] for a characterization of the logic connective implication in the classical propositional logic. These semilattices also have the property that on each principal filter of them an antitone involution is defined.

<sup>\*</sup>The first two authors are supported by the Czech Government via the project No. MSM 6198959214.

Motivated by this observation, the notion of a *join semilattice with section*ally antitone involutions in [2] and *join semilattice with sectionally antitone* bijections in [5] was defined. In this paper we introduce a further generalization of this concept defining the notion of a *commutative directoid with* sectionally antitone bijections. By means of these directoids we obtain "nice" algebraic structures, forming a variety with "nice" congruence properties.

The concept of directoid was introduced by J. Ježek and R. Quackenbush [8] and independently by V.M. Kopytov and Z.I. Dimitrov [9] in order to axiomatize algebraic structures defined on upward directed ordered sets. In a certain sense directoids generalize semilattices. For the reader's convenience, we repeat definitions and basic properties of these concepts.

An ordered set  $(A; \leq)$  is upward directed if  $U(x, y) \neq \emptyset$  for every  $x, y \in A$ , where  $U(x, y) = \{a \in A; x \leq a \text{ and } y \leq a\}$ . Elements of U(x, y) are referred to as common upper bounds of x, y. Of course, if  $(A; \leq)$  has a greatest element then it is upward directed.

Let  $(A; \leq)$  be an upward directed set and  $\sqcup$  denots a binary operation on A. The pair  $\mathcal{A} = (A; \sqcup)$  is called a  $\sqcup$ -directoid if

- (i)  $x \sqcup y \in U(x, y)$  for all  $x, y \in A$ ;
- (ii) if  $x \leq y$  then  $x \sqcup y = y$  and  $y \sqcup x = y$ .

The following axiomatization of directoids was established in [8]:

**Proposition 1.** A groupoid  $\mathcal{A} = (A; \sqcup)$  is a  $\sqcup$ -directoid if and only if it satisfies the following identities

- (D1)  $x \sqcup x = x;$
- (D2)  $(x \sqcup y) \sqcup x = x \sqcup y;$
- (D3)  $y \sqcup (x \sqcup y) = x \sqcup y;$
- (D4)  $x \sqcup ((x \sqcup y) \sqcup z) = (x \sqcup y) \sqcup z.$

A binary relation  $\leq$  defined on A by the rule

 $x \leq y$  if and only if  $x \sqcup y = y$ 

is an order and  $x \sqcup y \in U(x, y)$  for each  $x, y \in A$ .

A binary relation  $\leq$  will be called the *induced order* of  $(A; \sqcup)$ . A directoid  $\mathcal{A} = (A; \sqcup)$  is called *commutative* if it satisfies the identity

(D5) 
$$x \sqcup y = y \sqcup x$$
.

It was shown in [8] that commutative directoids are axiomatized by the identities (D1), (D4) and (D5).

Let us denote a greatest element of an ordered set  $(P; \leq)$  by 1. For each  $a \in P$  the interval [a, 1] will be called a *section*.

**Definition 1.** Let  $(P; \leq)$  be a poset with a greatest element 1.  $(P; \leq)$  is called a *poset with sectionally antitone bijections*, if for each element  $a \in S$  there exists a bijection  $f_a$  of the interval  $[a, 1] \subseteq P$  into itself such that

$$x \le y \quad \Leftrightarrow \quad f_a(y) \le f_a(x), \text{ for all } x, y \in [a, 1].$$

Of course, the inverse  $f_a^{-1}$  of  $f_a$  is also an antitone bijection on [a, 1]. If each  $f_a$  is an *involution*, i.e.,  $f_a^2(x) = x$ , for all  $x \in [a, 1]$ , then  $f_a^{-1} = f_a$  and  $(P; \leq)$  is called a *poset with sectionally antitone involutions*.

**Remark 1.** Any poset  $(P; \leq)$  with a greatest element possesses a commutative  $\sqcup$ -directoid  $(P; \sqcup)$ .

Indeed, take any choice function  $\varphi : 2^P \to P$ , such that for any nonempty subset  $A \subseteq P$ ,  $\varphi(A) \in A$ . Then define a binary operation  $\sqcup$  on P as follows

$$x \sqcup y = \begin{cases} \varphi(U(x,y)), & \text{if } x \text{ and } y \text{ are incomparable} \\ y & \text{if } x \leq y \\ x & \text{if } y \leq x. \end{cases}$$

It is evident that this operation satisfies the axioms (D1)-(D5) and that

$$x \le y \quad \Leftrightarrow \quad x \sqcup y = y.$$

Given a commutative  $\sqcup$ -directoid S with sectionally antitone bijections, we can introduce two new binary operations on S as follows:

(P) 
$$x \circ y = f_y(x \sqcup y), \qquad x * y = f_y^{-1}(x \sqcup y).$$

Since  $x \sqcup y \in [y, 1]$ ,  $\circ$  and \* are everywhere defined operations on the set S. Conversely, one can check immediately that for any  $a \in S$  and  $x \in [a, 1]$ 

(A) 
$$f_a(x) = x \circ a, \qquad f_a^{-1}(x) = x * a.$$

Clearly, if all the mappings  $f_a$  are involutions, then  $x \circ y = x * y$ , for all  $x, y \in S$ .

**Lemma 1.** Let S be a commutative  $\sqcup$ -directoid with sectionally antitone bijections and  $\circ$ , \* be operations defined by (P). Then

 $a \leq b \iff a \circ b = 1 \iff a * b = 1.$ 

**Proof.** Suppose  $a, b \in S$  and  $a \leq b$ . Since  $f_b$  is an antitone bijection on [b, 1], we have  $f_b(b) = 1$ . Then

(Q) 
$$a \circ b = f_b(a \sqcup b) = f_b(b) = 1, \quad a * b = f_b^{-1}(a \sqcup b) = f_b^{-1}(b) = 1.$$

To prove the converse implication we note that for all  $x \in [b, 1]$  it is  $b \leq x$ and thus  $f_b(b) \geq f_b(x)$ . Let  $a \circ b = 1$ . Then  $f_b(a \sqcup b) = 1 = f_b(b)$ , but  $f_b$  is a bijection, hence  $a \sqcup b = b$ , i.e.,  $a \leq b$ . Analogously we obtain  $a * b = 1 \Rightarrow a \leq b$ .

**Theorem 1.** Let S be a commutative  $\sqcup$ -directoid with sectionally antitone bijections and  $\circ$ , \* be operations defined by (P). Then

- (1)  $x \circ x = x * x = 1$ ,  $x \circ 1 = x * 1 = 1$ ,  $1 \circ x = 1 * x = x$ ;
- (2)  $(x \circ y) * y = (x * y) \circ y = (y \circ x) * x = (y * x) \circ x;$
- (3)  $x \circ ((((x \circ y) * y) \circ z) * z) = 1;$
- (4)  $(((((x \circ z) * z) \circ y) * y) \circ z) \circ (x \circ z) = (((((x \circ z) * z) \circ y) * y) * z) \circ (x * z) = 1.$

Moreover, each of the terms of (2) is equal to  $x \sqcup y$ .

## Proof.

- (1) By Lemma 1,  $x \circ x = x * x = 1$  and  $x \circ 1 = x * 1 = 1$ . We also conclude  $1 \circ x = f_x(1) = x$  and  $1 * x = f_x^{-1}(1) = x$ . Thus (1) is satisfied.
- (2)  $(x \circ y) * y = f_y^{-1}(f_y(x \sqcup y) \sqcup y) = f_y^{-1}(f_y(x \sqcup y)) = x \sqcup y \text{ since } f_y(x \sqcup y) \ge y.$ Analogously, we can check  $(x * y) \circ y = x \sqcup y, (y \circ x) * x = x \sqcup y,$  and  $(y * x) \circ x = x \sqcup y$ . Due to the fact that S is commutative, we can establish the last two equalities.

Commutative directoids with sectionally ...

- (3)  $x \circ ((((x \circ y) * y) \circ z) * z) = x \circ ((x \sqcup y) \sqcup z) = f_{(x \sqcup y) \sqcup z}(x \sqcup ((x \sqcup y) \sqcup z)) = f_{(x \sqcup y) \sqcup z}((x \sqcup y) \sqcup z) = 1$  by (D4).
- (4)  $((((x \circ z) * z) \circ y) * y) \circ z = ((x \sqcup z) \sqcup y) \circ z = f_z(((x \sqcup z) \sqcup y) \sqcup z)$ . Since  $f_z$  is antitone and  $z \le (x \sqcup z) \sqcup y$ ,  $x \sqcup z \le (x \sqcup z) \sqcup y$  we have:

$$f_z(((x \sqcup z) \sqcup y) \sqcup z) = f_z((x \sqcup z) \sqcup y) \le f_z(x \sqcup z) = x \circ z,$$

thus, by (Q):

$$\left(\left(\left(\left((x\circ z)*z\right)\circ y\right)*y\right)\circ z\right)\circ(x\circ z)=1.$$

Analogously we obtain:

$$\left(\left(\left(\left(x\circ z)*z\right)\circ y\right)*y\right)*z\right)\circ\left(x*z\right)=1.$$

**Theorem 2.** Let  $\mathcal{A} = (A; \circ, *, 1)$  be an algebra of type (2, 2, 0) satisfying the identities (1), (2) and (3). Define a binary relation  $\leq$  on A as follows:

(R) 
$$a \le b \text{ if and only if } a \circ b = 1.$$

Then  $\leq$  is a partial order on A with a greatest element 1 and  $(A; \leq)$  is a commutative  $\sqcup$ -directoid  $(A; \sqcup)$  where  $(x \circ y) * y = x \sqcup y$  and the following assertions are equivalent:

- (i) The algebra  $\mathcal{A}$  satisfies the identity (4);
- (ii) for any  $x, y, a \in A$ ,  $x \in [a, 1]$  the following implication holds:

(5)  $x \leq y \Rightarrow y \circ a \leq x \circ a \text{ and } y * a \leq x * a.$ 

Moreover, for any  $a \in A$  the mappings  $f_a(x) = x \circ a$ ,  $f_a^{-1}(x) = x * a$ are mutually inverse antitone bijections on [a, 1].

**Proof.** First we prove that the relation  $\leq$  defined by (R) is a partial order. Due to (1),  $\leq$  is reflexive. Suppose  $x \leq y$  and  $y \leq x$ . Then  $x \circ y = 1$  and  $y \circ x = 1$  hence, by (1) and (2),  $x = 1 * x = (y \circ x) * x = (x \circ y) * y = 1 * y = y$ , thus  $\leq$  is antisymmetrical. Suppose  $x \leq y$  and  $y \leq z$ . Then  $x \circ y = 1$  and  $y \circ z = 1$ . Hence  $x \circ z = x \circ (1*z) = x \circ ((y \circ z)*z) = x \circ (((1*y) \circ z)*z) = x \circ ((((x \circ y)*y) \circ z)*z) = 1$  by (3). Thus  $\leq$  is transitive, i.e., it is a partial order. Also (1) tells us that  $a \circ 1 = 1$  for all a, so 1 is the greatest element.

Define  $a \sqcup b = (a \circ b) * b$ . Put x = y = a, z = b in (3). Then, with (1), we can easily derive

$$a \circ ((a \circ b) * b) = 1.$$

Putting x = y = b, z = a in (3) we obtain analogously (using (1) and (2)) that

$$b \circ ((a \circ b) * b) = 1.$$

Thus  $a \leq a \sqcup b$  and  $b \leq a \sqcup b$ .

Suppose now  $a \leq b$ . Then  $a \sqcup b = (a \circ b) * b = 1 * b = b$ . Clearly,  $a \sqcup b = (a \circ b) * b = (b \circ a) * a = b \sqcup a$ , thus  $(A; \sqcup)$  is a commutative  $\sqcup$ -directoid.

(i)  $\Rightarrow$  (ii): Suppose now that also (4) is satisfied and  $x \in [a, 1], x \leq y$ . Then  $a \leq x$  and thus  $(x \circ a) * a = x \sqcup a = x$ . Therefore from (4) for z = a it follows:  $(((x \circ y) * y) \circ a) \circ (x \circ a) = 1$ , which yields:  $(y \circ a) \circ (x \circ a) = ((1 * y) \circ a) \circ (x \circ a) = (((x \circ y) * y) \circ a) \circ (x \circ a) = 1$ . Thus  $y \circ a \leq x \circ a$ . Analogously we obtain:  $y * a \leq x * a$ , i.e., the condition (5) holds.

Now we consider  $a \in A$ , and  $f_a$ ,  $f_a^{-1}$  defined by (A). Assume  $x \in [a, 1]$ . Then

$$f_a^{-1}(f_a(x)) = (x \circ a) * a = x \sqcup a = x,$$
  
$$f_a(f_a^{-1}(x)) = (x * a) \circ a = x \sqcup a = x,$$

using (3). Since  $x \in [a, 1]$ , by [5], also  $f_a(x) \in [a, 1]$ . Hence  $f_a$  and  $f_a^{-1}$  are bijections on [a, 1] (each being the inverse of the other).

For  $x \in [a, 1]$  and  $x \leq y$  we have by (5):

$$f_a(y) = y \circ a \le x \circ a = f_a(x),$$

$$f_a^{-1}(y) = y * a \le x * a = f_a^{-1}(x),$$

therefore  $f_a$  and  $f_a^{-1}$  are antitone bijections.

(ii)  $\Rightarrow$  (i): By the assumptions  $(A; \sqcup)$  is a commutative  $\sqcup$ -directoid with sectionally antitone bijections. Take  $x, y \in A$ . Since

$$f_y(x \sqcup y) = (x \sqcup y) \circ y = ((x \circ y) * y) \circ y = f_y(f_y^{-1}(x \circ y)) = x \circ y,$$
  
$$f_y^{-1}(x \sqcup y) = (x \sqcup y) * y = ((x * y) \circ y) * y = f_y^{-1}(f_y(x * y)) = x * y,$$

 $\circ$  and  $\ast$  can be also defined by (P). Applying Theorem 1, we obtain that the algebra (A;  $\circ$ ,  $\ast$ , 1) satisfies (4). ■

### Remark 2.

- (i) According to Theorem 1, to any commutative  $\sqcup$ -directoid  $(P; \leq)$  with sectionally antitone bijections it corresponds an algebra  $(P; \circ, *, 1)$  satisfying the axioms (1)-(4).
- (ii) In view of Theorem 2, to any algebra  $\mathcal{A} = (A; \circ, *, 1)$  satisfying the identities (1)-(4) it corresponds a commutative directoid  $(A; \sqcup)$  with sectionally antitone bijections, where  $\leq$  is the induced order of  $(A; \sqcup)$ .

The following example shows, that different algebras satisfying the identities (1)-(4) can be assigned to the same poset  $(P; \leq)$  with sectionally antitone bijections.

A poset  $(P; \leq)$  is called a *complete bipartite* poset if there exists two nonempty subset A, B of P such that  $A \cap B = \emptyset, A \cup B = P$  and such that for any  $x, y \in P$ 

$$x \leq y$$
 if and only if either  $x = y$  or  $x \in A$  and  $y \in B$ .

**Example 1.** Let us consider the finite poset  $P \cup \{1\}$ , where  $(P; \leq)$  is a complete bipartite poset and 1 is an element with the property  $x \leq 1$ , for all  $x \in P$ . Denote the sets A and B as follows:  $A = \{a_1, a_2, \ldots, a_n\}$ ,  $B = \{b_1, b_2, \ldots, b_m\}$ . Clearly  $P \cup \{1\} = \{a_1, a_2, \ldots, a_n, b_1, b_2, \ldots, b_m, 1\}$  and  $a_i < b_j < 1$  for all  $1 \leq i \leq n, 1 \leq j \leq m$ , see Figure 1. We show that  $(P \cup \{1\}; \leq)$  is a poset with antitone bijections. Take an arbitrary permutation  $\pi$  of the set  $\{b_1, b_2, \ldots, b_m\}$  and let us define an arbitrary bijection for each section [p, 1], (where  $p \in P \cup \{1\}$ ) as follows:

$$f_1(1) = 1,$$
  
 $f_{b_j}(1) = b_j, \quad f_{b_j}(b_j) = 1, \quad 1 \le j \le m,$ 

$$f_{a_i}(1) = a_i, \quad f_{a_i}(a_i) = 1, \quad 1 \le i \le n$$

and  $f_{a_i}(b_j) = \pi(b_j)$  for all  $1 \le i \le n$  and  $1 \le j \le m$ .

Further, a commutative directoid  $(P \cup \{1\}; \sqcup)$  can be defined on the set  $P \cup \{1\}$  as follows:

Fix an arbitrary element  $b_k$   $(1 \le k \le m)$ . Then define  $\sqcup$  in the following way:

$$x \sqcup y = y \quad \text{iff} \quad x \le y,$$

for any  $b_i, b_j, b_i \neq b_j$  let  $b_i \sqcup b_j = 1$ , for any  $a_i, a_j, a_i \neq a_j$  let  $a_i \sqcup a_j = b_k$ .

According to Remark 1,  $(P \cup \{1\}, \sqcup)$  is a commutative directoid with sectionally antitone bijections.



Figure 1

In view of Theorem 1, to any permutation  $\pi$  on the set  $\{b_1, \ldots, b_m\}$  and to any  $b_k \in \{b_1, b_2, \ldots, b_m\}$  there corresponds an algebra  $\mathcal{A}$  satisfying the identities (1)–(4). In fact we have  $a_i \circ a_i = b_i \circ b_i = a_i \circ b_j = b_j \circ 1 = 1 \circ 1 = 1$ and  $a_i * a_i = b_i * b_i = a_i * b_j = b_j * 1 = 1 * 1 = 1$  for all  $i \neq j$  such that  $1 \leq i \leq n$  and  $1 \leq j \leq m$ . Further, we have  $b_i \circ b_j = b_i * b_j = b_j$ , for all  $i \neq j$ such that  $1 \leq i, j \leq m, a_i \circ a_j = \pi(b_k), a_i * a_j = \pi^{-1}(b_k)$ , for all  $1 \leq i, j \leq n$ and  $b_j \circ a_i = \pi(b_j), b_j * a_i = \pi^{-1}(b_j)$ , for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

Let us recall that an algebra  $(L; \sqcup, \sqcap)$  of type (2, 2) is called a  $\lambda$ -lattice if its both reducts  $(L; \sqcup)$  and  $(L; \sqcap)$  are commutative directoids and hence the operations are connected by the absorption laws, see e.g. [11]. **Corollary 1.** Let  $\mathcal{A} = (A; \circ, *, 1)$  be an algebra of type (2, 2, 0) satisfying the identities (1), (2), (3) and (4) and let  $\leq$  be the induced order. Then  $(A; \leq)$  is a commutative  $\sqcup$ -directoid with 1, where  $x \sqcup y = (x \circ y) * y$  and for each  $a \in A$  the section [a, 1] becomes a  $\lambda$ -lattice where

$$x \sqcap_a y = (((x \circ a) \circ (y \circ a)) * (y \circ a)) * a.$$

**Proof.** In view of Theorem 2 (ii), the section [a, 1] is a commutative  $\sqcup$ directoid in which  $x \sqcup y = (x \circ y) * y$ . Since  $f_a, f_a^{-1}$  are antitone bijections it follows that letting  $x \sqcap_a y = f_a^{-1}(f_a(x) \sqcup f_a(y))$  we get a commutative  $\sqcap$ -directoid with the same order. Hence  $([a, 1]; \sqcup, \sqcap_a)$  is a  $\lambda$ -lattice.

We have

$$x \sqcap_{a} y = f_{a}^{-1}(f_{a}(x) \sqcup f_{a}(y)) = ((x \circ a) \sqcup (y \circ a)) * a =$$
$$= (((x \circ a) \circ (y \circ a)) * (y \circ a)) * a.$$

When  $\mathcal{A} = (A; \sqcup, \sqcap)$  is a  $\lambda$ -lattice then the induced operation  $\sqcap_a$  in a section [a, 1] need not coincide with the operation  $\sqcap$ , see e.g. the following:

**Example 2.** Let  $\mathcal{L} = (L; \sqcup, \sqcap)$ ,  $L = \{0, w, z, y, x, 1\}$  be a  $\lambda$ -lattice depicted in Figure 2. Pick  $w = x \sqcap y$  and  $z \sqcup w = x$ .





Further, let  $f_z(1) = z$ ,  $f_z(z) = 1$ ,  $f_z(x) = x$ ,  $f_z(y) = y$  and  $f_z^{-1} = f_z$ . Then  $x = x \circ z$ ,  $y = y \circ z$ ,  $z = (x \sqcup y) \circ z = (x \circ z) \sqcap_z (y \circ z)$ , i.e.,  $x \sqcap_z y = z$  but  $w = x \sqcap y = (x \circ z) \sqcap (y \circ z)$ , thus  $\sqcap_z$  is not the same as  $\sqcap$ .

An *implication algebra* [1] is an algebra  $(A; \circ)$  satisfying the identities

- (I1)  $(x \circ y) \circ x = x;$
- (I2)  $(x \circ y) \circ y = (y \circ x) \circ x;$
- (I3)  $x \circ (y \circ z) = y \circ (x \circ z).$

It is well-known that any implication algebra  $(A; \circ)$  contains an element  $1 \in A$  such that  $x \circ x = 1$ , for all  $x \in A$  (see e.g. [1]). The algebra  $(A; \circ, \circ, 1)$  of type (2,2,0) where the operation  $\circ$  is doubled will be called a *double implication algebra*.

It is easy to see that any double implication algebra satisfies the identities (1)-(4).

Indeed, for the algebra  $\mathcal{A} = (A; \circ, \circ, 1)$  the identity (2) is the same as (I2) and we have  $x \circ x = 1$ ,  $1 \circ x = x$  and  $x \circ 1 = (1 \circ x) \circ (1 \circ 1) =$  $1 \circ ((1 \circ x) \circ 1) = 1 \circ 1 = 1$ , i.e., (1) is also satisfied by  $\mathcal{A}$ . Clearly also (3) holds. According to [1], (R) defines a partial order  $\leq$  with the property (5) on any implication algebra  $(A; \circ)$ . Hence, by Theorem 2, we obtain that  $(A; \circ, \circ, 1)$  satisfies the identity (4), as well.

**Example 3.** Observe that for a two-element chain  $(\{0,1\};\leq)$  the algebra  $S_2 = (\{0,1\};\circ,\circ,1)$  (where, of course,  $f_0(0) = 1$ ,  $f_0(1) = 0$ ) is a double implication algebra. The operation  $\circ$  is given by the table

0	0	1
0	1	1
1	0	1.

It is proved in [10] that the implication algebras form a minimal quasivariety which is generated by the two-element implication algebra  $(\{0,1\};\circ)$ . Hence it is not hard to see that the variety generated by the algebra  $S_2$  is also a minimal quasivariety and it coincides with the variety of all double implication algebras. **Proposition 2.** The variety  $\mathcal{V}$  of the algebras  $(A; \circ, *, 1)$  of type (2, 2, 0) satisfying (1)–(4) contains a single minimal quasivariety, namely the variety of double implication algebras.

**Proof.** Let  $\mathcal{W}$  be a nontrivial subquasivariety of  $\mathcal{V}$  and  $\mathcal{A} = (A; \circ, *, 1)$ a nontrivial algebra in  $\mathcal{W}$ . In view of Theorem 1, the corresponding poset  $(A; \leq)$  is a commutative  $\sqcup$ -directoid with sectionally antitone bijections. As  $|\mathcal{A}| \geq 2$ , there exists an element  $a \in A$  with  $a \neq 1$ . In view of Theorem 1 we have  $a \circ a = a * a = 1 \circ 1 = 1 * 1 = 1$ ,  $a \circ 1 = a * 1 = 1$  and  $1 \circ a = 1 * a = a$ . Hence  $(\{a, 1\}; \circ, *, 1)$  is a subalgebra of  $(A; \circ, *, 1)$ . Since  $(\{a, 1\}; \leq)$  is a two-element chain, the algebra  $\mathcal{S}_2 = (\{a, 1\}; \circ, *, 1)$  is a double implication algebra with two elements (see Example 3). Denote the variety generated by it as  $\mathcal{V}_2$ . We already shown that  $\mathcal{V}_2$  is the variety of all double implication algebras. Since  $\mathcal{V}_2$  is a minimal quasivariety as well, we have  $\mathcal{V}_2 = Q(\mathcal{S}_2)$ , where  $Q(\mathcal{S}_2)$  denotes the quasivariety generated by the algebra  $\mathcal{S}_2$ . As  $\mathcal{S}_2 \in \mathcal{W}$  we get  $\mathcal{V}_2 = Q(\mathcal{S}_2) \subseteq \mathcal{W}$ , and this proves that  $\mathcal{V}_2$  is the unique minimal quasivariety contained in variety  $\mathcal{V}$ .

Let  $\mathcal{A} = (A; \circ, *, 1)$  be an algebra of type (2,2,0). A nonempty subset  $K \subseteq A$ is called a *congruence kernel* of  $\mathcal{A}$  if  $K = [1]_{\Theta} = \{x \in A; (x, 1) \in \Theta\}$  for some congruence  $\Theta$  of  $\mathcal{A}$ . Recall that  $\mathcal{A}$  is called 3-*permutable* if  $\Theta_1 \circ \Theta_2 \circ \Theta_1 =$  $\Theta_2 \circ \Theta_1 \circ \Theta_2$  holds for every  $\Theta_1, \Theta_2 \in Con\mathcal{A}$ . According to J. Hagemann and A. Mitschke [7], a variety  $\mathcal{V}$  of algebras is 3-permutable if and only if there exist ternary terms  $p_0, p_1, p_2$  and  $p_3$  in  $\mathcal{V}$  such that the following identities hold in  $\mathcal{V}$ :

(B) 
$$\begin{cases} p_0(x, y, z) = x, \ p_3(x, y, z) = z, \\ p_i(x, x, y) = p_{i+1}(x, y, y) \text{ for } i \in \{0, 1, 2\} \end{cases}$$

An algebra  $\mathcal{A}$  with a constant 1 is called *weakly regular* if every  $\Theta \in Con\mathcal{A}$  is determined by its kernel, i.e., if  $[1]_{\Theta} = [1]_{\Phi}$  implies  $\Theta = \Phi$  for every  $\Phi, \Theta \in Con\mathcal{A}$ . A variety  $\mathcal{V}$  is weakly regular if every algebra  $\mathcal{A} \in \mathcal{V}$  has this property. The following characterization of weakly regular varieties was established by B. Csákány in [6].

**Proposition 3** [6]. A variety  $\mathcal{V}$  with 1 is weakly regular if and only if there exists  $n \in \mathbb{N}$  and binary terms  $q_1(x, y), q_2(x, y), \ldots, q_n(x, y)$  such that

(C) 
$$q_1(x,y) = q_2(x,y) = \dots = q_n(x,y) = 1 \Leftrightarrow x = y$$

is satisfied for every algebra  $\mathcal{A} \in \mathcal{V}$ .

For other congruence conditions, the reader is asked to consult e.g. [3].

**Theorem 3.** The variety  $\mathcal{V}$  of the algebras  $(A; \circ, *, 1)$  of type (2, 2, 0) satisfying the identities (1)-(4) is weakly regular, 3-permutable, arithmetical at 1 and congruence distributive.

**Proof.** Consider the terms  $q_1(x, y) = x \circ y$  and  $q_2(x, y) = y \circ x$ . Then  $q_1(x, x) = q_2(x, x) = x \circ x = 1$ . If  $q_1(x, y) = 1$  and  $q_2(x, y) = 1$ , then by (R) we have  $x \leq y$  and  $y \leq x$  thus x = y. In view of Proposition 3, we conclude that  $\mathcal{V}$  is weakly regular (at 1).

Now, let  $p_0(x, y, z) = x$ ,  $p_3(x, y, z) = z$  and  $p_1(x, y, z) = (z * y) \circ x$ ,  $p_2(x, y, z) = (x * y) \circ z$ . It is easy to see that these terms satisfy the identities (B). Consequently,  $\mathcal{V}$  is congruence 3-permutable.

It is proved in [3] that a variety is arithmetical at 1 if and only if there exists a binary term b(x, y) of it such that b(x, x) = b(1, x) = 1 and b(x, 1) = x. Obviously, we can take  $b(x, y) = y \circ x$ .

Since  $\mathcal{V}$  is arithmetical at 1 it is congruence distributive at 1, i.e.,  $[1]_{\Theta \cap (\Phi \lor \Psi)} = [1]_{(\Theta \cap \Phi) \lor (\Theta \cap \Psi)}$ , for all  $\Theta, \Phi, \Psi \in Con\mathcal{A}$  for  $\mathcal{A} \in \mathcal{V}$ .

As  $\mathcal{V}$  is weakly regular, this equality implies  $\Theta \cap (\Phi \lor \Psi) = (\Theta \cap \Phi) \lor (\Theta \cap \Psi)$  (for all  $\Theta, \Phi, \Psi \in Con\mathcal{A}$  and  $\mathcal{A} \in \mathcal{V}$ ), thus the variety  $\mathcal{V}$  is congruence distributive.

#### References

- [1] J.C. Abbott, Semi-Boolean algebras, Matem. Vestnik 4 (1967), 177–198.
- [2] I. Chajda, Lattices and semilattices having an antitone bijection in any upper interval, Comment. Math. Univ. Carolinae 44 (2003), 577–585.
- [3] I. Chajda, G. Eigenthaler and H. Länger, Congruence Classes in Universal Algebra, Heldermann Verlag, Lemgo (Germany), 2003, ISBN 3-88538-226-1.
- [4] I. Chajda and M. Kolařík, Directoids with sectionally antitone involutions and skew MV-algebras, Math. Bohemica 132 (2007), 407–422.
- [5] I. Chajda and R. Radeleczki, Semilattices with sectionally antitone bijections, Novi Sad J. Math. 35 (2005), 93–101.

- [6] B. Csákány, Characterization of regular varieties, Acta Sci. Math. Szeged 31 (1970), 187–189.
- [7] J. Hagemann and A. Mitschke, On n-permutable congruences, Algebra Universalis 3 (1973), 8–12.
- [8] J. Ježek and R. Quackenbush, Directoids: algebraic models of up-directed sets, Algebra Universalis 27 (1990), 49–69.
- [9] V.M. Kopytov and Z.I. Dimitrov, On directed groups, Siberian Math. J. 30 (1989), 895–902. (Russian original: Sibirsk. Mat. Zh. 30 (6) (1988), 78–86.)
- [10] S. Radeleczki, The congruence lattice of implication algebras, Math. Pannonica 3 (1992), 115–123.
- [11] V. Snášel,  $\lambda$ -lattices, Math. Bohemica **122** (1997), 267–272.

Received 5 March 2007 Revised 27 March 2007