

REMARKS ON PSEUDO MV-ALGEBRAS*

IVAN CHAJDA AND MIROSLAV KOLAŘÍK

Department of Algebra and Geometry
Palacký University Olomouc
Tomkova 40, 779 00 Olomouc, Czech Republic

e-mail: chajda@inf.upol.cz

e-mail: kolarik@inf.upol.cz

Abstract

Pseudo MV-algebras (see e.g., [4, 6, 8]) are non-commutative extension of MV-algebras. We show that every pseudo MV-algebra is isomorphic to the algebra of action functions where the binary operation is function composition, zero is $x \wedge y$ and unit is x . Then we define the so-called difference functions in pseudo MV-algebras and show how a pseudo MV-algebra can be reconstructed by them.

Keywords: pseudo MV-algebra, action function, guard function, difference functions.

2000 Mathematics Subject Classification: 06D35, 08A02, 03G25.

1. INTRODUCTION

In 1958, C.C. Chang introduced the notion of MV-algebra as an algebraic counterpart of the Łukasiewicz propositional calculus.

Non-commutative MV-algebras, named pseudo MV-algebras, were introduced by G. Georgescu and A. Iorgulescu in [8].

*This work is supported by the Research and Development Council of the Czech Government via the project MSM6198959214.

Independently, starting from intervals of (not necessarily commutative) lattice-ordered groups, J. Rachůnek established in [12] the concept of a GMV-algebra (generalized MV-algebra).

Definition 1. A *Pseudo MV-algebra* is an algebra $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ of type $(2, 1, 1, 0, 0)$ satisfying the following axioms:

- (A1) $x \oplus (y \oplus z) = (x \oplus y) \oplus z$;
- (A2) $x \oplus 0 = x = 0 \oplus x$;
- (A3) $x \oplus 1 = 1 = 1 \oplus x$;
- (A4) $\neg 1 = 0 = \sim 1$;
- (A5) $\neg(\sim x \oplus \sim y) = \sim(\neg x \oplus \neg y)$;
- (A6) $x \oplus (y \odot \sim x) = y \oplus (x \odot \sim y) = (\neg y \odot x) \oplus y = (\neg x \odot y) \oplus x$;
- (A7) $\sim \neg x = x$

where the additional operation \odot is defined via

$$x \odot y = \sim(\neg x \oplus \neg y).$$

Note that this structure can be defined also (in term equivalent way) as an FL_ω -algebra that satisfies $x/(y \setminus x) = x \vee y = (x/y) \setminus x$; see [10].

If \oplus is commutative, then \sim coincides with \neg and $(A; \oplus, \neg, 0, 1)$ becomes an MV-algebra. For basic properties of MV-algebras and pseudo MV-algebras we refer to [5, 8, 12].

Remark 1. The definition of a pseudo MV-algebra usually contains the following axiom (see e.g., [4, 8, 12]):

$$(A8) \quad y \odot (x \oplus \sim y) = (\neg x \oplus y) \odot x.$$

However, F. Švrček shows recently in an unpublished preprint [11] that (A8) is redundant. More precisely, everyone can derive (A8) using the axioms of Definition 1 as follows.

We have

$$\begin{aligned} y \odot (x \oplus \sim y) &= \sim \neg y \odot (\sim \neg x \oplus \sim \sim \neg y) \\ &= \sim \neg y \odot \sim (\neg x \odot \sim \neg y) = \sim (\neg y \oplus (\neg x \odot \sim \neg y)). \end{aligned}$$

Further, by (A6), we obtain

$$\begin{aligned} \sim (\neg y \oplus (\neg x \odot \sim \neg y)) &= \sim ((\neg \neg x \odot \neg y) \oplus \neg x) \\ &= \sim (\neg \neg x \odot \neg y) \odot \sim \neg x = (\sim \neg \neg x \oplus \sim \neg y) \oplus x = (\neg x \oplus y) \odot x. \end{aligned}$$

Proposition 1 (see e.g., [8, 12]). *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra, let*

$$x \odot y = \sim (\neg x \oplus \neg y).$$

Then the following identities hold:

- (a) $\neg \sim x = x$;
- (b) $(x \odot y) \odot z = x \odot (y \odot z)$;
- (c) $x \odot \sim x = 0 = \neg x \odot x$;
- (d) $x \odot 0 = 0 = 0 \odot x$, $x \odot 1 = x = 1 \odot x$;
- (e) $\neg 0 = 1 = \sim 0$.

Proposition 2 (see [8]). *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra. For term functions*

$$x \vee y = x \oplus \sim (\neg y \oplus x) = \neg(x \oplus \sim y) \oplus y,$$

$$x \wedge y = x \odot \sim (\neg y \odot x) = \neg(x \odot \sim y) \odot x,$$

the algebra $\mathcal{L}(A) = (A; \vee, \wedge, 0, 1)$ is a bounded distributive lattice and the mappings $x \mapsto \neg x$ and $x \mapsto \sim x$ are mutually inverse antitone bijections on A .

The lattice order of $\mathcal{L}(A)$ is called the *induced order* of a given pseudo MV-algebra \mathcal{A} . Of course,

$$x \leq y \quad \text{iff} \quad \neg x \oplus y = 1 \quad \text{iff} \quad y \oplus \sim x = 1.$$

Proposition 3 (see [4]). *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra. For each $a \in A$, the mapping $f_a(x) = \neg x \oplus a$ is an antitone bijection on the interval $[a, 1]$ and its inverse mapping is given by $f_a^{-1}(x) = a \oplus \sim x$. Moreover, $x \oplus y = f_x^{-1}(\neg y \vee x)$.*

Note that the first part of proposition is obvious as $f_a(x) = x \setminus a$, $f_a^{-1}(x) = a/x$ and $a/(x \setminus a) = a \vee x = (a/x) \setminus a$.

2. A REPRESENTATION OF PSEUDO MV-ALGEBRAS BY MEANS OF BINARY FUNCTIONS

Cayley's Theorem provides a well-known representation of groups by means of certain unary functions (i.e. permutations) with composition as its binary operation. It was shown by W.C. Holland [9] that also ℓ -groups can be represented by monotonous permutations. A representation of Boolean algebras by binary functions, the so-called "guard functions" was presented in [1]. Also MV-algebras were represented by binary functions which are an extended version of action functions (see [3]). The aim of this section is to give a representation of pseudo MV-algebras by certain binary functions which can be considered as action functions.

At first we define an action function on pseudo MV-algebra.

By an *action* we mean any process which starts whenever certain conditions are satisfied. Formally, if a is a condition and x a process, by an action we mean the proposition

$$\text{IF } a = 1 \text{ DO } x, \quad \text{IF } a = 0 \text{ DO NOTHING.}$$

Consider a binary function

$$h_a(x, y) = x \wedge (a \oplus y)$$

defined on a pseudo MV-algebra $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$. One can easily see that

$$h_1(x, 0) = x \wedge (1 \oplus 0) = x$$

and

$$h_0(x, 0) = x \wedge (0 \oplus 0) = 0.$$

Hence, $h_a(x, 0)$ describes the aforementioned concept of action and hence $h_a(x, y)$ will be called an *action function* (cf. [3]).

For two action functions on \mathcal{A} we define the following composition

$$(h_a \circ h_b)(x, y) = h_a(x, h_b(x, y)).$$

Of course, $h_a(x, x) = x$ for each $a \in A$, thus this composition can be read as $(h_a \circ h_b)(x, y) = h_a(h_b(x, x), h_b(x, y))$.

The following result can be found in [6]. Note that it actually holds for all involutive residuated lattices (for details see [7]).

Lemma 1. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra, let $a, b, c \in A$. Then*

$$a \oplus (b \wedge c) = (a \oplus b) \wedge (a \oplus c).$$

Lemma 2. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra and h_a, h_b be action functions on \mathcal{A} . Then $h_a \circ h_b = h_{a \oplus b}$.*

Proof. By Theorem 3 in [12], $0 \leq a$ implies $x = 0 \oplus x \leq a \oplus x$. According to this fact and the previous definitions and Lemma 1, we have:

$$(h_a \circ h_b)(x, y) = h_a(x, h_b(x, y)) = x \wedge (a \oplus (x \wedge (b \oplus y)))$$

$$= x \wedge (a \oplus x) \wedge (a \oplus b \oplus y) = x \wedge (a \oplus b \oplus y) = h_{a \oplus b}(x, y).$$

■

Theorem 1. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra and $H(A) = \{h_a(x, y); a \in A\}$ the set of all action functions on \mathcal{A} . Define*

$$\neg h_a(x, y) = h_{\neg a}(x, y), \quad \sim h_a(x, y) = h_{\sim a}(x, y).$$

Then $\mathcal{H} = (H(A); \circ, \neg, \sim, x \wedge y, x)$ is a pseudo MV-algebra which is isomorphic to \mathcal{A} via $h_a(x, y) \mapsto a$.

Proof. We compute $h_0(x, y) = x \wedge (0 \oplus y) = x \wedge y$ and $h_1(x, y) = x \wedge (1 \oplus y) = x$. By the previous results the map $a \mapsto h_a$ is clearly a homomorphism of a pseudo MV-algebra \mathcal{A} onto $H(A)$ and, therefore, \mathcal{H} is also a pseudo MV-algebra. Moreover, if $h_a = h_b$ then $x \wedge (a \oplus y) = x \wedge (b \oplus y)$ for all x, y of \mathcal{A} . In particular, for $x = 1$ and $y = 0$ we obtain $a = b$, i.e. this mapping is one-to-one and hence \mathcal{H} is isomorphic to \mathcal{A} . ■

Similarly as for Boolean algebras [1], for q -algebras [2] and for MV-algebras [3], we can define *guard functions* $g_a(x, y)$ on a pseudo MV-algebra by

$$g_a(x, y) = (a \oplus x) \wedge (\neg a \oplus y).$$

The meaning of this is as follows. The element a is considered as a "condition" similarly as for action functions. Hence, if $a = 0$ then

$$g_0(x, y) = (0 \oplus x) \wedge (1 \oplus y) = x \wedge 1 = x$$

and if $a = 1$ then

$$g_1(x, y) = (1 \oplus x) \wedge (0 \oplus y) = 1 \wedge y = y.$$

Thus $g_a(x, y)$ makes decisions on the action x or y in dependency of the value of a . Of course, in many-valued logic we have also another values of a distinct from 0 and 1 and hence this guard function is much more complex. However, guard functions can be constructed by means of action functions as follows

$$g_a(x, y) = h_0(h_a(1, x), h_{\neg a}(1, y)).$$

On the contrary, action functions are not expressible in guard functions. Nevertheless, we can use a guard function to express an action function and its negations \neg , \sim in the function algebra \mathcal{H} as follows

$$h_a(x, y) = h_0(x, g_a(y, 1))$$

and

$$\neg h_a(x, y) = h_0(x, g_a(1, y)), \quad \sim h_a(x, y) = h_0(x, g_{\sim a}(1, y)).$$

Of course, we can define another sort of guard functions by

$$k_a(x, y) = (a \oplus x) \wedge (\sim a \oplus y)$$

whose behavior is similar.

3. DIFFERENCE FUNCTIONS IN PSEUDO MV-ALGEBRAS

The concept of difference function was already introduced for MV-algebras. Due to the fact that the binary operation of pseudo MV-algebra is not commutative and there are two unary operations \neg and \sim , we should define two difference functions as follows.

Definition 2. Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra. Define so-called *difference functions* $+_1, +_2$ as follows:

$$x +_1 y = (x \odot \sim y) \oplus (y \odot \sim x),$$

$$x +_2 y = (\neg y \odot x) \oplus (\neg x \odot y).$$

Of course, $+_1, +_2$ need not be commutative since \oplus does not have this property either. By (c) of Proposition 1 and (A2), the difference functions satisfy the expected properties:

$$x +_1 x = 0, \quad x +_2 x = 0.$$

Now, we give some basic properties of difference functions.

Lemma 3. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra and $+_1, +_2$ difference functions. Then*

- (i) $\neg x +_1 \neg y = y +_2 x$;
- (ii) $\sim x +_2 \sim y = y +_1 x$;
- (iii) $x +_1 y = 0 \Leftrightarrow x = y$;
- (iv) $x +_2 y = 0 \Leftrightarrow x = y$.

Proof.

$$(i): \neg x +_1 \neg y = (\neg x \odot \sim \neg y) \oplus (\neg y \odot \sim \neg x) = (\neg x \odot y) \oplus (\neg y \odot x) = y +_2 x.$$

$$(ii): \sim x +_2 \sim y = (\neg \sim y \odot \sim x) \oplus \neg \sim x \odot \sim y = (y \odot \sim x) \oplus (x \odot \sim y) = y +_1 x.$$

(iii): If $x = y$ then $x +_1 x = 0$. Conversely, let $x +_1 y = 0$. Then

$$(x \odot \sim y) \oplus (y \odot \sim x) = 0,$$

thus (according to Theorem 3 in [12])

$$(x \odot \sim y) \vee (y \odot \sim x) \leq (x \odot \sim y) \oplus (y \odot \sim x) = 0,$$

whence $x \odot \sim y = 0$ and $y \odot \sim x = 0$. By Theorem 5 in [12], $x \leq y$ and $y \leq x$, i.e. $x = y$.

(iv): In the same way as (iii).

■

Moreover, we can express the operation \oplus by means of $+_1$, $+_2$ and \odot as follows.

Lemma 4. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra and $+_1$, $+_2$ difference functions. Then $1 +_1 x = \sim x$, $x +_2 1 = \neg x$ and*

$$x \oplus y = (1 +_1 x) \odot (1 +_1 y) +_2 1.$$

Proof.

$$1 +_1 x = (1 \odot \sim x) \oplus (x \odot \sim 1) = \sim x \oplus 0 = \sim x,$$

$$x +_2 1 = (\neg 1 \odot x) \oplus (\neg x \odot 1) = 0 \oplus \neg x = \neg x.$$

Further,

$$(1 +_1 x) \odot (1 +_1 y) +_2 1 = (\sim x \odot \sim y) +_2 1$$

$$= \neg(\sim x \odot \sim y) = \neg \sim x \oplus \neg \sim y = x \oplus y.$$

■

Our next task is to set up axioms characterizing these difference functions and, further, to show that also conversely, a pseudo MV-algebra can be reconstructed by means of these difference functions and the binary operation \odot .

Theorem 2. *Let $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ be a pseudo MV-algebra and $+_1$, $+_2$ be the difference functions. Then the following identities (D1)–(D5), (M1)–(M3) are satisfied:*

$$(D1) \quad ((1 +_1 x) \odot (1 +_1 y) +_2 1) +_2 1 = (x +_2 1) \odot (y +_2 1),$$

$$1 +_1 ((1 +_1 x) \odot (1 +_1 y) +_2 1) = (1 +_1 x) \odot (1 +_1 y);$$

$$(D2) \quad (1 +_1 x) +_2 1 = x, \quad 1 +_1 (x +_2 1) = x;$$

$$(D3) \quad 1 +_1 1 = 0, \quad 1 +_2 1 = 0;$$

$$(D4) \quad 0 +_1 x = x = x +_1 0, \quad x +_2 0 = x = 0 +_2 x;$$

$$\begin{aligned} (D5) \quad & (1 +_1 x) \odot (1 +_1 (y \odot (1 +_1 x))) +_2 1 = (1 +_1 y) \odot (1 +_1 (x \odot (1 +_1 y))) +_2 1 \\ & = (1 +_1 ((y +_2 1) \odot x) \odot (1 +_1 y)) +_2 1 = (1 +_1 ((x +_2 1) \odot y) \\ & \odot (1 +_1 x)) +_2 1; \end{aligned}$$

$$(M1) \quad 1 \odot x = x = x \odot 1;$$

$$(M2) \quad 0 \odot x = 0 = x \odot 0;$$

$$(M3) \quad (x \odot y) \odot z = x \odot (y \odot z).$$

Proof.

$$(D1): \quad ((1 +_1 x) \odot (1 +_1 y) +_2 1) +_2 1 = \neg(\neg(\sim x \odot \sim y))$$

$$= \neg(\neg(\sim (\neg \sim x \oplus \neg \sim y))) \stackrel{(a)}{=} \neg(x \oplus y) \stackrel{(A7)}{=} \neg(\sim \neg x \oplus \sim \neg y)$$

$$\stackrel{(A5)}{=} \sim (\neg \neg x \oplus \neg \neg y) = \neg x \odot \neg y = (x +_2 1) \odot (y +_2 1),$$

$$1 +_1 ((1 +_1 x) \odot (1 +_1 y) +_2 1) = \sim (\neg(\sim x \odot \sim y))$$

$$\stackrel{(A7)}{=} \sim x \odot \sim y = (1 +_1 x) \odot (1 +_1 y);$$

$$(D2): \quad (1 +_1 x) +_2 1 = \neg \sim x \stackrel{(a)}{=} x,$$

$$1 +_1 (x +_2 1) = \sim \neg x = x \quad \text{by (A7);}$$

$$(D3): \quad 1 +_1 1 = (1 \odot \sim 1) \oplus (1 \odot \sim 1) \stackrel{(c)}{=} 0 \oplus 0 \stackrel{(A2)}{=} 0,$$

$$1 +_2 1 = (\neg 1 \odot 1) \oplus (\neg 1 \odot 1) \stackrel{(A4)}{=} (0 \odot 1) \oplus (0 \odot 1) \stackrel{(d)}{=} 0 \oplus 0 \stackrel{(A2)}{=} 0;$$

$$(D4): \quad 0 +_1 x = (0 \odot \sim x) \oplus (x \odot \sim 0) \stackrel{(d),(e)}{=} 0 \oplus (x \odot 1) \stackrel{(d)}{=} 0 \oplus x \stackrel{(A2)}{=} x,$$

$$\text{analogously } x +_1 0 = x,$$

$$x +_2 0 = (\neg 0 \odot x) \oplus (\neg x \odot 0) \stackrel{(d),(e)}{=} x \oplus 0 \stackrel{(A2)}{=} x,$$

$$\text{analogously } x +_2 0 = x;$$

(D5): Using Lemma 4 we have:

$$(1 +_1 x) \odot (1 +_1 (y \odot (1 +_1 x))) +_2 1 = x \oplus (y \odot \sim x),$$

$$(1 +_1 y) \odot (1 +_1 (x \odot (1 +_1 y))) +_2 1 = y \oplus (x \odot \sim y),$$

$$(1 +_1 ((y +_2 1) \odot x) \odot (1 +_1 y)) +_2 1 = (\neg y \odot x) \oplus y,$$

$$(1 +_1 ((x +_2 1) \odot y) \odot (1 +_1 x)) +_2 1 = (\neg x \odot y) \oplus x.$$

The rest of the proof of (D5) follows directly by (A6).

(M1), (M2) and (M3) follows immediately by (b) and (d) of Proposition 1.

■

Now, we can prove the converse.

Theorem 3. *Let A be a non-void set, $1 \in A$ and $+_1, +_2, \odot$ be binary operations on A satisfying the identities (D1)–(D5) and (M1), (M2), (M3). Then for*

$$x \oplus y = (1 +_1 x) \odot (1 +_1 y) +_2 1,$$

$$\sim x = 1 +_1 x,$$

$$\neg x = x +_2 1,$$

$$0 = 1 +_1 1,$$

the algebra $\mathcal{A} = (A; \oplus, \neg, \sim, 0, 1)$ is a pseudo MV-algebra.

Proof. We must verify the axioms (A1)–(A7).

$$\begin{aligned} \text{(A1): } & (x \oplus y) \oplus z = (1 +_1 (x \oplus y)) \odot (1 +_1 z) +_2 1 \\ &= (1 +_1 ((1 +_1 x) \odot (1 +_1 y) +_2 1)) \odot (1 +_1 z) +_2 1 \\ &\stackrel{(D2)}{=} ((1 +_1 x) \odot (1 +_1 y)) \odot (1 +_1 z) +_2 1 \\ &\stackrel{(M3)}{=} (1 +_1 x) \odot ((1 +_1 y) \odot (1 +_1 z)) +_2 1 \\ &\stackrel{(D2)}{=} (1 +_1 x) \odot (1 +_1 ((1 +_1 y) \odot (1 +_1 z) +_2 1)) +_2 1 \\ &= (1 +_1 x) \odot (1 +_1 (y \oplus z)) +_2 1 = x \oplus (y \oplus z). \end{aligned}$$

$$(A2): \quad x \oplus 0 = (1 +_1 x) \odot (1 +_1 0) +_2 1 \stackrel{(D4)}{=}$$

$$\stackrel{(D4)}{=} (1 +_1 x) \odot 1 +_2 1 \stackrel{(M1)}{=} (1 +_1 x) +_2 1 \stackrel{(D2)}{=} x,$$

analogously $0 \oplus x = x$.

$$(A3): \quad x \oplus 1 = (1 +_1 x) \odot (1 +_1 1) +_2 1 \stackrel{(D3)}{=}$$

$$\stackrel{(D3)}{=} (1 +_1 x) \odot 0 +_2 1 \stackrel{(M2)}{=} 0 +_2 1 \stackrel{(D4)}{=} 1,$$

analogously $1 \oplus x = 1$.

$$(A4): \quad \neg 1 = 1 +_2 1 \stackrel{(D3)}{=} 0, \quad \sim 1 = 1 +_1 1 \stackrel{(D3)}{=} 0.$$

$$(A5): \quad \neg(\sim x \oplus \sim y) = ((1 +_1 x) \oplus (1 +_1 y)) +_2 1$$

$$= ((1 +_1 (1 +_1 x)) \odot (1 +_1 (1 +_1 y)) +_2 1) +_2 1 \stackrel{(D1)}{=}$$

$$\stackrel{(D1)}{=} ((1 +_1 x) +_2 1) \odot ((1 +_1 y) +_2 1) \stackrel{(D2)}{=} x \odot y \stackrel{(D2)}{=}$$

$$\stackrel{(D2)}{=} (1 +_1 (x +_2 1)) \odot (1 +_1 (y +_2 1)) \stackrel{(D1)}{=}$$

$$\stackrel{(D1)}{=} 1 +_1 ((1 +_1 (x +_2 1)) \odot (1 +_1 (y +_2 1)) +_2 1)$$

$$= 1 +_1 ((x +_2 1) \oplus (y +_2 1)) = \sim (\neg x \oplus \neg y).$$

(A6): Follows directly by (D5).

(A7): $\sim \neg x = 1 +_1 (x +_2 1) \stackrel{(D2)}{=} x.$ ■

REFERENCES

- [1] S.L. Bloom, Z. Ésik and E. Manes, *A Cayley theorem for Boolean algebras*, Amer. Math. Monthly **97** (1990), 831–833.
- [2] I. Chajda, *A representation of the algebra of quasiordered logic by binary functions*, Demonstratio Mathem. **27** (1994), 601–607.
- [3] I. Chajda and H. Länger, *Action algebras*, Italian J. of Pure Appl. Mathem., submitted.
- [4] I. Chajda and J. Kühr, *Pseudo MV-algebras and meet-semilattices with sectionally antitone permutations*, Math. Slovaca **56** (2006), 275–288.
- [5] R.L.O. Cignoli, I.M.L. D'Ottaviano and D. Mundici, *Algebraic Foundations of Many-Valued Reasoning*, Kluwer Acad. Publ., Dordrecht-Boston-London 2000.
- [6] A. Dvurečenskij, *Pseudo MV-algebras are intervals in ℓ -groups*, J. Aust. Math. Soc. **72** (3) (2002), 427–445.
- [7] N. Galatos, P. Jipsen, T. Kowalski and H. Ono, *Residuated Lattices*, An Algebraic Glimpse at Substructural Logics, Elsevier 2007.
- [8] G. Georgescu and A. Iorgulescu, *Pseudo MV-algebras*, Multiple Valued Log. **6** (2001), 95–135.
- [9] A.M.W Glass and W.C. Holland, *Lattice-Ordered Groups*, Kluwer Acad. Publ., Dordrecht-Boston-London 1989.
- [10] P. Jipsen and C. Tsınakis, *A survey of Residuated Lattices, Ordered Algebraic Structures* (Martinez J., editor), Kluwer Academic Publishers, Dordrecht, 2002, 19–56.
- [11] J. Kühr and F. Švrček, *Operators on unital ℓ -groups*, preprint, 2007.

- [12] J. Rachůnek, *A non-commutative generalization of MV-algebras*, Czechoslovak Math. J. **52** (2002), 255–273.

Received 19 October 2007

Revised 1 April 2008