

HYPERIDENTITIES IN MANY-SORTED ALGEBRAS

KLAUS DENECKE AND SOMSAK LEKKOKSUNG

Universität Potsdam, Institut of Mathematics
Am Neuen Palais, 14415 Potsdam, Germany

e-mail: kdenecke@rz.uni-potsdam.de

Abstract

The theory of hyperidentities generalizes the equational theory of universal algebras and is applicable in several fields of science, especially in computers sciences (see e.g., [2, 1]). The main tool to study hyperidentities is the concept of a hypersubstitution. Hypersubstitutions of many-sorted algebras were studied in [3]. On the basis of hypersubstitutions one defines a pair of closure operators which turns out to be a conjugate pair. The theory of conjugate pairs of additive closure operators can be applied to characterize solid varieties, i.e., varieties in which every identity is satisfied as a hyperidentity (see [4]). The aim of this paper is to apply the theory of conjugate pairs of additive closure operators to many-sorted algebras.

Keywords: hypersubstitution; hyperidentity; heterogeneous algebra.

2000 Mathematics Subject Classification: 08A68, 08B15.

1. PRELIMINARIES

Hyperidentities in one-based algebras were considered by many authors (for references see e.g., [4, 2]). An identity $s \approx t$ is satisfied as a hyperidentity in the one-based algebra $\mathcal{A} = (A; (f_i^{\mathcal{A}})_{i \in I})$ of type τ if after any

replacements of the operation symbols occurring in s and t by terms of the same arity the arising equation is satisfied in \mathcal{A} . These replacements can be described by hypersubstitutions, i.e., mappings from the set of operation symbols into the set of all terms of type τ . Hypersubstitutions cannot only be applied to terms or equations but also to algebras. This gives a pair of additive closure operators which are related to each other by the so-called conjugate property and which form a conjugate pair of additive closure operators (see [4]). A variety of one-based algebras is called solid if every identity is satisfied as a hyperidentity. Characterizations of solid varieties are based on the theory of conjugate pairs of additive closure operators. For more background see [4].

In this paper we want to apply the theory of conjugate pairs of additive closure operators to many-sorted algebras and identities and want to define hyperidentities and solid varieties of many-sorted algebras.

Many-sorted algebras occur in various branches of mathematics. They have found their way into computer science through abstract data type specifications. Many-sorted algebras, varieties and quasivarieties of many-sorted algebras are the mathematical fundament of approaches to abstract data types in programming and specification languages. For basic concepts on many-sorted algebras we refer the reader to [5].

The concept of terms in many-sorted algebras was discussed in [5]. First we want to give a slightly different version of the definitions and results from [3].

Let I be a non-empty set, let $\mathbb{N}^+ := \mathbb{N} \setminus \{0\}$, $n \in \mathbb{N}^+$, let $I^* := \bigcup_{n \geq 1} I^n$ and $\Sigma \subseteq I^* \times I$. Then we define $\Sigma_n := \Sigma \cap I^{n+1}$. For $\gamma \in \Sigma$ let $\gamma(l)$ denote the l -th component of γ . Let K_γ be a set of indices with respect to γ . If $|K_\gamma| = 1$, we will drop the index.

Definition 1.1. Let $n \in \mathbb{N}^+$ and $X^{(n)} := (X_i^{(n)})_{i \in I}$ be an I -sorted set of variables, also called an n -element I -sorted alphabet, with $X_i^{(n)} := \{x_{i1}, \dots, x_{in}\}$, $i \in I$ and let $((f_\gamma)_k)_{k \in K_\gamma, \gamma \in \Sigma}$ be an indexed set of Σ -sorted operation symbols. Then for each $i \in I$ a set $W_n(i)$ which is called the set of all n -ary Σ -terms of sort i , is inductively defined as follows:

- (i) $W_0^n(i) := X_i^{(n)}$, $i \in I$,

- (ii) $W_{l+1}^n(i) := W_l^n(i) \cup \{f_\gamma(t_{k_1}, \dots, t_{k_n}) \mid \gamma = (k_1, \dots, k_n; i) \in \Sigma, t_{k_j} \in W_l^n(k_j), 1 \leq j \leq n\}, l \in \mathbb{N}$. (Here we inductively assume that the sets $W_l^n(i)$ are already defined for all sorts $i \in I$).

Then $W_n(i) := \bigcup_{l=0}^{\infty} W_l^n(i)$ and we set $W(i) := \bigcup_{n \in \mathbb{N}^+} W_n(i)$. Let $X_i := \bigcup_{n \in \mathbb{N}^+} X_i^{(n)}$ and $X := (X_i)_{i \in I}$. Let $W_\Sigma(X) := (W(i))_{i \in I}$. The set $W_\Sigma(X)$ is called I -sorted set of all Σ -terms and its elements are called I -sorted Σ -terms.

For any $n \in \mathbb{N}^+, i \in I$ we set $\Lambda_n(i) := \{(w; i) \in I^{n+1} \mid w \in I^n, \exists m \in \mathbb{N}^+, \exists \alpha \in \Sigma_m, \exists j (1 \leq j \leq m)(\alpha(j) = i)\}$. Let $\Lambda(i) := \bigcup_{n=1}^{\infty} \Lambda_n(i)$ and we set $\Lambda := \bigcup_{i \in I} \Lambda(i)$.

To define many-sorted hypersubstitutions we need the following superposition operation for I -sorted Σ -terms.

Definition 1.2. Let $t \in W(i), t_j \in W(k_j)$ where $1 \leq j \leq n, n \in \mathbb{N}$. Then the superposition operation

$$S_\beta : W(i) \times W(k_1) \times \dots \times W(k_n) \rightarrow W(i)$$

for $\beta = (k_1, \dots, k_n; i) \in \Lambda$, is defined inductively as follows:

1. If $t = x_{ij} \in X_i$, then

- 1.1 $S_\beta(x_{ij}, t_1, \dots, t_n) := x_{ij}$ for $i \neq k_j$ and

- 1.2 $S_\beta(x_{ij}, t_1, \dots, t_n) := t_j$ for $i = k_j$.

2. If $t = f_\gamma(s_1, \dots, s_m) \in W(i)$ for $\gamma = (i_1, \dots, i_m; i) \in \Sigma$ and $s_q \in W_n(i_q), 1 \leq q \leq m, m \in \mathbb{N}$, and if we assume that $S_{\beta_q}(s_q, t_1, \dots, t_n)$ with $\beta_q = (k_1, \dots, k_n; i_q) \in \Lambda$ are already defined, then $S_\beta(f_\gamma(s_1, \dots, s_m), t_1, \dots, t_n) := f_\gamma(S_{\beta_1}(s_1, t_1, \dots, t_n), \dots, S_{\beta_m}(s_m, t_1, \dots, t_n))$.

Definition 1.3. Let $i \in I$ and $((f_\gamma)_k)_{k \in K_\gamma, \gamma \in \Sigma}$ be an indexed set of Σ -sorted operation symbols. Let $\Sigma_m(i) := \{\gamma \in \Sigma_m \mid \gamma(m+1) = i\}$, $m \in \mathbb{N}^+$ and let

$$\Sigma(i) := \bigcup_{m \geq 1} \Sigma_m(i).$$

Any mapping

$$\sigma_i : \{(f_\gamma)_k \mid k \in K_\gamma, \gamma \in \Sigma(i)\} \rightarrow W(i), i \in I,$$

which preserves arities, is said to be a Σ -hypersubstitution of sort i . Let $\Sigma(i)\text{-Hyp}$ be the set of all Σ -hypersubstitutions of sort i . The I -sorted mapping $\sigma := (\sigma_i)_{i \in I}$ is called an I -sorted Σ -hypersubstitution. Let $\Sigma\text{-Hyp}$ be the set of all I -sorted Σ -hypersubstitutions. Any I -sorted Σ -hypersubstitution σ can inductively be extended to an I -sorted mapping $\hat{\sigma} := (\hat{\sigma}_i)_{i \in I}$. The I -sorted mapping

$$\hat{\sigma} : W_\Sigma(X) \rightarrow W_\Sigma(X)$$

is defined by the following steps: For each $i \in I$ we define

- (i) $\hat{\sigma}_i[x_{ij}] := x_{ij}$ for any variable $x_{ij} \in X_i$.
- (ii) $\hat{\sigma}_i[f_\gamma(t_1, \dots, t_n)] := S_\gamma(\sigma_i(f_\gamma), \hat{\sigma}_{k_1}[t_1], \dots, \hat{\sigma}_{k_n}[t_n])$, where $\gamma = (k_1, \dots, k_n; i) \in \Sigma$ and $t_q \in W(k_q)$, $1 \leq q \leq n$, $n \in \mathbb{N}$, assumed that $\hat{\sigma}_{k_q}[t_q]$, are already defined.

Using the extension $\hat{\sigma}_i$, we define $(\sigma_1)_i \circ_i (\sigma_2)_i := (\hat{\sigma}_1)_i \circ (\hat{\sigma}_2)_i$. Then we have $((\sigma_1)_i \circ_i (\sigma_2)_i)^\wedge = (\hat{\sigma}_1)_i \circ (\hat{\sigma}_2)_i$. Together with the identity mapping $(\sigma_{id})_i$ the set $\Sigma(i)\text{-Hyp}$ forms a monoid (see [3]).

Now we want to describe the connection between heterogeneous algebras and Σ -terms.

Let A be an I -sorted set. Then \mathcal{A} is said to be a Σ -algebra if it has the form

$$\mathcal{A} = \left(A; (((f_\gamma)_k)^A)_{k \in K_\gamma, \gamma \in \Sigma} \right)$$

where $((f_\gamma)_k)^A : A_{k_1} \times \cdots \times A_{k_n} \rightarrow A_i$ if $\gamma = (k_1, \dots, k_n; i) \in \Sigma$. Let $\text{Alg}(\Sigma)$ be the collection of all Σ -algebras. To connect Σ -terms with Σ -algebras we need to consider operations on I -sorted sets. Let A be an I -sorted set, $n \in \mathbb{N}^+$, $(\omega; i) \in I^* \times I$. Then ω is called input sequence on A and i is called output sort.

Definition 1.4. Let A be an I -sorted set, let $\omega = (k_1, \dots, k_n) \in I^n$, $n \in \mathbb{N}^+$ be an input sequence on A . Then we define the q -th n -ary projection operation

$$e_q^{\omega, A} : A_{k_1} \times \cdots \times A_{k_n} \rightarrow A_{k_q}, 1 \leq q \leq n$$

of the input sequence ω on A by

$$e_q^{\omega, A}(a_1, \dots, a_n) := a_q.$$

We denote by

$$O^{(\omega, i)}(A) := \{f \mid f : A_{k_1} \times \cdots \times A_{k_n} \rightarrow A_i\}$$

the set of all n -ary operations on A with input sequence ω and output sort i .

In particular we denote by

$$O^\omega(A) := (O^{(\omega, i)}(A))_{i \in I}$$

the I -sorted set of all n -ary operations on A with the same input sequence ω .

Finally we introduce

$$O(A) := \bigcup_{\omega \in I^*} O^\omega(A)$$

as the I -sorted set of all finitary operations on the I -sorted set A .

Definition 1.5. Let A be an I -sorted set and let $\omega = (s_1, \dots, s_n), \omega' = (s'_1, \dots, s'_m)$ be input sequences on A . Then the superposition operation

$$S_{\omega'}^{\omega,i} : O^{(\omega,i)}(A) \times O^{(\omega',s_1)}(A) \times \cdots \times O^{(\omega',s_n)}(A) \rightarrow O^{(\omega',i)}(A)$$

is defined by

$$S_{\omega'}^{\omega,i}(f, g_1, \dots, g_n) := f[g_1, \dots, g_n], \text{ with}$$

$$f[g_1, \dots, g_n](a_1, \dots, a_m) := f(g_1(a_1, \dots, a_m), \dots, g_n(a_1, \dots, a_m))$$

for all $(a_1, \dots, a_m) \in A_{s'_1} \times \cdots \times A_{s'_m}$.

Using these composition operations we may consider a many-sorted algebra, which satisfies similar identities as clones in the one-sorted case.

Theorem 1.6. *Let A be an I -sorted set. Then the many-sorted algebra*

$$\left((O^\omega(A))_{\omega \in I^*}; \left(S_{\omega'}^{\omega,i} \right)_{(\omega,i),(\omega',i) \in I^* \times I}, \left(e_j^{\omega,A} \right)_{\omega \in I^*, 1 \leq j \leq |\omega|} \right)$$

(where $|\omega|$ is the length of the sequence ω) satisfies the following identities:

$$1) \quad S_{\omega''}^{\omega,i} \left(f, S_{\omega''}^{\omega',s_1}(g_1, h_1, \dots, h_m), \dots, S_{\omega''}^{\omega',s_n}(g_n, h_1, \dots, h_m) \right)$$

$$= S_{\omega''}^{\omega',i} \left(S_{\omega'}^{\omega,i}(f, g_1, \dots, g_n), h_1, \dots, h_m \right) \text{ where}$$

$$\omega = (s_1, \dots, s_n) \in I^n, \quad \omega' = (s'_1, \dots, s'_m) \in I^m, \quad \omega'' = (s''_1, \dots, s''_p) \in I^p,$$

and

$$f \in O^{(\omega, i)}(A), \quad g_j \in O^{(\omega', s_j)}(A), \quad h_k \in O^{(\omega'', s'_k)}(A) \quad \text{for } 1 \leq j \leq n,$$

$$1 \leq k \leq m, \quad m, n \in \mathbb{N}.$$

$$2) \quad S_{\omega'}^{\omega, s_j} \left(e_j^{\omega, A}, g_1, \dots, g_n \right) = g_j \quad \text{where } \omega = (s_1, \dots, s_n) \in I^n, \omega' \in I^m,$$

and

$$g_j \in O^{(\omega', s_j)}(A), \quad 1 \leq j \leq n, m, n \in \mathbb{N}^+.$$

$$3) \quad S_{\omega}^{\omega, i} \left(f, e_1^{\omega, A}, \dots, e_n^{\omega, A} \right) = f \quad \text{where } f \in O^{(\omega, i)}(A), \omega \in I^n, n \in \mathbb{N}^+.$$

The proofs are similar to the proofs of the corresponding propositions for Σ -terms (see [3]).

2. I -SORTED IDENTITIES AND MODEL CLASSES

Definition 2.1. Let $n \in \mathbb{N}^+$ and $X^{(n)}$ be an n -element I -sorted alphabet and let A be an I -sorted set. Let $\mathcal{A} \in \text{Alg}(\Sigma)$ be a Σ -algebra, and $t \in W_n(i), i \in I$. Let $f := (f_i)_{i \in I}$, where $f_i : X_i^{(n)} \rightarrow A_i$ is an I -sorted evaluation mapping of variables from $X^{(n)}$ by elements in A . Each mapping f_i can be extended in a canonical way to a mapping $\bar{f}_i : W_n(i) \rightarrow A_i$. Then $t^{\mathcal{A}} : A^{X^{(n)}} \rightarrow A_i$ is defined by

$$t^{\mathcal{A}}(f) := \bar{f}_i(t) \quad \text{for all } f \in A^{X^{(n)}},$$

where \bar{f}_i is the extension of the evaluation mapping $f_i : X_i^{(n)} \rightarrow A_i$. The operation $t^{\mathcal{A}}$ is called the n -ary Σ -term operation on \mathcal{A} induced by the n -ary Σ -term t of sort i . We have $x_{k_q q}^{\mathcal{A}} = e_q^{\omega, A}, 1 \leq q \leq n$, where $\omega = (k_1, \dots, k_n)$, since for $f \in A^{X^{(n)}}$ we have

$$\begin{aligned}
x_{k_q q}^A(f) &= \bar{f}_{k_q}(x_{k_q q}) \\
&= f_{k_q}(x_{k_q q}) \\
&= e_q^{\omega, A}(a_1, \dots, a_{q-1}, f_{k_q}(x_{k_q q}), a_{q+1}, \dots, a_n)
\end{aligned}$$

for all $a_j \in A_{k_j}$ such that $j \in \{1, \dots, q-1, q+1, \dots, n\}$.

Let $W^A(i)$ be the set of all Σ -term operations on \mathcal{A} induced by the Σ -terms of sort i . We set $W_\Sigma^A(X) := (W^A(i))_{i \in I}$ and call it the I -sorted set of Σ -term operations on \mathcal{A} induced by the Σ -terms.

Definition 2.2. Let $t \in W(i)$, $t_j \in W(k_j)$ where $1 \leq j \leq n, n \in \mathbb{N}$. Then the superposition operation

$$S_\alpha^A : W^A(i) \times W^A(k_1) \times \dots \times W^A(k_n) \rightarrow W^A(i)$$

where $\alpha = (k_1, \dots, k_n; i) \in \Lambda$, is inductively defined in the following way:

1) If $t = x_{ij} \in X_i$, then

$$1.1) \quad S_\alpha^A(x_{ij}^A, t_1^A, \dots, t_n^A) := x_{ij}^A \text{ for } i \neq k_j \text{ and}$$

$$1.2) \quad S_\alpha^A(x_{ij}^A, t_1^A, \dots, t_n^A) := t_j^A \text{ for } i = k_j.$$

2) If $t = f_\gamma(s_1, \dots, s_m) \in W(i)$ where $\gamma = (i_1, \dots, i_m; i) \in \Sigma$, $s_q \in W(i_q), 1 \leq q \leq m, m \in \mathbb{N}$ and assume that $S_{\alpha_q}^A(s_q^A, t_1^A, \dots, t_n^A)$, where $\alpha_q = (k_1, \dots, k_n; i_q) \in \Lambda$, are already defined, then

$$\begin{aligned}
& S_{\alpha}^A \left((f_{\gamma}(s_1, \dots, s_m))^A, t_1^A, \dots, t_n^A \right) \\
& := f_{\gamma}^A \left(S_{\alpha_1}^A (s_1^A, t_1^A, \dots, t_n^A), \dots, S_{\alpha_m}^A (s_m^A, t_1^A, \dots, t_n^A) \right).
\end{aligned}$$

Example 2.3. Let $I = \{1, 2\}$, $X^{(2)} = (X_i^{(2)})_{i \in I}$, $\Sigma = \{(1, 2; 1), (2, 1; 2)\}$. Let \mathcal{A} be a Σ -algebra and let $t = f_{(1,2;1)}(f_{(1,2;1)}(x_{11}, x_{21}), f_{(2,1;2)}(x_{22}, x_{11})) \in W(1)$, $t_1 \in W(2)$, and $t_2 \in W(1)$. Then

$$\begin{aligned}
S_{(2,1;1)}^A \left(t^A t_1^A t_2^A \right) &= S_{(2,1;1)}^A \left((f_{(1,2;1)}(f_{(1,2;1)}(x_{11}, x_{21}), f_{(2,1;2)}(x_{22}, x_{11})))^A t_1^A t_2^A \right) \\
&= f_{(1,2;1)}^A \left(S_{(2,1;1)}^A ((f_{(1,2;1)}(x_{11}, x_{21}))^A t_1^A t_2^A), \right. \\
&\quad \left. S_{(1,2;2)}^A ((f_{(2,1;2)}(x_{22}, x_{11}))^A, t_1^A, t_2^A) \right) \\
&= f_{(1,2;1)}^A \left(f_{(1,2;1)}^A \left(S_{(2,1;1)}^A (x_{11}^A, t_1^A, t_2^A), S_{(2,1;2)}^A (x_{21}^A, t_1^A, t_2^A) \right), \right. \\
&\quad \left. f_{(2,1;2)}^A \left(S_{(2,1;2)}^A (x_{22}^A, t_1^A, t_2^A), S_{(2,1;1)}^A (x_{11}^A, t_1^A, t_2^A) \right) \right) \\
&= f_{(1,2;1)}^A \left(f_{(1,2;1)}^A (x_{11}^A, t_1^A), f_{(2,1;2)}^A (x_{22}^A, x_{11}^A) \right).
\end{aligned}$$

Proposition 2.4. Let \mathcal{A} be a Σ -algebra and $f_{\gamma}(t_1, \dots, t_n) \in W_n(i)$ where $\gamma = (i_1, \dots, i_n, i) \in \Sigma$, $t_q \in W_n(i_q)$, $1 \leq q \leq n$, $n \in \mathbb{N}$. Then

$$\left(f_{\gamma}(t_1, \dots, t_n) \right)^A = f_{\gamma}^A(t_1^A, \dots, t_n^A).$$

Proof. Let $f \in A^{X^{(n)}}$, then

$$\begin{aligned}
\left(f_\gamma(t_1, \dots, t_n)\right)^{\mathcal{A}}(f) &= \bar{f}_i\left(f_\gamma(t_1, \dots, t_n)\right) \\
&= f_\gamma^{\mathcal{A}}\left(\bar{f}_{i_1}(t_1), \dots, \bar{f}_{i_n}(t_n)\right) \\
&= f_\gamma^{\mathcal{A}}\left(t_1^{\mathcal{A}}(f), \dots, t_n^{\mathcal{A}}(f)\right) \\
&= f_\gamma^{\mathcal{A}}\left(t_1^{\mathcal{A}}, \dots, t_n^{\mathcal{A}}\right)(f). \quad \blacksquare
\end{aligned}$$

Lemma 2.5. *Let \mathcal{A} be a Σ -algebra. For $t \in W(i)$, $t_j \in W(k_j)$, $1 \leq j \leq n$, $n \in \mathbb{N}$ we have:*

$$S_\alpha^{\mathcal{A}}\left(t^{\mathcal{A}}, t_1^{\mathcal{A}}, \dots, t_n^{\mathcal{A}}\right) = \left(S_\alpha(t, t_1, \dots, t_n)\right)^{\mathcal{A}}$$

where $\alpha = (k_1, \dots, k_n; i) \in \Lambda$.

Proof. We will give a proof by induction on the complexity of the Σ -term t .

1) If $t = x_{ij} \in X_i$, then

1.1) for $i \neq k_j$,

$$\begin{aligned}
S_\alpha^{\mathcal{A}}\left(t^{\mathcal{A}}, t_1^{\mathcal{A}}, \dots, t_n^{\mathcal{A}}\right) &= S_\alpha^{\mathcal{A}}\left(x_{ij}^{\mathcal{A}}, t_1^{\mathcal{A}}, \dots, t_n^{\mathcal{A}}\right) \\
&= x_{ij}^{\mathcal{A}} \\
&= \left(S_\alpha(x_{ij}, t_1, \dots, t_n)\right)^{\mathcal{A}} \\
&= \left(S_\alpha(t, t_1, \dots, t_n)\right)^{\mathcal{A}},
\end{aligned}$$

1.2) and for $i = k_j$,

$$\begin{aligned}
S_\alpha^A(t^A, t_1^A, \dots, t_n^A) &= S_\alpha^A(x_{ij}^A, t_1^A, \dots, t_n^A) \\
&= t_j^A \\
&= \left(S_\alpha(x_{ij}, t_1, \dots, t_n)\right)^A \\
&= \left(S_\alpha(t, t_1, \dots, t_n)\right)^A.
\end{aligned}$$

- 2) If $t = f_\gamma(s_1, \dots, s_m) \in W(i)$, where $\gamma = (i_1, \dots, i_m; i) \in \Sigma$ and $s_q \in W(i_q)$, $1 \leq q \leq m$, $m \in \mathbb{N}$, and if we assume that the equations

$$S_{\alpha_q}^A(s_q^A, t_1^A, \dots, t_n^A) = \left(S_{\alpha_q}(s_q, t_1, \dots, t_n)\right)^A,$$

where $\alpha_q = (k_1, \dots, k_n; i_q) \in \Lambda$, are satisfied, then for $f \in A^{X^{(n)}}$ we have

$$\begin{aligned}
&S_\alpha^A(t^A, t_1^A, \dots, t_n^A)(f) \\
&= S_\alpha^A\left(\left(f_\gamma(s_1, \dots, s_m)\right)^A, t_1^A, \dots, t_n^A\right)(f) \\
&= f_\gamma^A\left(S_{\alpha_1}^A\left((s_1^A, t_1^A, \dots, t_n^A)(f)\right), \dots, S_{\alpha_m}^A\left(s_m^A, t_1^A, \dots, t_n^A\right)(f)\right) \\
&= f_\gamma^A\left(\left(S_{\alpha_1}(s_1, t_1, \dots, t_n)\right)^A(f), \dots, \left(S_{\alpha_m}(s_m, t_1, \dots, t_n)\right)^A(f)\right) \\
&= f_\gamma^A\left(\bar{f}_{i_1}\left(S_{\alpha_1}(s_1, t_1, \dots, t_n)\right), \dots, \bar{f}_{i_m}\left(S_{\alpha_m}(s_m, t_1, \dots, t_n)\right)\right)
\end{aligned}$$

$$\begin{aligned}
&= \bar{f}_i \left(f_\gamma \left(S_{\alpha_1}(s_1, t_1, \dots, t_n), \dots, S_{\alpha_m}(s_m, t_1, \dots, t_n) \right) \right) \\
&= \left(f_\gamma \left(S_{\alpha_1}(s_1, t_1, \dots, t_n), \dots, S_{\alpha_m}(s_m, t_1, \dots, t_n) \right) \right)^{\mathcal{A}}(f) \\
&= \left(S_\alpha \left(f_\gamma(s_1, \dots, s_m), t_1, \dots, t_n \right) \right)^{\mathcal{A}}(f) \\
&= \left(S_\alpha(t, t_1, \dots, t_n) \right)^{\mathcal{A}}(f). \quad \blacksquare
\end{aligned}$$

Now we can define equations and identities.

Definition 2.6. A Σ -equation of sort i in X is a pair (s_i, t_i) of elements from $W(i)$, $i \in I$. Such pairs are more commonly written as $s_i \approx_i t_i$. The Σ -equation $s_i \approx_i t_i$ is said to be a Σ -identity of sort i in the Σ -algebra \mathcal{A} if $s_i^{\mathcal{A}} = t_i^{\mathcal{A}}$, that is, if the Σ -term operations induced by s_i and t_i , respectively, on the Σ -algebra \mathcal{A} are equal.

In this case we also say that the Σ -equation $s_i \approx_i t_i$ is satisfied or modelled by the Σ -algebra \mathcal{A} , and write $\mathcal{A} \models_i s_i \approx_i t_i$. If the Σ -equation $s_i \approx_i t_i$ is satisfied by every Σ -algebra \mathcal{A} of a class K_0 of Σ -algebras, we write $K_0 \models_i s_i \approx_i t_i$. For a set $F(i)$ of equations of sort i we write $\mathcal{A} \models_i F(i)$ if $\mathcal{A} \models_i s_i \approx_i t_i$ for all $(s_i, t_i) \in F(i)$.

Example 2.7. Let $I = \{1, 2\}$, $X^{(2)} := (X_i^{(2)})_{i \in I}$ be a 2-element I -sorted alphabet, and $\Sigma = \{(1, 1; 1), (2, 1; 1)\}$. Let $\mathcal{V} = (A; f_{(2,1;1)}^{\mathcal{V}}, f_{(1,1;1)}^{\mathcal{V}})$ where $f_{(2,1;1)}^{\mathcal{V}}, f_{(1,1;1)}^{\mathcal{V}}$ correspond to $\circ, +$, respectively, and $A := (V, \mathbb{R})$ is the universe of a real vector space. Then the Σ -equation

$$\begin{aligned}
&f_{(2,1;1)} \left(x_{21}, f_{(1,1;1)}(x_{11}, x_{12}) \right) \\
&\approx_1 f_{(1,1;1)} \left(f_{(2,1;1)}(x_{21}, x_{11}), f_{(2,1;1)}(x_{21}, x_{12}) \right) \in W(1)^2
\end{aligned}$$

is a Σ -identity of sort 1 in \mathcal{V} , that is,

$$\begin{aligned} \mathcal{V} \models_1 f_{(2,1;1)}(x_{21}, f_{(1,1;1)}(x_{11}, x_{12})) \\ \approx_1 f_{(1,1;1)}(f_{(2,1;1)}(x_{21}, x_{11}), f_{(2,1;1)}(x_{21}, x_{12})) \end{aligned}$$

since for $f \in A^{X^{(2)}}$ we have

$$\begin{aligned} f_{(2,1;1)}(x_{21}, f_{(1,1;1)}(x_{11}, x_{12}))^{\mathcal{V}}(f) &= \bar{f}_1(f_{(2,1;1)}(x_{21}, f_{(1,1;1)}(x_{11}, x_{12}))) \\ &= f_{(2,1;1)}^{\mathcal{V}}(\bar{f}_2(x_{21}), \bar{f}_1(f_{(1,1;1)}(x_{11}, x_{12}))) \\ &= f_{(2,1;1)}^{\mathcal{V}}(\bar{f}_2(x_{21}), f_{(1,1;1)}^{\mathcal{V}}(\bar{f}_1(x_{11}), \bar{f}_1(x_{12}))) \\ &= f_{(2,1;1)}^{\mathcal{V}}(f_2(x_{21}), f_{(1,1;1)}^{\mathcal{V}}(f_1(x_{11}), f_1(x_{12}))) \end{aligned}$$

and

$$\begin{aligned} f_{(1,1;1)}(f_{2,1;1}(x_{21}, x_{11}), f_{(2,1;1)}(x_{21}, x_{12}))^{\mathcal{V}}(f) \\ &= \bar{f}_1(f_{(1,1;1)}(f_{(2,1;1)}(x_{21}, x_{11}), f_{(2,1;1)}(x_{21}, x_{12}))) \\ &= f_{(1,1;1)}^{\mathcal{V}}(\bar{f}_1(f_{(2,1;1)}(x_{21}, x_{11})), \bar{f}_1(f_{(2,1;1)}(x_{21}, x_{12}))) \\ &= f_{(1,1;1)}^{\mathcal{V}}(f_{(2,1;1)}^{\mathcal{V}}(\bar{f}_2(x_{21}), \bar{f}_1(x_{11})), f_{(2,1;1)}^{\mathcal{V}}(\bar{f}_2(x_{21}), \bar{f}_1(x_{12}))) \\ &= f_{(1,1;1)}^{\mathcal{V}}(f_{(2,1;1)}^{\mathcal{V}}(f_2(x_{21}), f_1(x_{11})), f_{(2,1;1)}^{\mathcal{V}}(f_2(x_{21}), f_1(x_{12}))). \end{aligned}$$

Therefore,

$$\begin{aligned} & \left(f_{(2,1;1)} \left(x_{21}, f_{(1,1;1)}(x_{11}, x_{12}) \right) \right)^\nu \\ &= \left(f_{(1,1;1)} \left(f_{2,1;1}(x_{21}, x_{11}), f_{(2,1;1)}(x_{21}, x_{12}) \right) \right)^\nu. \end{aligned}$$

Now we extend the usual Galois-connection between identities and algebras to the many-sorted case.

Let $K_0 \subseteq \text{Alg}(\Sigma)$ and $L(i) \subseteq W(i)^2$. Then a mapping

$$\Sigma(i)\text{-Id} : P(\text{Alg}(\Sigma)) \rightarrow P(W(i)^2)$$

is defined by

$$\Sigma(i)\text{-Id}K_0 := \left\{ (s_i, t_i) \in W(i)^2 \mid (\forall \mathcal{A} \in K_0)(\mathcal{A} \models_i s_i \approx_i t_i) \right\}$$

and a mapping $\Sigma(i)\text{-Mod} : P(W(i)^2) \rightarrow P(\text{Alg}(\Sigma))$ is defined by

$$\Sigma(i)\text{-Mod}L(i) := \{ \mathcal{A} \in \text{Alg}(\Sigma) \mid (\forall (s_i, t_i) \in L(i))(\mathcal{A} \models_i s_i \approx_i t_i) \}.$$

In the next propositions, we will show that these two mappings satisfy the Galois-connection properties.

Proposition 2.8. *Let $i \in I$ and let $K_0, K_1, K_2 \subseteq \text{Alg}(\Sigma)$. Then*

- (1) $K_1 \subseteq K_2 \Rightarrow \Sigma(i)\text{-Id}K_2 \subseteq \Sigma(i)\text{-Id}K_1,$
- (2) $K_0 \subseteq \Sigma(i)\text{-Mod}\Sigma(i)\text{-Id}K_0.$

Proof.

- (1) Assume that $K_1 \subseteq K_2$ and let $s_i \approx_i t_i \in \Sigma(i)\text{-Id}K_2$. Then for all $\mathcal{A} \in K_2$, we have $\mathcal{A} \models_i s_i \approx_i t_i$. Because of $K_1 \subseteq K_2$, we obtain $\mathcal{A} \models_i s_i \approx_i t_i$, for all $\mathcal{A} \in K_1$. This means that $s_i \approx_i t_i \in \Sigma(i)\text{-Id}K_1$, and then $\Sigma(i)\text{-Id}K_2 \subseteq \Sigma(i)\text{-Id}K_1$.
- (2) Let $\mathcal{A} \in K_0$. Then $\mathcal{A} \models_i \Sigma(i)\text{-Id}K_0$, means that $\mathcal{A} \in \Sigma(i)\text{-Mod}\Sigma(i)\text{-Id}K_0$ and then $K_0 \subseteq \Sigma(i)\text{-Mod}\Sigma(i)\text{-Id}K_0$.

■

Proposition 2.9. *Let $L(i), L_1(i), L_2(i) \subseteq W(i)^2$ be subsets of the set of all Σ -equations of sort $i \in I$. Then*

- (1) $L_1(i) \subseteq L_2(i) \Rightarrow \Sigma(i)\text{-Mod}L_2(i) \subseteq \Sigma(i)\text{-Mod}L_1(i)$,
- (2) $L(i) \subseteq \Sigma(i)\text{-Id}\Sigma(i)\text{-Mod}L(i)$.

Proof.

- (1) Assume that $L_1(i) \subseteq L_2(i)$ and let $\mathcal{A} \in \Sigma(i)\text{-Mod}L_2(i)$. Then $\mathcal{A} \models_i s_i \approx_i t_i$ for all $s_i \approx_i t_i \in L_2(i)$, but we have $L_1(i) \subseteq L_2(i)$, so that $\mathcal{A} \models_i s_i \approx_i t_i$ for all $s_i \approx_i t_i \in L_1(i)$. It follows that $\mathcal{A} \in \Sigma(i)\text{-Mod}L_1(i)$ and then $\Sigma(i)\text{-Mod}L_2(i) \subseteq \Sigma(i)\text{-Mod}L_1(i)$.
- (2) Let $s_i \approx_i t_i \in L(i)$. Then we have $\Sigma(i)\text{-Mod}L(i) \models_i s_i \approx_i t_i$, that is $s_i \approx_i t_i \in \Sigma(i)\text{-Id}\Sigma(i)\text{-Mod}L(i)$ and then $L(i) \subseteq \Sigma(i)\text{-Id}\Sigma(i)\text{-Mod}L(i)$.

■

From both propositions, we have that $(\Sigma(i)\text{-Mod}, \Sigma(i)\text{-Id})$ is a Galois connection between $\text{Alg}(\Sigma)$ and $W(i)^2$ with respect to the relation

$$\models_i := \left\{ (\mathcal{A}, (s_i, t_i)) \in \text{Alg}(\Sigma) \times W(i)^2 \mid \mathcal{A} \models_i s_i \approx_i t_i \right\}.$$

The fixed points with respect to the closure operator $\Sigma(i)\text{-Mod}\Sigma(i)\text{-Id}$ are called Σ -varieties of sort i and the fixed points with respect to the closure operator $\Sigma(i)\text{-Id}\Sigma(i)\text{-Mod}$ are called Σ -equational theories of sort i .

3. APPLICATION OF Σ -HYPERSUBSTITUTIONS

Now we apply Σ -hypersubstitutions to many-sorted algebras and to many-sorted equations.

Definition 3.1. Let A be an I -sorted set, let $\mathcal{A} := (A; (((f_\gamma)_k)^{\mathcal{A}})_{k \in K_\gamma, \gamma \in \Sigma})$ be a Σ -algebra and let $\sigma \in \Sigma\text{-Hyp}$. Then we define the Σ -algebra

$$\sigma(\mathcal{A}) := \left(A; \left((\sigma_i((f_\gamma)_k))^{\mathcal{A}} \right)_{k \in K_\gamma, \gamma \in \Sigma(i), i \in I} \right).$$

This Σ -algebra is called the Σ -algebra derived from \mathcal{A} and σ , for short derived Σ -algebra.

For illustration we consider the following example.

Example 3.2. Let $I = \{1, 2\}$, $\Sigma = \{(1, 2, 1), (2, 1, 2)\}$, $K_{(1,2,1)} = \{1, 2\}$, $A = (A_1, A_2)$, $\mathcal{A} = ((A_1, A_2); ((f_{(1,2,1)})_1)^{\mathcal{A}}, ((f_{(1,2,1)})_2)^{\mathcal{A}}, f_{(2,1,2)}^{\mathcal{A}})$. Let $\sigma = (\sigma_1, \sigma_2) \in \Sigma\text{-Hyp}$. Then we have

$$\begin{aligned} & \sigma(\mathcal{A}) \\ &= \left((A_1, A_2); \left(\sigma_1((f_{(1,2,1)})_1) \right)^{\mathcal{A}}, \left(\sigma_1((f_{(1,2,1)})_2) \right)^{\mathcal{A}}, \left(\sigma_2(f_{(2,1,2)}) \right)^{\mathcal{A}} \right). \end{aligned}$$

Theorem 3.3. Let A be an I -sorted set and $\mathcal{A} := (A; (((f_\gamma)_k)^{\mathcal{A}})_{k \in K_\gamma, \gamma \in \Sigma})$ be a Σ -algebra. Let $\sigma \in \Sigma\text{-Hyp}$ and $t \in W(i), i \in I$. Then $t^{\sigma(\mathcal{A})} = (\hat{\sigma}_i[t])^{\mathcal{A}}$.

Proof. We will give a proof by induction on the complexity of the Σ -term t .

- 1) If $t = x_{ij} \in X_i$ where $1 \leq j \leq n, n \in \mathbb{N}$, then for $f \in A^{X^{(n)}}$ we have

$$\begin{aligned}
t^{\sigma(\mathcal{A})}(f) &= x_{ij}^{\sigma(\mathcal{A})}(f) \\
&= \bar{f}_i(x_{ij}) \\
&= x_{ij}^{\mathcal{A}}(f) \\
&= (\hat{\sigma}_i[x_{ij}])^{\mathcal{A}}(f) \\
&= (\hat{\sigma}_i[t])^{\mathcal{A}}(f).
\end{aligned}$$

2) If $t = f_\gamma(s_1, \dots, s_m) \in W(i)$ where $\gamma = (i_1, \dots, i_m; i) \in \Sigma$, $s_q \in W(i_q)$, $1 \leq q \leq m$, $m \in \mathbb{N}$ and assume that $s_q^{\sigma(\mathcal{A})} = \hat{\sigma}_{i_q}[s_q]^{\mathcal{A}}$ are satisfied, then for $f \in A^{X^{(n)}}$ we have

$$\begin{aligned}
t^{\sigma(\mathcal{A})}(f) &= (f_\gamma(s_1, \dots, s_m))^{\sigma(\mathcal{A})}(f) \\
&= \bar{f}_i(f_\gamma(s_1, \dots, s_m)) \\
&= f_\gamma^{\sigma(\mathcal{A})}(\bar{f}_{i_1}(s_1), \dots, \bar{f}_{i_m}(s_m)) \\
&= f_\gamma^{\sigma(\mathcal{A})}(s_1^{\sigma(\mathcal{A})}(f), \dots, s_m^{\sigma(\mathcal{A})}(f)) \\
&= \sigma_i(f_\gamma)^{\mathcal{A}}(\hat{\sigma}_{i_1}[s_1]^{\mathcal{A}}(f), \dots, \hat{\sigma}_{i_m}[s_m]^{\mathcal{A}}(f)) \\
&= \sigma_i(f_\gamma)^{\mathcal{A}}(\hat{\sigma}_{i_1}[s_1]^{\mathcal{A}}, \dots, \hat{\sigma}_{i_m}[s_m]^{\mathcal{A}})(f) \\
&= S_\gamma^{\mathcal{A}}(\sigma_i(f_\gamma)^{\mathcal{A}}, \hat{\sigma}_{i_1}[s_1]^{\mathcal{A}}, \dots, \hat{\sigma}_{i_m}[s_m]^{\mathcal{A}})(f) \\
&= (S_\gamma(\sigma_i(f_\gamma), \hat{\sigma}_{i_1}[s_1], \dots, \hat{\sigma}_{i_m}[s_m]))^{\mathcal{A}}(f) \text{ by Lemma 2.5} \\
&= (\hat{\sigma}_i[f_\gamma(s_1, \dots, s_m)])^{\mathcal{A}}(f) \\
&= (\hat{\sigma}_i[t])^{\mathcal{A}}(f). \quad \blacksquare
\end{aligned}$$

Lemma 3.4. *Let $\mathcal{A} \in \text{Alg}(\Sigma)$, $\sigma_1, \sigma_2 \in \Sigma\text{-Hyp}$. Then we have*

$$\left((\sigma_1)_i(f_\gamma)\right)^{\sigma_2(\mathcal{A})} = \left((\sigma_2)_i \circ_i (\sigma_1)_i(f_\gamma)\right)^{\mathcal{A}},$$

for $\gamma \in \Sigma(i), i \in I$.

Proof. By Theorem 3.3, we have

$$\begin{aligned} \left((\sigma_1)_i(f_\gamma)\right)^{\sigma_2(\mathcal{A})} &= \left((\hat{\sigma}_2)_i[(\sigma_1)_i(f_\gamma)]\right)^{\mathcal{A}} \\ &= \left((\hat{\sigma}_2)_i \circ (\sigma_1)_i(f_\gamma)\right)^{\mathcal{A}} \\ &= \left((\sigma_2)_i \circ_i (\sigma_1)_i(f_\gamma)\right)^{\mathcal{A}}. \end{aligned}$$

■

Let σ_1, σ_2 be elements in $\Sigma\text{-Hyp}$. Then we set $\sigma_1 \diamond \sigma_2 := ((\sigma_1)_i \circ_i (\sigma_2)_i)_{i \in I}$.

Lemma 3.5. *Let A be an I -sorted set, let $\mathcal{A} = (A; ((f_\gamma)_k)^{\mathcal{A}})_{k \in K_\gamma, \gamma \in \Sigma}$ be a Σ -algebra, and $\sigma_1, \sigma_2 \in \Sigma\text{-Hyp}$. Then we have*

$$\sigma_1(\sigma_2(\mathcal{A})) = (\sigma_2 \diamond \sigma_1)(\mathcal{A}).$$

Proof. By Lemma 3.4, we have

$$\begin{aligned} \sigma_1(\sigma_2(\mathcal{A})) &= \left(A; \left((\sigma_1)_i((f_\gamma)_k)^{\sigma_2(\mathcal{A})}\right)_{k \in K_\gamma, \gamma \in \Sigma(i), i \in I}\right) \\ &= \left(A; \left((\sigma_2)_i \circ_i (\sigma_1)_i((f_\gamma)_k)^{\mathcal{A}}\right)_{k \in K_\gamma, \gamma \in \Sigma(i), i \in I}\right) \\ &= (\sigma_2 \diamond \sigma_1)(\mathcal{A}). \end{aligned}$$

■

Theorem 3.6. *Let A be an I -sorted set, $\mathcal{A} := (A; (((f_\alpha)_k)^A)_{k \in K_\alpha, \alpha \in \Sigma})$, and $\sigma_{id} \in \Sigma\text{-Hyp}$. Then we have*

$$\sigma_{id}(\mathcal{A}) = \mathcal{A}.$$

Proof. We will show that $((\sigma_{id})_i(f_\alpha)_k)^A = f_\alpha^A$ for all $k \in K_\alpha, \alpha \in \Sigma$. Assume that $\alpha = (k_1, \dots, k_n; i) \in \Sigma$ and $\omega = (k_1, \dots, k_n) \in I^n$. Then

$$\begin{aligned} \left((\sigma_{id})_i(f_\alpha) \right)^A &= \left(f_\alpha(x_{k_1 1}, \dots, x_{k_n n}) \right)^A \\ &= f_\alpha^A(x_{k_1 1}^A, \dots, x_{k_n n}^A) \\ &= f_\alpha^A(e_1^{\omega, A}, \dots, e_n^{\omega, A}) \\ &= f_\alpha^A. \end{aligned}$$

■

Definition 3.7. A Σ -algebra \mathcal{A} is said to hypersatisfy the Σ -identity $s_i \approx_i t_i$ of sort $i \in I$, if for every Σ -hypersubstitution of sort i , i.e., $\sigma_i \in \Sigma(i)\text{-Hyp}$, the Σ -identity $\hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i]$ holds in \mathcal{A} .

In this case we say that the Σ -identity $s_i \approx_i t_i$ of sort i is satisfied as a Σ -hyperidentity of sort i in \mathcal{A} and write $\mathcal{A} \models_{\Sigma\text{-hyp}} s_i \approx_i t_i$, that is

$$\mathcal{A} \models_{\Sigma\text{-hyp}} s_i \approx_i t_i \Leftrightarrow \forall \sigma_i \in \Sigma(i)\text{-Hyp} \quad (\mathcal{A} \models_i \hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i]).$$

Let us consider the following example.

Example 3.8. Let $I = \{1, 2\}$, $X^{(2)} := (X_i^{(2)})_{i \in I}$ and let $\Sigma = \{(1, 1; 1), (2, 2; 2)\}$. Let $\mathcal{B}_i := (B_i; \circ_i)$ be bands. Then $f_{(i, i, i)}(x_{ij}, x_{ij}) \approx_i x_{ij}$ are hyperidentities in $\mathcal{B}_i, i \in I$. Let $\mathcal{B} := (B; \circ)$ be a double band, where $B := (B_i)_{i \in I}$, $\circ := (\circ_i)_{i \in I}$. Then $f_{(i, i, i)}(x_{ij}, x_{ij}) \approx_i x_{ij}$ are Σ -hyperidentities of sort i in \mathcal{B} .

Let $K_0 \subseteq Alg(\Sigma)$ be a set of Σ -algebras, and let $L(i) \subseteq W(i)^2$ be a set of Σ -equations of sort i . Then we define a mapping

$$H\Sigma(i)-Id : P(Alg(\Sigma)) \rightarrow P(W(i)^2)$$

by

$$H\Sigma(i)-IdK_0 := \left\{ (s_i, t_i) \in W(i)^2 \mid (\forall \mathcal{A} \in K_0) \left(\mathcal{A} \models_{\Sigma-hyp} s_i \approx_i t_i \right) \right\}$$

and a mapping $H\Sigma(i)-Mod : P(W(i)^2) \rightarrow P(Alg(\Sigma))$ by

$$H\Sigma(i)-ModL(i) := \left\{ \mathcal{A} \in Alg(\Sigma) \mid (\forall (s_i, t_i) \in L(i)) \left(\mathcal{A} \models_{\Sigma-hyp} s_i \approx_i t_i \right) \right\}.$$

We get that $(H\Sigma(i)-Mod, H\Sigma(i)-Id)$ is also a Galois connection between $Alg(\Sigma)$ and $W(i)^2$ with respect to the relation

$$\models_{\Sigma-hyp} := \left\{ (\mathcal{A}, (s_i, t_i)) \in Alg(\Sigma) \times W(i)^2 \mid \mathcal{A} \models_{\Sigma-hyp} s_i \approx_i t_i \right\}.$$

Definition 3.9. Let $K_0 \subseteq Alg(\Sigma)$ be a subclass of Σ -algebras and let $L(i) \subseteq W(i)^2$ be a set of Σ -equations of sort i . Then we set

$$\chi^{\Sigma-E(i)}[s_i \approx_i t_i] := \{ \hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i] \mid \sigma_i \in \Sigma(i)-Hyp \}$$

and

$$\chi^{\Sigma-A}[\mathcal{A}] := \{ \sigma(\mathcal{A}) \mid \sigma \in \Sigma-Hyp \}.$$

We define two operators

$$\chi^{\Sigma-E(i)} : P(W(i)^2) \rightarrow P(W(i)^2)$$

by

$$\chi^{\Sigma-E(i)}[L(i)] := \bigcup_{s_i \approx_i t_i \in L(i)} \chi^{\Sigma-E(i)}[s_i \approx_i t_i]$$

and

$$\chi^{\Sigma-A} : P(\text{Alg}(\Sigma)) \rightarrow P(\text{Alg}(\Sigma))$$

by

$$\chi^{\Sigma-A}[K_0] := \bigcup_{\mathcal{A} \in K_0} \chi^{\Sigma-A}[\mathcal{A}].$$

Proposition 3.10. *Let $L(i), L_k(i) \subseteq W(i)^2$ be sets of Σ -equations of sort $i \in I$ with $k = 1, 2$. Then*

- (i) $L(i) \subseteq \chi^{\Sigma-E(i)}[L(i)]$,
- (ii) $L_1(i) \subseteq L_2(i) \Rightarrow \chi^{\Sigma-E(i)}[L_1(i)] \subseteq \chi^{\Sigma-E(i)}[L_2(i)]$,
- (iii) $\chi^{\Sigma-E(i)}[L(i)] = \chi^{\Sigma-E(i)}[\chi^{\Sigma-E(i)}[L(i)]]$.

Proof.

- (i) Let $s_i \approx_i t_i \in L(i)$. Then since $s_i = (\hat{\sigma}_{id})_i[s_i]$ and $t_i = (\hat{\sigma}_{id})_i[t_i]$, we have $(\hat{\sigma}_{id})_i[s_i] = s_i \approx_i t_i = (\hat{\sigma}_{id})_i[t_i] \in \chi^{\Sigma-E(i)}[L(i)]$ and then $L(i) \subseteq \chi^{\Sigma-E(i)}[L(i)]$.
- (ii) Assume that $L_1(i) \subseteq L_2(i)$ and let $\hat{\sigma}[s_i] \approx_i \hat{\sigma}[t_i] \in \chi^{\Sigma-E(i)}[L_1(i)]$. Then $s_i \approx_i t_i \in L_1(i)$ but $L_1(i) \subseteq L_2(i)$, so that $s_i \approx_i t_i \in L_2(i)$ and $\hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i] \in \chi^{\Sigma-E(i)}[L_2(i)]$. We have $\chi^{\Sigma-E(i)}[L_1(i)] \subseteq \chi^{\Sigma-E(i)}[L_2(i)]$.

- (iii) By (i) we have $\chi^{\Sigma-E(i)}[L(i)] \subseteq \chi^{\Sigma-E(i)}[\chi^{\Sigma-E(i)}[L(i)]]$. Let $\hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i] \in \chi^{\Sigma-E(i)}[\chi^{\Sigma-E(i)}[L(i)]]$. Then $s_i \approx_i t_i \in \chi^{\Sigma-E(i)}[L(i)]$, and there exists $\rho_i \in \Sigma(i)\text{-Hyp}$ and $u_i \approx_i v_i \in L(i)$ such that $s_i = \hat{\rho}_i[u_i]$ and $t_i = \hat{\rho}_i[v_i]$, and we have

$$\begin{aligned} \hat{\sigma}_i[s_i] &= \hat{\sigma}_i[\hat{\rho}_i[u_i]] \\ &= (\hat{\sigma}_i \circ \hat{\rho}_i) [u_i] \\ &= (\sigma_i \circ_i \rho_i)^\wedge [u_i] \\ &= \hat{\lambda}_i[u_i], \text{ where } \lambda_i = \sigma_i \circ_i \rho_i \in \Sigma(i)\text{-Hyp}, \end{aligned}$$

and

$$\begin{aligned} \hat{\sigma}_i[t_i] &= \hat{\sigma}_i[\hat{\rho}_i[v_i]] \\ &= (\hat{\sigma}_i \circ \hat{\rho}_i) [v_i] \\ &= (\sigma_i \circ_i \rho_i)^\wedge [v_i] \\ &= \hat{\lambda}_i[v_i]. \end{aligned}$$

Then we set

$$\hat{\lambda}_i[u_i] = \hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i] = \hat{\lambda}_i[v_i] \in \chi^{\Sigma-E(i)}[L(i)],$$

and then

$$\chi^{\Sigma-E(i)}[\chi^{\Sigma-E(i)}[L(i)]] \subseteq \chi^{\Sigma-E(i)}[L(i)].$$

■

Proposition 3.11. *Let $K_0, K_1, K_2 \subseteq \text{Alg}(\Sigma)$ be classes of Σ -algebras. Then*

- (i) $K_0 \subseteq \chi^{\Sigma-A}[K_0]$,
- (ii) $K_1 \subseteq K_2 \Rightarrow \chi^{\Sigma-A}[K_1] \subseteq \chi^{\Sigma-A}[K_2]$,
- (iii) $\chi^{\Sigma-A}[K_0] = \chi^{\Sigma-A}[\chi^{\Sigma-A}[K_0]]$.

Proof.

- (i) Let $\mathcal{A} \in K_0$. Then since $\mathcal{A} = \sigma_{id}(\mathcal{A}) \in \chi^{\Sigma-A}[K_0]$, we have $K_0 \subseteq \chi^{\Sigma-A}[K_0]$.
- (ii) Assume that $K_1 \subseteq K_2$ and let $\sigma(\mathcal{A}) \in \chi^{\Sigma-A}[K_1]$. Then $\mathcal{A} \in K_1$ by our assumption that $\mathcal{A} \in K_2$, with $\sigma(\mathcal{A}) \in \chi^{\Sigma-A}[K_2]$, and then $\chi^{\Sigma-A}[K_1] \subseteq \chi^{\Sigma-A}[K_2]$.
- (iii) By (i), we have $\chi^{\Sigma-A}[K_0] \subseteq \chi^{\Sigma-A}[\chi^{\Sigma-A}[K_0]]$. We will show that $\chi^{\Sigma-A}[\chi^{\Sigma-A}[K_0]] \subseteq \chi^{\Sigma-A}[K_0]$. Let $\sigma(\mathcal{A}) \in \chi^{\Sigma-A}[\chi^{\Sigma-A}[K_0]]$. Then $\mathcal{A} \in \chi^{\Sigma-A}[K_0]$, and there exists $\rho \in \Sigma\text{-Hyp}$ and $\mathcal{B} \in K_0$ such that $\mathcal{A} = \rho(\mathcal{B})$. We have

$$\begin{aligned} \sigma(\mathcal{A}) &= \sigma(\rho(\mathcal{B})) \\ &= (\rho \diamond \sigma)(\mathcal{B}) \\ &= \lambda(\mathcal{B}), \text{ where } \lambda = \rho \diamond \sigma \in \Sigma\text{-Hyp}. \end{aligned}$$

Thus we have $\sigma(\mathcal{A}) = \lambda(\mathcal{B}) \in \chi^{\Sigma-A}[K_0]$ and then $\chi^{\Sigma-A}[\chi^{\Sigma-A}[K_0]] \subseteq \chi^{\Sigma-A}[K_0]$. ■

Lemma 3.12. *Let $\mathcal{A} \in \text{Alg}(\Sigma)$ be a Σ -algebra, let $s_i \approx_i t_i \in W(i)^2$ be a Σ -equation of sort $i \in I$, and $\sigma \in \Sigma\text{-Hyp}$. Then*

$$\sigma(\mathcal{A}) \models_i s_i \approx_i t_i \iff \mathcal{A} \models_i \hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i].$$

Proof. We obtain

$$\begin{aligned} \sigma(\mathcal{A}) \models_i s_i \approx_i t_i &\iff s_i^{\sigma(\mathcal{A})} = t_i^{\sigma(\mathcal{A})} \\ &\iff \hat{\sigma}_i[s_i]^{\mathcal{A}} = \hat{\sigma}_i[t_i]^{\mathcal{A}} \\ &\iff \mathcal{A} \models_i \hat{\sigma}_i[s_i] \approx_i \hat{\sigma}_i[t_i]. \end{aligned}$$

■

The next theorem needs the concept of a conjugate pair of additive closure operators (see [4]).

Theorem 3.13. *The pair $(\chi^{\Sigma-A}, \chi^{\Sigma-E(i)})$ is a conjugate pair of completely additive closure operators of sort i with respect to the relation \models_i .*

Proof. By Definition 3.9, Propositions 3.10–3.11, and Lemma 3.12. ■

Now we may apply the theory of conjugate pairs of additive closure operators (see e.g., [4]) and obtain the following propositions:

Lemma 3.14 ([4]). *For all $K_0 \subseteq \text{Alg}(\Sigma)$ and for all $L(i) \subseteq W(i)^2$ the following properties hold:*

- (i) $H\Sigma(i)\text{-Mod}L(i) = \Sigma(i)\text{-Mod}\chi^{\Sigma-E(i)}[L(i)],$
- (ii) $H\Sigma(i)\text{-Mod}L(i) \subseteq \Sigma(i)\text{-Mod}L(i),$
- (iii) $\chi^{\Sigma-A}[H\Sigma(i)\text{-Mod}L(i)] = H\Sigma(i)\text{-Mod}L(i),$

- (iv) $\chi^{\Sigma-E(i)}[\Sigma(i)-IdH\Sigma(i)-ModL(i)] = \Sigma(i)-IdH\Sigma(i)-ModL(i),$
- (v) $H\Sigma(i)-ModH\Sigma(i)-IdK_0 = \Sigma(i)-Mod\Sigma(i)-Id\chi^{\Sigma-A}[K_0],$ and
- (i)' $H\Sigma(i)-IdK_0 = \Sigma(i)-Id\chi^{\Sigma-A}[K_0],$
- (ii)' $H\Sigma(i)-IdK_0 \subseteq \Sigma(i)-IdK_0,$
- (iii)' $\chi^{\Sigma-E(i)}[H\Sigma(i)-IdK_0] = H\Sigma(i)-IdK_0,$
- (iv)' $\chi^{\Sigma-A}[\Sigma(i)-ModH\Sigma(i)-IdK_0] = \Sigma(i)-ModH\Sigma(i)-IdK_0,$
- (v)' $H\Sigma(i)-IdH\Sigma(i)-ModL(i) = \Sigma(i)-Id\Sigma(i)-Mod\chi^{\Sigma-E(i)}[L(i)].$

4. I -SORTED SOLID Σ -VARIETIES

Definition 4.1. Let $K_0 \subseteq Alg(\Sigma)$ be a subclass of Σ -algebras. Then K_0 is called a solid model class of sort i or a solid Σ -variety of sort i if every Σ -identity of sort i is satisfied as a Σ -hyperidentity of sort i :

$$K_0 \models_{\Sigma-hyp}^i \Sigma(i)-IdK_0.$$

K_0 is called an I -sorted solid model class if every Σ -identity of sort i is satisfied as a Σ -hyperidentity of sort i for all $i \in I$, that is,

$$K_0 \models_{\Sigma-hyp}^i \Sigma(i)-IdK_0 \text{ for all } i \in I.$$

$L(i)$ is said to be a Σ -equational theory of sort i if there exists a class of Σ -algebras K_0 such that $L(i) = \Sigma(i)-IdK_0$. Then we set $L := (L(i))_{i \in I}$. This I -sorted set is called I -sorted Σ -equational theory.

Using the propositions of Lemma 3.14 one obtains the following characterization of solid Σ -varieties of sort i and solid Σ -equational theories of sort i (see e.g., [4]).

Theorem 4.2 ([4]). *Let K_0 be a Σ -variety of sort i . Then the following properties are equivalent:*

- (i) $K_0 = H\Sigma(i)\text{-}ModH\Sigma(i)\text{-}IdK_0$,
- (ii) $\chi^{\Sigma-A}[K_0] = K_0$,
- (iii) $\Sigma(i)\text{-}IdK_0 = H\Sigma(i)\text{-}IdK_0$,
- (iv) $\chi^{\Sigma-E(i)}[\Sigma(i)\text{-}IdK_0] = \Sigma(i)\text{-}IdK_0$.

Theorem 4.3 ([4]). *Let $L(i)$ be a Σ -equational theory of sort i . Then the following properties are equivalent:*

- (i) $L(i) = H\Sigma(i)\text{-}IdH\Sigma(i)\text{-}ModL(i)$,
- (ii) $\chi^{\Sigma-E(i)}[L(i)] = L(i)$,
- (iii) $\Sigma(i)\text{-}ModL(i) = H\Sigma(i)\text{-}ModL(i)$,
- (iv) $\chi^{\Sigma-A}[\Sigma(i)\text{-}ModL(i)] = \Sigma(i)\text{-}ModL(i)$.

5. I -SORTED COMPLETE LATTICES

Let $\mathcal{H}(i)$ be the class of all fixed points with respect to the closure operator $\Sigma(i)\text{-}Mod\Sigma(i)\text{-}Id$:

$$\mathcal{H}(i) := \{K_0 \subseteq Alg(\Sigma) \mid K_0 = \Sigma(i)\text{-}Mod\Sigma(i)\text{-}IdK_0\},$$

that is, $\mathcal{H}(i)$ is the class of all Σ -varieties of sort i . Then $\mathcal{H}(i)$ forms a complete lattice of Σ -varieties of sort i . Let $\mathcal{Hy}(i)$ be the class of all fixed points with respect to the closure operator $H\Sigma(i)\text{-}ModH\Sigma(i)\text{-}Id$:

$$\mathcal{Hy}(i) := \{K_0 \subseteq Alg(\Sigma) \mid K_0 = H\Sigma(i)\text{-}ModH\Sigma(i)\text{-}IdK_0\},$$

that is, $\mathcal{Hy}(i)$ is the class of all solid Σ -varieties of sort i . Then $\mathcal{Hy}(i)$ forms a complete lattice of solid Σ -varieties of sort i and $\mathcal{Hy}(i)$ is a complete sublattice of $\mathcal{H}(i)$. We set $\mathcal{H} := (\mathcal{H}(i))_{i \in I}$ and $\mathcal{Hy} := (\mathcal{Hy}(i))_{i \in I}$. \mathcal{H} is called an I -sorted complete lattice. \mathcal{Hy} is called an I -sorted complete sublattice of \mathcal{H} , since for every $i \in I$, $\mathcal{Hy}(i)$ is a complete sublattice of $\mathcal{H}(i)$. Dually

Let $\mathcal{L}(i)$ be the class of all fixed points with respect to the closure operator $\Sigma(i)\text{-}Id\Sigma(i)\text{-}Mod$:

$$\mathcal{L}(i) := \{L(i) \subseteq W(i)^2 \mid L(i) = \Sigma(i)\text{-}Id\Sigma(i)\text{-}ModL(i)\},$$

that is, $\mathcal{L}(i)$ is the class of all Σ -equational theories of sort i . Then $\mathcal{L}(i)$ forms a complete lattice of Σ -equational theories of sort i . Let $\mathcal{Ly}(i)$ be the class of all fixed points with respect to the closure operator $H\Sigma(i)\text{-}IdH\Sigma(i)\text{-}Mod$:

$$\mathcal{Ly}(i) := \{L(i) \subseteq W(i)^2 \mid L(i) = H\Sigma(i)\text{-}IdH\Sigma(i)\text{-}ModL(i)\},$$

that is, $\mathcal{Ly}(i)$ is the class of all solid Σ -equational theories of sort i . Then $\mathcal{Ly}(i)$ forms a complete lattice of solid Σ -equational theories of sort i and $\mathcal{Ly}(i)$ is a complete sublattice of $\mathcal{L}(i)$. We set $\mathcal{L} := (\mathcal{L}(i))_{i \in I}$ and $\mathcal{Ly} := (\mathcal{Ly}(i))_{i \in I}$. \mathcal{L} is called an I -sorted complete lattice. \mathcal{Ly} is called an I -sorted complete sublattice of \mathcal{L} , since for every $i \in I$, $\mathcal{Ly}(i)$ is a complete sublattice of $\mathcal{L}(i)$.

REFERENCES

- [1] P. Baltazar, *M-Solid Varieties of Languages*, Acta Cybernetica **18** (2008) 719–731.
- [2] K. Denecke and S.L. Wismath, *Hyperidentities and Clones*, Gordon and Breach 2000.
- [3] K. Denecke and S. Lekkoksung, *Hypersubstitutions of Many-Sorted Algebras*, Asian-European J. Math. Vol. **I** (3) (2008) 337–346.

- [4] J. Koppitz and K. Denecke, *M-solid Varieties of Algebras*, Springer 2005.
- [5] H. Lugowski, *Grundzüge der Universellen Algebra*, Teubner-Verlag, Leipzig 1976.

Received 20 May 2009

Revised 10 September 2009