

A NOTE ON HYPERVECTOR SPACES

SANJAY ROY

Department of Mathematics
South Bantra Ramkrishna Institution
West Bengal, India

e-mail: sanjappuremath@gmail.com

AND

TAPAS K. SAMANTA

Department of Mathematics, Uluberia College
West Bengal, India

e-mail: mumpu_tapas5@yahoo.co.in

Abstract

The main aim of this paper is to generalize the concept of vector space by the hyperstructure. We generalize some definitions such as hypersubspaces, linear combination, Hamel basis, linearly dependence and linearly independence. A few important results like deletion theorem, extension theorem, dimension theorem have been established in this hypervector space.

Keywords: hyperoperation, hyperfield, hypervector spaces, linear dependent, linear independent.

2010 Mathematics Subject Classification: 14L17, 20N20.

1. INTRODUCTION

The concept of hyperstructure was first introduced by Marty [4] in 1934 at the 8th congress of scandinavian Mathematicians and then he established the definition of hypergroup [2] in 1935 to analyse its properties and applied them to groups of rational algebraic functions. Also he was motivated to introduce this structure to study several problems of the non-commutative algebra. Then several researchers have been working on this new field of modern algebra and developed it. M. Krasner [1], a great researcher in this area, introduced the notions of hyperring and hyperfield to use it as a technical tool in the study of the approximation of valued fields. Later on it has been developed and generalized by other researchers. Then the notion of the hypervector spaces was introduced by M. Scafati Tallini [8] in 1988.

In the definition [16] of hypervector spaces, M. Scafati Tallini has considered the field as a usual field. In this paper, we have generalized the definition of hypervector space by considering the field as a hyperfield and considering the multiplication structure of a vector by a scalar as hyperstructure like M. Scafati Tallini. We again call it a hypervector space. Then we have established a few basic properties in this hypervector space and thereafter the notions of linear combinations, linearly dependence, linearly independence, Hamel basis, etc. are introduced and several important properties like deletion theorem, extension theorem etc. are developed.

2. PRELIMINARIES

We quote some definitions and proofs of a few results which will be needed in the sequel.

Definition 2.1 [5]. A *hyperoperation* over a non-empty set X is a mapping of $X \times X$ into the set of all non-empty subsets of X .

Definition 2.2 [5]. A non-empty set X with exactly one hyperoperation ' $\#$ ' is a *hypergroupoid*.

Let $(X, \#)$ be a hypergroupoid. For every point $x \in X$ and every non-empty subset A of X , we define $x \# A = \bigcup_{a \in A} \{x \# a\}$.

Definition 2.3 [5]. A hypergroupoid $(X, \#)$ is called a *semihypergroup* if $x \# (y \# z) = (x \# y) \# z$ for all $x, y, z \in X$.

Definition 2.4 [5]. A hypergroupoid $(X, \#)$ is called a *hypergroup* if

- (i) $x \# (y \# z) = (x \# y) \# z$.
- (ii) $\exists 0 \in X$ such that for every $a \in X$, there is unique element $b \in X$ for which $0 \in a \# b$ and $0 \in b \# a$. Here b is denoted by $-a$.
- (iii) For all $a, b, c \in X$ if $a \in b \# c$, then $b \in a \# (-c)$.

Result 2.5 [5]. In a hypergroup $(X, \#)$, $-(-a) = a$, $\forall a \in X$.

Proof. Since $0 \in a \# (-a)$ and $0 \in (-a) \# a$, $\forall a \in X$ i.e $0 \in (-a) \# a$ and $0 \in a \# (-a)$, $\forall a \in X$. Hence $-(-a) = a$, $\forall a \in X$.

Result 2.6 [5]. $0 \# a = \{a\}$, $\forall a \in X$, if $(X, \#)$ is a commutative hypergroup.

Proof. Let $a \in X$, then $0 \in a \# (-a)$

$$\Rightarrow a \in 0 \# (-(-a)) \text{ [by Definition 2.4]}$$

$$\Rightarrow a \in 0 \# a$$

we now show that $0 \# a = \{a\}$.

Let $b \in 0 \# a$

$$\Rightarrow 0 \in b \# (-a) = (-a) \# b$$

$$\Rightarrow b = -(-a)$$

$$\Rightarrow b = a.$$

This completes the proof.

Result 2.7 [5]. In a commutative hypergroup $(X, \#)$, 0 is unique.

Proof. Let there be another element $e \in X$ such that

$$e \in a \# (-a), e \in (-a) \# a, \forall a \in X.$$

$$\Rightarrow a \# e = \{a\}, \forall a \in X.$$

$$\Rightarrow 0 \# e = \{0\}. \text{ Again } 0 \# e = \{e\}.$$

Hence $e = 0$.

This completes the proof.

Note 2.8. In a hypergroup, if the element 0 is unique, then 0 is called the zero element of the hypergroup and b is called the additive inverse of a if $0 \in a \# b$ and $0 \in b \# a$.

3. HYPERVECTOR SPACE

Definition 3.1. A *hyperring* is a non-empty set endowed with a hyper-addition ' \oplus ' and a multiplication ' \cdot ' such that (X, \oplus) is a commutative hypergroup and (X, \cdot) is a semigroup and the multiplication is distributive with respect of the hyperaddition, both from the left and from the right side and $a \cdot 0 = 0 \cdot a = 0, \forall a \in X$, where 0 is the zero element of the hyperring.

Definition 3.2. A *hyperfield* is a non-empty set X endowed with a hyper-addition ' \oplus ' and a multiplication ' \cdot ' such that

- (i) (X, \oplus, \cdot) is a hyperring.
- (ii) \exists an element $1 \in X$, called the identity element such that $a \cdot 1 = a, \forall a \in X$.
- (iii) For each non zero element $a \in X, \exists$ an element $a^{-1} \in X$ such that $a \cdot a^{-1} = 1$.
- (iv) $a \cdot b = b \cdot a, \forall a, b \in X$.

Definition 3.3. Let (F, \oplus, \cdot) be a hyperfield and $(V, \#)$ be an additive commutative hypergroup. Then V is said to be a *hypervector space* over the hyperfield F if there exist a hyperoperation $*$: $F \times V \rightarrow P^*(V)$ such that

- (i) $a * (\alpha \# \beta) \subseteq a * \alpha \# a * \beta, \quad \forall a \in F \text{ and } \forall \alpha, \beta \in V.$
- (ii) $(a \oplus b) * \alpha \subseteq a * \alpha \# b * \alpha, \quad \forall a, b \in F \text{ and } \forall \alpha \in V.$
- (iii) $(a \cdot b) * \alpha = a * (b * \alpha), \quad \forall a, b \in F \text{ and } \forall \alpha \in V.$
- (iv) $1_F * \alpha = \{\alpha\} \text{ and } 0 * \alpha = \{\theta\}, \quad \forall \alpha \in V$ where 1_F is the identity element of F , 0 is the zero element of F and θ is the zero vector of V and $P^*(V)$ is the set of all non-empty subsets of V .

A hypervector space is called *strongly right distributive hypervector space* (respectively, *strongly left distributive hypervector space*), if equality holds in (i) (respectively, in (ii)).

A hypervector space is called a *good hypervector space* if equality holds in both (i) and (ii).

Remark 3.4. By a hypervector space V , we mean a hypervector space $(V, \#, *)$ and by a hyperfield F , we mean a hyperfield (F, \oplus, \cdot) .

Remark 3.5. Let V be a hypervector space over a hyperfield F . Let $a, b \in F$ and $\alpha, \beta \in V$, then by $a * \alpha \# b * \beta$, we mean $(a * \alpha) \# (b * \beta)$.

Example 3.6. Let (F, \oplus, \cdot) be an hyperfield and $V = F \times F$. Let us define a hyperoperation $\#'$ on V as follows

$$(a_1, a_2) \# (b_1, b_2) = (a_1 \oplus b_1, a_2 \oplus b_2) = \{(x, y) : x \in a_1 \oplus b_1 \text{ and } y \in a_2 \oplus b_2\}.$$

Then we prove that $(V, \#)$ is an additive commutative hypergroup. Now we define a scalar multiplication $*$: $F \times V \rightarrow P^*(V)$ by $a * ((a_1, a_2)) = \{(a \cdot a_1, a \cdot a_2)\}$, where $a \in F$ and $(a_1, a_2) \in V$. Then we easily verify that

- (i) $a * ((a_1, a_2) \# (b_1, b_2)) = (a * (a_1, a_2)) \# (a * (b_1, b_2))$,
- (ii) $(a \oplus b) * (a_1, a_2) = (a * (a_1, a_2)) \# (b * (a_1, a_2))$,
- (iii) $(a \cdot b) * (a_1, a_2) = a * (b * (a_1, a_2))$,
- (iv) $1_F * (a_1, a_2) = (a_1, a_2)$ and $0 * (a_1, a_2) = (0, 0) = \theta$,

for all $a, b \in F$ and for all $(a_1, a_2), (b_1, b_2) \in V$.

Result 3.7. Let $(V, \#, *)$ be a hypervector space over a hyperfield (F, \oplus, \cdot) . Then

- (i) $k * \theta = \{\theta\}$, $\forall k \in F, \theta$ being the zero vector of V .
- (ii) Let $k \in F$ and $\alpha \in V$ be such that $k * \alpha = \{\theta\}$, then either $k = 0$, or $\alpha = \theta$.
- (iii) $-\alpha \in (-1_F) * \alpha$, $\forall \alpha \in V, 1_F$ being the identity element of F .

Proof.

$$(i) \quad k * \theta = k * (0 * \theta),$$

[by axiom (iv), we have $0 * \alpha = \theta$, $\forall \alpha \in V$] $= (k \cdot 0) * \theta = 0 \cdot \theta = \theta$.

$$(ii) \quad \text{Let } k \in F \text{ and } \alpha \in V \text{ be such that } k * \alpha = \{\theta\}.$$

If $k = 0$, then $0 * \alpha = \theta$.

If $k \neq 0$, then $k^{-1} \in F$.

$$\text{Therefore } k * \alpha = \theta \Rightarrow k^{-1} * (k * \alpha) = k^{-1} * \theta$$

$$\Rightarrow (k^{-1} \cdot k) * \alpha = \theta \Rightarrow 1_F * \alpha = \theta \Rightarrow \alpha = \theta.$$

This completes the proof.

$$(iii) \quad \text{Let } \alpha \in V, \text{ then}$$

$$(1_F \oplus (-1_F)) * \alpha \subseteq 1_F * \alpha \# (-1_F) * \alpha = \alpha \# (-1_F) * \alpha.$$

$$\text{Since } 0 \in 1_F \oplus (-1_F) \Rightarrow \theta = 0 * \alpha \in (1_F \oplus (-1_F)) * \alpha$$

$$\Rightarrow \theta \in \alpha \# (-1_F) * \alpha.$$

$$\text{Therefore } -\alpha \in (-1_F) * \alpha, \forall \alpha \in V.$$

4. HYPERSUBSPACES

Definition 4.1. A subset W of a hypervector space V over a hyperfield F is called a *hypersubspace* of V if W is a hypervector space over F with the hyperoperations of addition and the scalar multiplication defined on V .

Therefore a subset W of a hypervector space V is a hypersubspace of V if and only if the following four properties hold.

$$(i) \quad \alpha \# \beta \subseteq W, \quad \forall \alpha, \beta \in W,$$

$$(ii) \quad a * \alpha \subseteq W, \quad \forall \alpha \in W \text{ and } \forall a \in F,$$

$$(iii) \quad W \text{ has a zero vector,}$$

$$(iv) \quad \text{each vector of } W \text{ has an additive inverse.}$$

Theorem 4.2. *Let V be a hypervector space and W is a subset of V . Then W is a hypersubspace of V if and only if the following three conditions hold:*

- (i) W is non-empty.
- (ii) $\alpha \# \beta \subseteq W, \forall \alpha, \beta \in W$.
- (iii) $a * \alpha \subseteq W, \forall a \in F$ and $\forall \alpha \in W$.

Proof. If W is a hypersubspace of V , then obviously the conditions (i), (ii) and (iii) hold.

Conversely, let W be a subset of V such that W satisfies the three conditions (i), (ii) and (iii).

To proof that W is a hypersubspace of V . it is enough to prove that (1) W has a zero vector. (2) Each vector in W has an additive inverse. Since W is non-empty, let $\alpha \in W$. Now $0 \in F$, therefore by the condition (iii) we get $0 * \alpha \subseteq W \Rightarrow \theta \in W$. Therefore W has a zero vector. Again, since $-1_F \in F$, therefore $(-1_F) * \alpha \subseteq W \Rightarrow -\alpha \in W$. Hence each vector in W has an additive inverse.

Theorem 4.3. *W be a hypersubspace of a hypervector space V if and only if (i) W is non-empty. (ii) $a * \alpha \# b * \beta \subseteq W, \forall a, b \in F$ and $\forall \alpha, \beta \in W$.*

Proof. If W is a hypersubspace of V , then obviously W satisfies the conditions (i) and (ii). Conversely, let W satisfy the conditions (i) and (ii).

Since $1_F \in F$, let $\alpha, \beta \in W$, then by (ii) $1_F * \alpha \# 1_F * \beta \subseteq W \Rightarrow \alpha \# \beta \subseteq W$. Let $a \in F$ and $\alpha, \beta \in V$, since $0 \in F$, therefore by (ii) $a * \alpha \# 0 * \beta \subseteq W \Rightarrow a * \alpha \# \theta \subseteq W \Rightarrow a * \alpha \subseteq W$ [because $(W, \#)$ is a commutative hypergroup]. Hence W is a hypersubspace of V .

Example 4.4. Let (F, \oplus, \cdot) be a hyperfield and $V = F \times F$.

Then $(V, \#, *)$ is a hypervector space, where the hyperoperations ' $\#$ ' and ' $*$ ' are defined by $(a_1, a_2) \# (b_1, b_2) = (a_1 \oplus b_1, a_2 \oplus b_2) = \{(x, y) : x \in a_1 \oplus b_1 \text{ and } y \in a_2 \oplus b_2\}$ and $a * (a_1, a_2) = \{(a.a_1, a.a_2)\}$, $\forall (a_1, a_2), (b_1, b_2) \in V$ and $\forall a \in F$.

Let $W = F \times \{0\} \subseteq V$. We now show that W is a hypersubspace of V , since $\theta = (0, 0) \in W$. Now let $\alpha = (a_1, 0)$, $\beta = (b_1, 0) \in W$ and $a, b \in F$.

Then

$$\begin{aligned}
& a * \alpha \# b * \beta \\
&= a*(a_1, 0) \# b*(b_1, 0) \\
&= \{(a.a_1, a.0)\} \# \{(b.b_1, b.0)\} \\
&= \{(a.a_1, 0)\} \# \{(b.b_1, 0)\} \\
&= (a.a_1 \oplus b.b_1, 0 \oplus 0) \\
&= (a.a_1 \oplus b.b_1, 0) \subseteq W \text{ [because } a.a_1 \oplus b.b_1 \subseteq F].
\end{aligned}$$

Theorem 4.5. *The intersection of two hypersubspaces of a hypervector space V over a hyperfield F is again a hypersubspace of V .*

Proof. Obvious.

Theorem 4.6. *The intersection of any family of hypersubspaces of a hypervector space V over a hyperfield F is again a hypersubspace of V .*

Proof. Obvious.

In the above example, we take $W_1 = F \times \{0\} \subseteq V$ and $W_2 = \{0\} \times F \subseteq V$. Then by the same procedure of the above example, we can prove that W_1 and W_2 are hypersubspaces of V .

Let $(a, 0) \in W_1$ and $(0, b) \in W_2$.

Then $(a, 0) \# (0, b) = (a \oplus 0, 0 \oplus b) = (a, b)$.

But $\{(a, b)\}$ is not a subset of $W_1 \cup W_2$.

Therefore $W_1 \cup W_2$ is not a hypersubspace of V .

Note 4.7. *The union of two hypersubspaces of a hypervector space V is not necessarily a hypersubspace of V .*

Theorem 4.8. *Let W_1 and W_2 be two hyperspaces of a hypervector space V . Then we prove that $W_1 \# W_2 = \cup\{\alpha \# \beta, \alpha \in W_1, \beta \in W_2\}$ is a hypersubspace of V .*

Proof. Since $\theta \in W_1$ and $\theta \in W_2$. Then $\{\theta \# \theta\} \subseteq W_1 \# W_2 \Rightarrow \{\theta\} \subseteq W_1 \# W_2 \Rightarrow \theta \in W_1 \# W_2$, therefore $W_1 \# W_2$ is non-empty.

Let $\alpha, \beta \in W_1 \# W_2$, Then $\exists \alpha_1, \alpha_2 \in W_1$ and $\beta_1, \beta_2 \in W_2$ such that $\alpha \in \alpha_1 \# \beta_1$ and $\beta \in \alpha_2 \# \beta_2$. Let $a, b \in F$. Now

$$\begin{aligned}
 a * \alpha \# b * \beta &\subseteq a * (\alpha_1 \# \beta_1) \# b * (\alpha_2 \# \beta_2) \\
 &\subseteq (a * \alpha_1 \# a * \beta_1) \# (b * \alpha_2 \# b * \beta_2) \\
 &= a * \alpha_1 \# (a * \beta_1 \# b * \alpha_2) \# b * \beta_2 \\
 &= a * \alpha_1 \# (b * \alpha_2 \# a * \beta_1) \# b * \beta_2 \\
 &= (a * \alpha_1 \# b * \alpha_2) \# (a * \beta_1 \# b * \beta_2) \\
 &\subseteq W_1 \# W_2.
 \end{aligned}$$

Hence $W_1 \# W_2$ is a hypersubspace of V .

Definition 4.9. If W_1 and W_2 are two hypersubspaces of a hypervector space V , then the hypersubspace $W_1 \# W_2$ is called the *hyperlinear sum* or *linear sum* of the hyperspaces W_1 and W_2 .

If $W_1 \cap W_2 = \{\theta\}$ then $W_1 \# W_2$ is called the *direct sum* of the hypersubspaces W_1 and W_2 .

Theorem 4.10. *The hypersubspace $W_1 \# W_2$ is the smallest hypersubspace of V containing the hypersubspaces W_1 and W_2 .*

Proof. Let W be a hypersubspace of V such that $W_1 \subseteq W$ and $W_2 \subseteq W$. Let $\gamma \in W_1 \# W_2$, then $\exists \alpha \in W_1$ and $\beta \in W_2$ such that $\gamma \in \alpha \# \beta$. Since $W_1 \subseteq W$ and $W_2 \subseteq W$. Therefore $\alpha, \beta \in W$. Again since W is a hypersubspace of V .

Therefore $\alpha \# \beta \subseteq W \Rightarrow \gamma \in W$.

Hence $W_1 \# W_2 \subseteq W$

This completes the proof.

If V is a strongly left distributive hypervector space over a hyperfield F , then it can be easily prove that $W = \cup\{a * \alpha, a \in F\}$ forms a hypersubspace of V , where $\alpha \in V$. This hypersubspace is said to be generated by the vector α and α is said to be a *generator* of the hypersubspace W . This hypersubspace is usually denoted by $HL(\alpha)$.

Then from the previous theorem it can be easily proved that if $\alpha, \beta \in V$, then the set $W = \cup\{a * \alpha \# b * \beta, a, b \in F\}$ is a hypersubspace of V . This hypersubspace is called the *hyperlinear span* of the vectors α and β , it is usually denoted by $HL(\alpha, \beta)$.

Theorem 4.11. *Let V be a strongly left distributive hypervector space over the hyperfield F and $\alpha_1, \alpha_2, \dots, \alpha_n \in V$. Then*

$$W = \cup\{a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n / a_1, a_2, \dots, a_n \in F\}$$

is a hypersubspace of V . In fact W is the smallest hypersubspace of V containing $\alpha_1, \alpha_2, \dots, \alpha_n$.

Proof. Since F is non-empty, therefore W is non-empty. Let $w_1, w_2 \in W$. Then $\exists a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n \in F$ such that

$$w_1 \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n$$

and

$$w_2 \in b_1 * \alpha_1 \# b_2 * \alpha_2 \# \dots \# b_n * \alpha_n,$$

therefore

$$w_1 \# w_2 \subseteq (a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n) \# (b_1 * \alpha_1 \# b_2 * \alpha_2 \# \dots \# b_n * \alpha_n)$$

$$\text{or } w_1 \# w_2 \subseteq (a_1 * \alpha_1 \# b_1 * \alpha_1) \# (a_2 * \alpha_2 \# b_2 * \alpha_2) \# \dots \# (a_n * \alpha_n \# b_n * \alpha_n)$$

$$\text{or } w_1 \# w_2 \subseteq (a_1 \oplus b_1) * \alpha_1 \# (a_2 \oplus b_2) * \alpha_2 \# \dots \# (a_n \oplus b_n) * \alpha_n.$$

Therefore $w_1 \# w_2 \subseteq W$.

Next let $w \in W$ and $a \in F$.

Then $\exists a_1, a_2, \dots, a_n \in F$ such that

$$w \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n.$$

$$\text{Therefore } a * w \subseteq a * (a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n)$$

$$\subseteq a * (a_1 * \alpha_1) \# a * (a_2 * \alpha_2) \# \dots \# a * (a_n * \alpha_n)$$

$$= (a.a_1) * \alpha_1 \# (a.a_2) * \alpha_2 \# \dots \# (a.a_n) * \alpha_n \subseteq W.$$

Therefore $a * w \subseteq W$, $\forall a \in F$ and $\forall w \in W$.

Hence W is a hypersubspace of V .

Next, since $0, 1 \in F$. Let $i \in \{1, 2, 3, \dots, n\}$.

Therefore

$$0 * \alpha_1 \# 0 * \alpha_2 \# \dots \# 0 * \alpha_{i-1} \# 1 * \alpha_i \# 0 * \alpha_{i+1} \# \dots \# 0 * \alpha_n \subseteq W$$

$$\Rightarrow \theta \# \theta \# \dots \# \theta \# \alpha_i \# \theta \# \dots \# \theta \subseteq W$$

$$\Rightarrow \alpha_i \in W, [\text{As } \theta \# \theta = \{\theta\} \text{ and } \theta \# \alpha_i = \alpha_i \# \theta = \alpha_i],$$

therefore $\alpha_i \in W$, for all $i = 1, 2, \dots, n$.

Let P be a hypersubspace of V containing $\alpha_1, \alpha_2, \dots, \alpha_n$.

Let $\alpha \in W$, then $\exists a_1, a_2, \dots, a_n \in F$ such that

$$\alpha \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n \subseteq P.$$

Therefore $\alpha \in P \Rightarrow W \subseteq P$.

Hence W is the smallest hyperspace of V containing $\alpha_1, \alpha_2, \dots, \alpha_n$.

Note 4.12. The linear combination of a null vector is the set $\{\theta\}$.

5. LINEAR DEPENDENCE AND LINEAR INDEPENDENCE

Definition 5.1. Let V be a hypervector space over a hyperfield F and $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a finite subset of V . Then S is said to be *linearly dependent* if there exist the scalars $a_1, a_2, \dots, a_n \in F$ (not all zero) such that $\theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n$.

Otherwise S is said to be *linearly independent*. i.e, if S is linearly independent and $\theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n$.

Then $a_1 = a_2 = \dots = a_n = 0$

Definition 5.2. Let V be a hypervector space over a hyperfield F and $S \subseteq V$. Then S is said to be linearly dependent if S contains a finite subset which is linearly dependent. Otherwise S is linearly independent.

Result 5.3. Any singleton set of non-null vector of a hypervector space V is linearly independent.

Proof. Let α be a non-null vector of a hypervector space V . If possible let $\theta \in a * \alpha$ for some $a \in F$. We now show that $a = 0$. If $a \neq 0$, then $a^{-1} \in F$.

Now, since $\theta \in a * \alpha \Rightarrow a^{-1} * \theta \subseteq a^{-1} * (a * \alpha)$

$$\Rightarrow \theta \in a^{-1} * (a * \alpha) \quad [\text{As } a^{-1} * \theta = \theta]$$

$$\Rightarrow \theta \in (a^{-1}.a) * \alpha$$

$$\Rightarrow \theta \in 1_F * \alpha$$

$$\Rightarrow \theta \in \{\alpha\} \Rightarrow \alpha = \theta, \text{ which is a contradiction.}$$

Hence $a = 0$. This completes the proof.

Result 5.4. Any set of vectors containing the null vector is always linearly dependent.

Proof. Obvious.

Definition 5.5. Let V be a hypervector space over a hyperfield F . Then the vector $\alpha \in V$ is said to be a *linear combination* of the vectors $\alpha_1, \alpha_2, \dots, \alpha_n \in V$ if there exist $a_1, a_2, \dots, a_n \in F$ such that

$$\alpha \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n.$$

Theorem 5.6. Let V be a hypervector space over a hyperfield F and $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a subset of V . Then S is linearly dependent if and only if at least one of S can be expressed as a linear combination of the remaining other members of S .

Proof. Let S is linearly dependent. Then there exist a_1, a_2, \dots, a_n (not all zero) $\in F$ such that

$$(i) \quad \theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n, \quad \dots$$

since $\#$ is commutative. Therefore without loss of generality we assume that $a_1 \neq 0$. Then $a_1^{-1} \in F$. Therefore from (i) we get

$$\begin{aligned}
a_1^{-1} * \theta &\subseteq a_1^{-1} * (a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n) \\
\Rightarrow \{\theta\} &\subseteq \{a_1^{-1} * (a_1 * \alpha_1)\} \# \{a_1^{-1} * (a_2 * \alpha_2)\} \# \dots \# \{a_1^{-1} * (a_n * \alpha_n)\} \\
&= (a_1^{-1}.a_1) * \alpha_1 \# (a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n \\
&= 1_F * \alpha_1 \# (a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n \\
&= \alpha_1 \# (a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n \\
&= \alpha_1 \# ((a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n).
\end{aligned}$$

Then \exists an element $\beta \in (a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n$ such that $\theta \in \alpha_1 \# \beta = \beta \# \alpha_1$.

This implies that $\alpha_1 = -\beta$. So $\alpha_1 \in (-1_F) * \beta$. Therefore

$$\begin{aligned}
\alpha_1 \in (-1_F) * \beta &\subseteq (-1_F) * ((a_1^{-1}.a_2) * \alpha_2 \# \dots \# (a_1^{-1}.a_n) * \alpha_n) \\
&\subseteq (-1_F) * ((a_1^{-1}.a_2) * \alpha_2) \# \dots \# (-1_F) * ((a_1^{-1}.a_n) * \alpha_n) \\
&= ((-1_F).(a_1^{-1}.a_2)) * \alpha_2 \# \dots \# ((-1_F).(a_1^{-1}.a_n)) * \alpha_n \\
&= (-a_1^{-1}.a_2) * \alpha_2 \# \dots \# (-a_1^{-1}.a_n) * \alpha_n
\end{aligned}$$

i.e $\alpha_1 \in (-a_1^{-1}.a_2) * \alpha_2 \# \dots \# (-a_1^{-1}.a_n) * \alpha_n \Rightarrow \alpha_1 \in HL(\alpha_2, \alpha_2, \dots, \alpha_n)$.

This completes the proof of the necessary part of the theorem.

Converse part.

Without loss of generality we assume that $\alpha_1 \in Hl(\alpha_2, \alpha_3, \dots, \alpha_n)$.

Then $\exists a_2, a_3, \dots, a_n \in F$ such that

$$\begin{aligned}
 & \alpha_1 \in a_2 * \alpha_2 \# a_3 * \alpha_3 \# \dots \# a_n * \alpha_n \\
 & \Rightarrow -\alpha_1 \# \alpha_1 \subseteq -\alpha_1 \# a_2 * \alpha_2 \# a_3 * \alpha_3 \# \dots \# a_n * \alpha_n \\
 & \Rightarrow \theta \in -\alpha_1 \# a_2 * \alpha_2 \# a_3 * \alpha_3 \# \dots \# a_n * \alpha_n \text{ [as } \theta \in -\alpha_1 \# \alpha_1 \text{]} \\
 & \Rightarrow \theta \in (-1_F) * \alpha_1 \# a_2 * \alpha_2 \# a_3 * \alpha_3 \# \dots \# a_n * \alpha_n \\
 & \Rightarrow \theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n, \text{ where } a_1 = -1_F (\neq 0).
 \end{aligned}$$

Therefore the set $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is linearly dependent.

Theorem 5.7. *The non-zero vectors $\alpha_1, \alpha_2, \dots, \alpha_n$ of a hypervector space are linearly dependent if and only if one of them, say α_i , is a linear combination of the previous vectors, i.e $\alpha_i \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_{i-1} * \alpha_{i-1}$, for some $a_1, a_2, \dots, a_{i-1} \in F$.*

Proof. First we suppose that $\alpha_1, \alpha_2, \dots, \alpha_n$ are linearly dependent. Then we get a set of scalars a_1, a_2, \dots, a_n (not all zero) $\in F$ such that

$$(i) \quad \theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n \quad \dots$$

Let k be the largest integer such that $a_k \neq 0$, here we see that $k \neq 1$.

If $k = 1$, then we see that $\alpha_1 = \theta$, which contradicts the fact that $\alpha_1, \alpha_2, \dots, \alpha_n$ are non-zero vectors. Since k be the largest integer such that $a_k \neq 0$, it follows that $a_i = 0$ for all $i, k < i \leq n$. Hence (i) reduces to the following form

$$\begin{aligned}
 & \theta \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_k * \alpha_k \\
 & \Rightarrow \alpha_k \in (-a_k^{-1} . a_1) * \alpha_1 \# (-a_k^{-1} . a_2) * \alpha_2 \# \dots \# (-a_k^{-1} . a_{k-1}) * \alpha_{k-1},
 \end{aligned}$$

[By the procedure of the proof of the Theorem 5.6] conversely, we suppose that $\alpha_i \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_{i-1} * \alpha_{i-1}$, for some scalars $a_1, a_2, \dots, a_{i-1} \in F$.

Therefore the set $\{\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i\}$ of vectors is linearly dependent. So $\{\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i, \alpha_{i+1}, \dots, \alpha_n\}$ of vectors is also linearly dependent. This completes the proof.

Theorem 5.8 (Deletion Theorem). *Let V be a strongly left distributive hypervector space over the hyperfield F and V be generated by a linearly dependent set $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq V$. Then V can also be generated by a suitable proper subset of $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$.*

Proof. Since V is generated by $\alpha_1, \alpha_2, \dots, \alpha_n$, therefore we have $V = HL(\alpha_1, \alpha_2, \dots, \alpha_n)$.

Again, since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is linearly dependent. It follows that one of the vectors $\alpha_1, \alpha_2, \dots, \alpha_n$, say α_i , can be expressed as a linear combination of the remaining others.

Then we get the scalars $c_1, c_2, \dots, c_{i-1}, c_{i+1}, \dots, c_n \in F$ such that

$$\alpha_i \in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_{i-1} * \alpha_{i-1} \# c_{i+1} * \alpha_{i+1} \# \dots \# c_n * \alpha_n.$$

We now show that

$$HL(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n) = HL(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i, \alpha_{i+1}, \dots, \alpha_n).$$

It is obvious that

$$HL(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n) \subseteq HL(\alpha_1, \alpha_2, \dots, \alpha_i, \dots, \alpha_n) = V.$$

Let $\alpha \in V$, then there exist the scalars a_1, a_2, \dots, a_n such that

$$\begin{aligned} \alpha &\in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_i * \alpha_i \# \dots \# a_n * \alpha_n \\ \Rightarrow \alpha &\in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_{i-1} * \alpha_{i-1} \# a_i * (c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \\ &\# c_{i-1} * \alpha_{i-1} \# c_{i+1} * \alpha_{i+1} \# \dots \# c_n * \alpha_n) \# a_{i+1} * \alpha_{i+1} \# \dots \# a_n * \alpha_n \\ \Rightarrow \alpha &\in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_{i-1} * \alpha_{i-1} \# (a_i \cdot c_1) * \alpha_1 \# (a_i \cdot c_2) * \alpha_2 \# \dots \# \\ &(a_i \cdot c_{i-1}) * \alpha_{i-1} \# (a_i \cdot c_{i+1}) * \alpha_{i+1} \# \dots \# (a_i \cdot c_n) * \alpha_n \# a_{i+1} * \alpha_{i+1} \# \dots \# a_n * \alpha_n \\ \Rightarrow \alpha &\in (a_1 \oplus a_i \cdot c_1) * \alpha_1 \# (a_2 \oplus a_i \cdot c_2) * \alpha_2 \# \dots \# (a_{i-1} \oplus a_i \cdot c_{i-1}) * \alpha_{i-1} \# \\ &(a_{i+1} \oplus a_i \cdot c_{i+1}) * \alpha_{i+1} \# \dots \# (a_n \oplus a_i \cdot c_n) * \alpha_n. \end{aligned}$$

Hence $\alpha \in \text{HL}(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n)$.

Therefore $V = \text{HL}(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n)$.

This completes the proof.

Theorem 5.9. *If $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a linearly independent set of generators of a hypervector space V , then no proper subset of S can be a spanning set of V .*

Proof. Obvious.

Theorem 5.10. *Suppose $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ generates a hypervector space V . If $\{\beta_1, \beta_2, \dots, \beta_m\}$ is linearly independent, then $m \leq n$ and V is generated by a set of the form $\{\beta_1, \beta_2, \dots, \beta_m, \alpha_{i_1}, \alpha_{i_2}, \dots, \alpha_{i_{n-m}}\}$.*

Proof. Obvious.

6. BASIS OR HAMEL BASIS

Definition 6.1 Let V be a hypervector space over the hyperfield F and S be a subset of V . S is said to be a *basis*, or *Hamel basis* if

- (i) S is linearly independent.
- (ii) Every elements of V can be expressed as a finite linear combination of a few elements of S .

If S is a basis of the hypervector space V and S is finite, then the hypervector space V is said to be a finite dimensional hypervector space and the number of elements in S is called the dimension of the hypervector space V . Usually the dimension of V is denoted by $\dim(V)$.

Again if S is infinite Set, then V is said to be an infinite dimensional hypervector space.

Theorem 6.2. *If $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a basis of a finite dimensional strongly left distributive hypervector space V over a hyperfield F , then every non-null vector $\alpha \in V$ has a unique representation.*

Proof. Since S is a basis of V and $\alpha \in V$, there exist $a_1, a_2, \dots, a_n \in F$ such that

$$(i) \quad \alpha \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n. \quad \dots$$

If possible let $\alpha \in b_1 * \alpha_1 \# b_2 * \alpha_2 \# \dots \# b_n * \alpha_n$, for some $b_1, b_2, \dots, b_n \in F$.

Therefore

$$\begin{aligned} -\alpha &\in (-1_F * \alpha) \subseteq (-1_F) * (b_1 * \alpha_1 \# b_2 * \alpha_2 \# \dots \# b_n * \alpha_n) \\ \Rightarrow -\alpha &\in ((-1_F) * (b_1 * \alpha_1)) \# ((-1_F) * (b_2 * \alpha_2)) \# \dots \# ((-1_F) * (b_n * \alpha_n)) \\ &= ((-1_F).b_1) * \alpha_1 \# ((-1_F).b_2) * \alpha_2 \# \dots \# ((-1_F).b_n) * \alpha_n \\ &= (-b_1) * \alpha_1 \# (-b_2) * \alpha_2 \# \dots \# (-b_n) * \alpha_n. \end{aligned}$$

Therefore

$$(ii) \quad -\alpha \in (-b_1) * \alpha_1 \# (-b_2) * \alpha_2 \# \dots \# (-b_n) * \alpha_n. \quad \dots$$

From (i) and (ii) we get

$$\begin{aligned} \alpha \# -\alpha &\subseteq a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_n * \alpha_n \\ &\# (-b_1) * \alpha_1 \# (-b_2) * \alpha_2 \# \dots \# (-b_n) * \alpha_n \end{aligned}$$

Therefore

$$\theta \in \alpha \# -\alpha \subseteq (a_1 \oplus (-b_1)) * \alpha_1 \# (a_2 \oplus (-b_2)) * \alpha_2 \# \dots \# (a_n \oplus (-b_n)) * \alpha_n.$$

Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a basis of V and

$$\theta \in (a_1 \oplus (-b_1)) * \alpha_1 \# (a_2 \oplus (-b_2)) * \alpha_2 \# \dots \# (a_n \oplus (-b_n)) * \alpha_n.$$

Then $0 \in a_i \oplus (-b_i)$, for all $i = 1, 2, \dots, n$. Again (F, \oplus) is commutative.

Therefore

$$a_i = -(-b_i), \text{ for all } i = 1, 2, \dots, n \text{ i.e. } a_i = b_i, \text{ for all } i = 1, 2, \dots, n.$$

This completes the proof.

Theorem 6.3 (Extension Theorem). *A linearly independent set of vectors in a finite dimension hypervector space V over a hyperfield F is either a basis, or it can be extended to a basis of V .*

Proof. Let $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set of vectors in V .

Now $\text{HL}(S)$ being a smallest hypersubspace of V containing S . It therefore follows that $\text{HL}(S) \subseteq V$.

If $\text{HL}(S) = V$, then S is a basis of V .

If $\text{HL}(S)$ is a proper hypersubspace of V , we show that S can be extended to a basis of V .

Let $\alpha \in V \setminus \text{HL}(S)$ and $S_1 = \{\alpha_1, \alpha_2, \dots, \alpha_n, \alpha\}$. Now we consider the following relation:

$$(i) \quad \theta \in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_n * \alpha_n \# c * \alpha, \quad \dots$$

for some $c_1, c_2, \dots, c_n, c \in F$. We now claim that $c = 0$.

If $c \neq 0$, then $c^{-1} \in F$.

Therefore

$$\begin{aligned} c^{-1} * \theta &\subseteq c^{-1} * (c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_n * \alpha_n \# c * \alpha) \\ \Rightarrow \theta &\in (c^{-1}.c_1) * \alpha_1 \# (c^{-1}.c_2) * \alpha_2 \# \dots \# (c^{-1}.c_n) * \alpha_n \# (c^{-1}.c) * \alpha \\ \Rightarrow \theta &\in (c^{-1}.c_1) * \alpha_1 \# (c^{-1}.c_2) * \alpha_2 \# \dots \# (c^{-1}.c_n) * \alpha_n \# 1_F * \alpha \\ \Rightarrow \theta &\in (c^{-1}.c_1) * \alpha_1 \# (c^{-1}.c_2) * \alpha_2 \# \dots \# (c^{-1}.c_n) * \alpha_n \# \alpha. \end{aligned}$$

Then $\exists \beta \in (c^{-1}.c_1) * \alpha_1 \# (c^{-1}.c_2) * \alpha_2 \# \dots \# (c^{-1}.c_n) * \alpha_n$ such that $\theta \in \beta \# \alpha = \alpha \# \beta \Rightarrow \alpha = -\beta \in (-1_F * \beta)$.

Therefore

$$\alpha \in (-1_F) * ((c^{-1}.c_1) * \alpha_1 \# (c^{-1}.c_2) * \alpha_2 \# \dots \# (c^{-1}.c_n) * \alpha_n)$$

$$\begin{aligned} &\Rightarrow \alpha \in (-c^{-1}.c_1) * \alpha_1 \# (-c^{-1}.c_2) * \alpha_2 \# \dots \# (-c^{-1}.c_n) * \alpha_n \\ &\Rightarrow \alpha \in HL(\alpha_1, \alpha_2, \dots, \alpha_n) \end{aligned}$$

i.e $\alpha \in HL(S)$, which is contradiction, since $\alpha \in V \setminus HL(S)$.

Hence we see that $c = 0$.

Substituting $c = 0$ in (i) we get

$$\begin{aligned} &\theta \in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_n * \alpha_n \# 0 * \alpha \\ &\Rightarrow \theta \in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_n * \alpha_n \# \theta \\ &\Rightarrow \theta \in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_n * \alpha_n \\ &\Rightarrow c_1 = c_2 = \dots = c_n = 0, \text{ as } \alpha_1, \alpha_2, \dots, \alpha_n \text{ are linearly independent.} \end{aligned}$$

Thus we observe that S_1 is linearly independent.

If $HL(S_1) = V$, then S_1 is a basis of V and the theorem is proved.

If $HL(S_1)$ is a proper subspace of V , then again we can take a vector $\beta \in V \setminus HL(S_1)$ and proceed as before.

Since V is a finite dimensional hypervector space, after a finite number of steps we come to a finite set of linearly independent vectors which generate the hypervector space V .

This completes the proof.

Theorem 6.4. *If $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a maximal linearly independent subset of a hypervector space V , then $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a basis of V .*

Proof. Obvious.

Theorem 6.5. *Let U and W be two finite dimensional hypersubspace of a strongly left distributive hypervector space V over the hyperfield F , then $U \# W$ is also a finite dimensional hypersubspace of V and*

$$\dim(U \# W) = \dim(U) + \dim(W) - \dim(U \cap W).$$

Proof. Here we see that $U \cap W$ is a hypersubspace of both U and W . So $U \cap W$ is a finite dimensional hypersubspace of V . Let $\dim(U \cap W) = r$, $\dim(U) = m$ and $\dim(W) = n$. then we have $r \leq m, n$.

Again let $\{\alpha_1, \alpha_2, \dots, \alpha_r\}$ be a basis of $U \cap W$.

Since $\{\alpha_1, \alpha_2, \dots, \alpha_r\}$ is a linearly independent set of vectors in U and $r \leq m = \dim(U)$, then it follows that either $\{\alpha_1, \alpha_2, \dots, \alpha_r\}$ is a basis of U or it can be extended to a basis for U .

Let $\{\alpha_1, \alpha_2, \dots, \alpha_r, \beta_1, \beta_2, \dots, \beta_{m-r}\}$ be a basis for U .

By similar arguments we can suppose that $\{\alpha_1, \alpha_2, \dots, \alpha_r, \gamma_1, \gamma_2, \dots, \gamma_{n-r}\}$ is a basis of W .

Let $S = \{\alpha_1, \alpha_2, \dots, \alpha_r, \beta_1, \beta_2, \dots, \beta_{m-r}, \gamma_1, \gamma_2, \dots, \gamma_{n-r}\}$. We now show that S is a basis of $U \# W$.

First we consider the following relation

$$(i) \quad \begin{aligned} \theta \in & a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_r * \alpha_r \# b_1 * \beta_1 \# b_2 * \beta_2 \\ & \# \dots \# b_{m-r} * \beta_{m-r} \# c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r}, \quad \dots \end{aligned}$$

for some $a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_{m-r}, c_1, c_2, \dots, c_{n-r} \in F$.

Then

$$(ii) \quad \begin{aligned} \exists \alpha \in & a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_r * \alpha_r \\ & \# b_1 * \beta_1 \# b_2 * \beta_2 \# \dots \# b_{m-r} * \beta_{m-r} \quad \dots \end{aligned}$$

So $\theta \in \alpha \# c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r}$.

Then $\exists \beta \in c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r}$.

So $\theta \in \alpha \# \beta = \beta \# \alpha$, as $\#$ is commutative.

$$\begin{aligned}
& \Rightarrow -\alpha = \beta \\
\text{(iii)} \quad & \Rightarrow -\alpha \in c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r} \quad \dots
\end{aligned}$$

$$\begin{aligned}
& \Rightarrow (-1_F) * (-\alpha) \subseteq (-1_F) * (c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r}) \\
& \Rightarrow -(-\alpha) \in ((-1_F).c_1) * \gamma_1 \# ((-1_F).c_2) \\
\text{(iv)} \quad & * \gamma_2 \# \dots \# ((-1_F).c_{n-r}) * \gamma_{n-r} \\
& \Rightarrow \alpha \in (-c_1) * \gamma_1 \# (-c_2) * \gamma_2 \# \dots \# (-c_{n-r}) * \gamma_{n-r} \quad \dots
\end{aligned}$$

From (ii) and (iv) we see that $\alpha \in U \cap W$.

Since $\alpha \in U \cap W$ and $\{\alpha_1, \alpha_2, \dots, \alpha_r\}$ is a basis of $U \cap W$, we have

$$\text{(v)} \quad \alpha \in d_1 * \alpha_1 \# d_2 * \alpha_2 \# \dots \# d_r * \alpha_r, \quad \dots$$

for some $d_1, d_2, \dots, d_r \in F$.

Therefore from (iii) and (v) we get

$$\begin{aligned}
& \alpha \# (-\alpha) \subseteq d_1 * \alpha_1 \# d_2 * \alpha_2 \# \dots \# d_r \\
& * \alpha_r \# c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r} \\
& \Rightarrow \theta \in d_1 * \alpha_1 \# d_2 * \alpha_2 \# \dots \# d_r \\
& * \alpha_r \# c_1 * \gamma_1 \# c_2 * \gamma_2 \# \dots \# c_{n-r} * \gamma_{n-r} \\
& \Rightarrow d_1 = d_2 = \dots = d_r = c_1 = c_2 = \dots = c_{n-r} = 0, \\
& \text{as } \{\alpha_1, \alpha_2, \dots, \alpha_r, \gamma_1, \gamma_2, \dots, \gamma_{n-r}\}
\end{aligned}$$

is a linearly independent set of vectors.

Then the relation (i) reduces to the following

$$\begin{aligned} \theta &\in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_r \\ &\quad * \alpha_r \# b_1 * \beta_1 \# b_2 * \beta_2 \# \dots \# b_{m-r} * \beta_{m-r} \\ \Rightarrow a_1 &= a_2 = \dots = a_r = b_1 = b_2 = \dots = b_{m-r} = 0, \\ &\text{as } \{\alpha_1, \alpha_2, \dots, \alpha_r, \beta_1, \beta_2, \dots, \beta_{m-r}\} \end{aligned}$$

is a linearly independent set of vectors.

Thus S is a linearly independent set of vectors V .

We now show that S generates $U \# W$.

Let $\alpha \in U \# W$.

Then we have $\alpha \in \beta \# \gamma$, for some $\beta \in U$ and for some $\gamma \in W$.

Since $\beta \in U$ and $\{\alpha_1, \alpha_2, \dots, \alpha_r, \beta_1, \beta_2, \dots, \beta_{m-r}\}$ is a basis of U .

Therefore

$$\begin{aligned} \beta &\in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_r * \alpha_r \# b_1 * \beta_1 \# b_2 * \beta_2 \\ &\quad \# \dots \# b_{m-r} * \beta_{m-r}, \text{ for some } a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_{m-r} \in F. \end{aligned}$$

Again since $\gamma \in W$ and $\{\alpha_1, \alpha_2, \dots, \alpha_r, \gamma_1, \gamma_2, \dots, \gamma_{n-r}\}$ is a basis of W .

Therefore

$$\begin{aligned} \gamma &\in c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_r * \alpha_r \# d_1 * \gamma_1 \# d_2 * \gamma_2 \\ &\quad \# \dots \# d_{n-r} * \gamma_{n-r}, \text{ for some } c_1, c_2, \dots, c_r, d_1, d_2, \dots, d_{n-r} \in F. \end{aligned}$$

Now $\alpha \in \beta \# \gamma$.

Therefore

$$\begin{aligned}
& \alpha \in a_1 * \alpha_1 \# a_2 * \alpha_2 \# \dots \# a_r * \alpha_r \# b_1 * \beta_1 \# b_2 * \beta_2 \# \dots \# b_{m-r} * \beta_{m-r} \\
& \# c_1 * \alpha_1 \# c_2 * \alpha_2 \# \dots \# c_r * \alpha_r \# d_1 * \gamma_1 \# d_2 * \gamma_2 \# \dots \# d_{n-r} * \gamma_{n-r} \\
& \Rightarrow \alpha \in (a_1 \oplus c_1) * \alpha_1 \# (a_2 \oplus c_2) * \alpha_2 \# \dots \# (a_r \oplus c_r) * \alpha_r \# b_1 \\
& * \beta_1 \# b_2 * \beta_2 \# \dots \# b_{m-r} * \beta_{m-r} \# d_1 * \gamma_1 \# d_2 * \gamma_2 \# \dots \# d_{n-r} * \gamma_{n-r}.
\end{aligned}$$

This shows that every vector of $U \# W$ can be expressed as a linear combination of the vectors of S . Hence S is a basis of $U \# W$, which proves that $U \# W$ is a finite dimensional hypervector space of V .

Now

$$\begin{aligned}
\dim(U \# W) &= |S| \\
&= r + m - r + n - r \\
&= m + n - r \\
&= \dim(U) + \dim(W) - \dim(U \cap W).
\end{aligned}$$

This completes the proof.

Acknowledgements

The authors are grateful to the referees for their valuable suggestions in rewriting the paper in the present form.

REFERENCES

- [1] M. Krasner, *A class of hyperrings and hyperfields*, Intern. J. Math. and Math. Sci. **6** (2) (1983), 307–312. doi:10.1155/S0161171283000265

- [2] F. Marty, *Role de la notion d'hypergroupe dans l'étude des groupes non abéliens*, Comptes Rendus Acad. Sci. Paris Math. **201** (1935), 636–638.
- [3] F. Marty, *Sur les groupes et hypergroupes attachés à une fraction rationnelle*, Ann. Sci. de l'Ecole Norm. Sup. **53** (3) (1936), 82–123.
- [4] F. Marty, *Sur une généralisation de la notion de groupe.*, in: pages 45–49, “8^{ème} congrès des Mathématiciens Scandinaves”, Stockholm 1934.
- [5] A. Nakassis, *Expository and survey article recent results in hyperring and hyperfield theory*, J. Math. and Math. Sci. **11** (2) (1988), 209–220. doi:10.1155/S0161171288000250
- [6] P. Raja and S.M. Vaezpour, *Normed Hypervector spaces*, Iranian Journal of Mathematical Sciences and Informatics **2** (2007), 35–44.
- [7] M.K. Sen, *Utpal Dasgupta, Hypersemiring*, Bull. Cal. Math. Soc. **100** (2) (2008), 143–156.
- [8] M.S. Tallini, *A-ipermoduli e spazi ipervettoriali*, Rivista di Mat. Pura e Applicata, Università di Udine **3** (1988), 1–46.
- [9] M.S. Tallini, *Hypervector spaces*, Fourth Int. Congress on AHA, Xanthi, Greece (1990), 167–174.
- [10] M.S. Tallini, *Spazi ipervettoriali fortemente distributivi a sinistra*, Rend. Mat. Univ. Roma La Sapienza **11** (VII) (1991), 1–16.
- [11] M.S. Tallini, *Matroidal Hypervector spaces*, Journal of Geometry **42** (1991), 132–140. doi:10.1007/BF01231873
- [12] M.S. Tallini, *La categoria degli spazi ipervettoriali*, Quaderni Sem. Geom. Comb. Dip. Mat. “G. Castelnuovo” Univ. Di Roma La Sapienza, **110** (1993), 1–17.
- [13] M.S. Tallini, *Spazi ipervettoriali deboli e norme in tali spazi*. Quaderni Sem. Geom. Comb. Dip. Mat. “G. Castelnuovo”, Univ. Di Roma La Sapienza, **111** (1993), 1–14.
- [14] M.S. Tallini, *La categoria degli spazi ipervettoriali*, Rivista di Mat. Pura e Applicata, Univ. di Udine, **15** (1994), 97–109.
- [15] M.S. Tallini, *Weak hypervector spaces and norms in such spaces*, Proc. Int. Congress Algebraic Hyperstructures and Applications, Iasi, Romania, Luglio 1993, Hadronic Press, Palm Harbor, Florida (U.S.A.) (1994), 109–206.

- [16] M.S. Tallini, *Characterization of Remarkable Hypervector spaces*, Proc. of 8th int. Congress on Algebraic Hyperstructures and Applications, Samotraki, Greece, Sept. 1–9 2002, Spanidis Press, Xanthi, Greece, ISBN 960-87499-5-6 (2003), 231–237.

Received 23 March 2010
Revised 9 September 2010