

***bi-BL*-ALGEBRA**

MAHDEIEH ABBASLOO

Department of Mathematics, Science and Research Branch
Islamic Azad University, Kerman, Iran

e-mail: m.abbasloo@iauba.ac.ir

AND

ARSHAM BORUMAND SAEID

Department of Mathematics
Shahid Bahonar University of Kerman
Kerman, Iran

e-mail: arsham@mail.uk.ac.ir

Abstract

In this paper, we introduce the notion of a *bi-BL*-algebra, *bi*-filter, *bi*-deductive system and *bi*-Boolean elements of a *bi-BL*-algebra and deal with *bi*-filters in *bi-BL*-algebra. We study this structure and construct the quotient of *bi-BL*-algebra. Also present a classification for examples of proper *bi-BL*-algebras.

Keywords: *bi-BL*-algebra, *bi*-filter, *bi*-deductive system, *bi*-Boolean elements of a *bi-BL*-algebra.

2010 Mathematics Subject Classification: 03B47, 03G25, 06D99.

1. INTRODUCTION

*bi*structure is a tool as this answers a major problem faced by all algebraic structures - groups, semigroups, loops, groupoids etc. that is the union of two subgroups, or two subrings, or two subsemigroups etc. do not form any algebraic structure but all of them find a nice *bialgebraic* structure as *bigroups*, *birings*, *bisemigroups* etc. Except for this *bialgebraic* structure these would remain only as sets without any nice algebraic structure on them. Further when these *bialgebraic* structures are defined on them they enjoy not only the inherited qualities of the

algebraic structure from which they are taken but also several distinct algebraic properties that are not present in algebraic structures.

The study of *bialgebraic* structures started recently. The study of *bigroups* was carried out in 1994–1996. Further research on *bigroups* and fuzzy *bigroups* was published in 1998. In the year 1999, *bivector* spaces were introduced. In 2001, concept of free De Morgan *bisemigroups* and *bisemilattices* was studied. It is said by Zoltan Esik that these *bialgebraic* structures like *bigroups*, *bisemigroups*, *bilinear* rings help in the construction of finite machines or finite automaton and semi automaton. The notion of non-associative *bialgebraic* structures was first introduced in the year 2003, [19].

BL-algebra have been invented by P. Hajek [9] in order to provide an algebraic proof of the completeness theorem of "Basic Logic" (*BL*, for short) arising from the continuous triangular norms, familiar in the fuzzy Logic framework. The language of propositional Hajek basic logic [9] contains the binary connectives \odot and \rightarrow and the constant $\bar{0}$.

Axioms of *BL* are:

- (A₁) $(\phi \rightarrow \chi) \rightarrow ((\chi \rightarrow \psi) \rightarrow (\phi \rightarrow \psi))$
- (A₂) $(\phi \odot \chi) \rightarrow \phi$
- (A₃) $(\phi \odot \chi) \rightarrow (\chi \odot \phi)$
- (A₄) $(\phi \odot (\phi \rightarrow \chi)) \rightarrow (\chi \odot (\chi \rightarrow \phi))$
- (A_{5a}) $(\phi \rightarrow (\chi \rightarrow \psi)) \rightarrow ((\phi \odot \chi) \rightarrow \psi)$
- (A_{5b}) $((\phi \odot \chi) \rightarrow \psi) \rightarrow (\phi \rightarrow (\chi \rightarrow \psi))$
- (A₆) $((\phi \rightarrow \chi) \rightarrow \psi) \rightarrow (((\chi \rightarrow \phi) \rightarrow \psi) \rightarrow \psi)$
- (A₇) $\bar{0} \rightarrow \omega$.

In this paper, we generalize the notion of *BL*-algebra and introduce notion of *bi-BL*-algebra and study it. The notions of *bi-filter*, *bi-deductive* system and *bi-Boolean* elements of a *bi-BL*-algebra are introduced and studied this structure in detail. We construct the quotient of *bi-BL*-algebra, also present classes of examples of proper *bi-BL*-algebras.

2. PRELIMINARIES

2.1. Definitions and Theorems

Definition 2.1 [9]. A *BL*-algebra is an algebra $(A, \wedge, \vee, \odot, \rightarrow, 0, 1)$ with four binary operations $\wedge, \vee, \odot, \rightarrow$ and two constants $0, 1$ such that:

- (BL1) $(A, \wedge, \vee, \rightarrow, 0, 1)$ is a bounded lattice,
- (BL2) $(A, \odot, 1)$ is a commutative monoid,
- (BL3) \odot and \rightarrow form an adjoint pair i.e, $a \odot b \leq c$ if and only if $a \leq b \rightarrow c$,

- (BL4) $a \wedge b = a \odot (a \rightarrow b)$,
 (BL5) $(a \rightarrow b) \vee (b \rightarrow a) = 1$,
 for all $a, b, c \in A$.

A BL-algebra is called an MV-algebra if $x^{--} = x$, for all $x \in A$, where $x^- = x \rightarrow 0$.

Definition 2.2 [9]. A filter of a BL-algebra A is a nonempty subset F of A , such that for all $x, y \in A$, we have

- (1) $x, y \in F$ implies $x \odot y \in F$,
- (2) $x \in F$ and $x \leq y$ imply $y \in F$.

Definition 2.3 [17]. A non-empty subset D of BL-algebra A is called a deductive system if

- (1) $1 \in D$,
- (2) If $x \in D$ and $x \rightarrow y \in D$ imply $y \in D$.

Proposition 2.4 [17]. A non-empty subset F of BL-algebra is a deductive system if and only if F is a filter.

Theorem 2.5 [9]. Let F be a filter of a BL-algebra A . Define: $x \equiv_F y$ if and only if $x \rightarrow y \in F$ and $y \rightarrow x \in F$. Then \equiv_F is a congruence relation on A . The set of all congruence classes is denoted by $\frac{A}{F}$, i.e., $\frac{A}{F} := \{[x] | x \in A\}$, where $[x] = \{y \in A | x \equiv_F y\}$. Define $\bullet, \rightarrow, \sqcap, \sqcup$ on $\frac{A}{F}$ as follows:
 $[x] \bullet [y] = [x \odot y]$, $[x] \rightarrow [y] = [x \rightarrow y]$, $[x] \sqcap [y] = [x \wedge y]$, $[x] \sqcup [y] = [x \vee y]$.
 Therefore $(\frac{A}{F}, \sqcap, \sqcup, \bullet, \rightarrow, [1], [0])$ is a BL-algebra with respect to F .

Definition 2.6 [9]. Let L be a BL-algebra. An element $a \in L$ is called complemented if there is an $b \in L$ such that $a \vee b = 1$ and $a \wedge b = 0$; If such element b exists it is called a complement of a . We will denote the set of all complement in L by $B(L)$.

For any BL-algebra A , $B(A)$ denotes the Boolean algebra of all complement elements in $L(A)$ (hence $B(A) = B(L(A))$).

Definition 2.7 [7, 9, 18]. Let A and B are BL-algebras. A function $f : A \rightarrow B$ is called homomorphism of BL-algebras if and only if:

- (1) $f(0) = 0$,
- (2) $f(x * y) = f(x) * f(y)$,
- (3) $f(x \rightarrow y) = f(x) \rightarrow f(y)$,

for all $x, y \in A$.

3. *bi-BL*-ALGEBRA

3.1. Definition and some examples

Definition 3.1. A *bi-BL*-algebra is an algebra $(L, \wedge, \vee, \odot, \rightarrow, 0, 1)$ with four binary operations and two constants if $L = L_1 \cup L_2$ where L_1 and L_2 are proper subsets of L and

- (i) $(L_1, \wedge, \vee, \odot, \rightarrow, 0, 1)$ is a non-trivial *BL*-algebra,
- (ii) $(L_2, \wedge, \vee, \odot, \rightarrow, 0, 1)$ is a non-trivial *BL*-algebra.

Definition 3.2. If L is a *bi-BL*-algebra and also a *BL*-algebra, then we say that L is a super *BL*-algebra.

Definition 3.3. A *bi-BL*-algebra $L = L_1 \cup L_2$ is said to be finite if it has a finite number of elements and if L has infinite number of elements, then L is said to be infinite *bi-BL*-algebra.

Example 3.4. Let $L_1 = \{0, a, c, 1\}$ and $L_2 = \{0, b, c, 1\}$. Define \odot and \rightarrow as follow:

L_1	\odot	0	a	c	1
	0	0	0	0	0
	a	0	a	a	a
	c	0	a	c	c
	1	0	a	c	1

	\rightarrow	0	a	c	1
	0	1	1	1	1
	a	0	1	1	1
	c	0	a	1	1
	1	0	a	c	1

L_2	\odot	0	b	c	1
	0	0	0	0	0
	b	0	b	b	b
	c	0	b	c	c
	1	0	b	c	1

	\rightarrow	0	b	c	1
	0	1	1	1	1
	b	0	1	1	1
	c	0	b	1	1
	1	0	b	c	1

For L , whose tables are the following:

L	\odot	0	a	b	c	1
	0	0	0	0	0	0
	a	0	a	0	a	a
	b	0	0	b	b	b
	c	0	a	b	b	b
	1	0	a	b	c	1

	\rightarrow	0	a	b	c	1
	1	1	1	1	1	1
	a	b	1	b	1	1
	b	a	a	1	1	1
	c	0	a	b	1	1
	1	0	a	b	c	1

Then L_1 and L_2 are BL -algebras and $L = L_1 \cup L_2$ is a bi - BL -algebra but L is not a BL -algebra since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$. In this example $L_1 \cap L_2 \neq \{0, 1\}$.

Example 3.5. Let $L_1 = \{0, a, b, c, d, 1\}$ and $L_2 = \{0, d, e, 1\}$. Define \odot and \rightarrow as follow:

L_1	\odot	0	a	b	c	d	1
	0	0	0	0	0	0	0
	a	0	a	c	c	d	a
	b	0	c	b	c	d	b
	c	0	c	c	c	d	c
	d	0	d	d	d	0	d
	1	0	a	b	c	d	1
	\rightarrow	0	a	b	c	d	1
	0	1	1	1	1	1	1
	a	0	1	b	b	d	1
	b	0	a	1	a	d	1
	c	0	1	1	1	d	1
	d	d	1	1	1	1	1
	1	0	a	b	c	d	1

L_2	\odot	0	d	e	1
	0	0	0	0	0
	d	0	0	d	d
	e	0	d	e	e
	1	0	d	e	1
	\rightarrow	0	d	e	1
	0	1	1	1	1
	d	d	1	1	1
	e	0	d	1	1
	1	0	d	e	1

For L , whose tables are the following:

L	\odot	0	a	b	c	d	e	1
	0	0	0	0	0	0	0	0
	a	0	a	c	c	d	e	a
	b	0	c	b	c	d	b	b
	c	0	c	c	c	d	e	c
	d	0	d	d	d	0	d	d
	e	0	e	b	e	d	e	e
	1	0	a	b	c	d	e	1
	\rightarrow	0	a	b	c	d	e	1
	0	1	1	1	1	1	1	1
	a	0	1	b	b	d	e	1
	b	0	a	1	a	d	d	1
	c	0	1	1	1	d	e	1
	d	d	1	1	1	1	1	1
	e	0	d	b	d	d	1	1
	1	0	a	b	c	d	e	1

Then L_1 and L_2 are BL -algebras and $L = L_1 \cup L_2$ is a bi - BL -algebra but L is not a BL -algebra since $(a \rightarrow e) \vee (e \rightarrow a) = e \vee d = e \neq 1$. In this case, $L_1 \cap L_2 \neq \{0, 1\}$.

Example 3.6. Let $L_1 = \{0, a, c, 1\}$ and $L_2 = \{0, b, c, d, 1\}$. Define \odot and \rightarrow as follow:

L_1	\odot	0	a	c	1
	0	0	0	0	0
	a	0	a	a	a
	c	0	a	a	c
	1	0	a	c	1
	\rightarrow	0	a	c	1
	0	1	1	1	1
	a	0	1	1	1
	c	0	c	1	1
	1	0	a	c	1

L_2	\odot	<hr/>			
		0	b	d	1
	0	0	0	0	0
	b	0	0	b	b
	d	0	b	d	d
	1	0	b	d	1

\rightarrow	<hr/>			
	0	b	d	1
0	1	1	1	1
b	b	1	1	1
d	0	b	1	1
1	0	b	d	1

For L , whose tables are the following:

	\odot	0	a	b	c	d	1		\rightarrow	0	a	b	c	d	1
	0	0	0	0	0	0	0		0	1	1	1	1	1	1
	a	0	a	0	a	0	a		a	d	1	d	1	d	1
L	b	0	0	0	0	b	b		b	c	c	1	1	1	1
	c	0	a	0	a	b	c		c	b	c	d	1	d	1
	d	0	0	b	b	d	d		d	a	a	c	c	1	1
	1	0	a	b	c	d	1		1	0	a	b	c	d	1

Then L_1 and L_2 are BL -algebras. $L = L_1 \cup L_2$ is a bi - BL -algebra also L is a super BL -algebra. In this case, $L_1 \cap L_2 \neq \{0, 1\}$.

Remark 3.7. Special case of bi - BL -algebra:

A non-empty set $(L, \wedge, \vee, \odot, \rightarrow, 0, 1)$ is called a bi - BL -algebra if $L = L_1 \cup L_2$ where L_1 and L_2 are proper subsets of L (denote the least element by 0 and the greatest element by 1) and

- (i) $(L_1, \wedge, \vee, \odot, \rightarrow, 0_1, 1_1)$ is non-trivial a BL -algebra,
- (ii) $(L_2, \wedge, \vee, \odot, \rightarrow, 0_2, 1_2)$ is a non-trivial BL -algebra.

Now, we present classes of examples of proper bi - BL -algebras which is similar to BL -algebras [11]:

3.2. Classes of examples of bi - BL -algebras

We start details with the linearly ordered set(chain).

$$L_{n+1} = \{0, 1, 2, \dots, n\},$$

($n \geq 1$), organized as a lattice with $\wedge = \min$ and $\vee = \max$, and organized term equivalent:

$$\mathcal{L}_{n+1} = (L_{n+1}, \odot, ^-, n),$$

with:

$$x \odot y = \max(0, x + y - n), x^- = x \rightarrow 0, \quad (0 = n^-),$$

hence $x \rightarrow y = \max\{z | x \odot z \leq y\} = (x \odot y^-)^- = \min(n, y - x + n)$. Hence, for $n = 1, \dots, 6$, we have the linearly ordered MV -algebras $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}_4, \mathcal{L}_5, \mathcal{L}_6$, [11].

3.2.1. Classes of examples of finite, linearly ordered *bi-BL*-algebras

The examples are one of the following forms:

1. Linearly ordered $MV \cup$ linearly ordered MV ,
2. Linearly ordered $MV \cup$ linearly ordered BL or linearly ordered $BL \cup$ linearly ordered MV ,
3. Linearly ordered $BL \cup$ linearly ordered BL .

• (1) **Examples of the form: Linearly ordered $MV \cup$ linearly ordered MV .**

Denote $\mathcal{H}_{m+1,n+1} = \mathcal{L}_{m+1} \cup \mathcal{L}_{n+1}$, for $m, n \geq 1$.

1. Example of the form: $\mathcal{H}_{2,n+1} = \mathcal{L}_2 \cup \mathcal{L}_{n+1}$ for $n \geq 1$.

Denote $H_{2,n+1} = L_2 \cup L_{n+1} = \{-1, 0\} \cup \{0, 1, 2, \dots, n\} = \{-1, 0, 1, 2, \dots, n\}$.

For $n = 1, 2, 3, 4, 5$, since elements are from integer numbers then we have the linearly ordered *bi-BL*-algebras $\mathcal{H}_{2,2} = \mathcal{L}_2 \cup \mathcal{L}_2$, $\mathcal{H}_{2,3} = \mathcal{L}_2 \cup \mathcal{L}_3$, $\mathcal{H}_{2,4} = \mathcal{L}_2 \cup \mathcal{L}_4$, $\mathcal{H}_{2,5} = \mathcal{L}_2 \cup \mathcal{L}_5$, $\mathcal{H}_{2,6} = \mathcal{L}_2 \cup \mathcal{L}_6$, whose tables are the following:

$\mathcal{H}_{2,2}$	\odot	-1	0	1	\rightarrow	-1	0	1
	-1	-1	-1	-1	-1	1	1	1
	0	-1	0	0	0	-1	1	1
	1	-1	0	1	1	-1	0	1

$\mathcal{H}_{2,3}$	\odot	-1	0	1	2	\rightarrow	-1	0	1	2
	-1	-1	-1	-1	-1	-1	2	2	2	2
	0	-1	0	0	0	0	-1	2	2	2
	1	-1	0	0	1	1	-1	1	2	2
	2	-1	0	1	2	2	-1	0	1	2

$\mathcal{H}_{2,4}$	\odot	-1	0	1	2	3	\rightarrow	-1	0	1	2	3
	-1	-1	-1	-1	-1	-1	-1	3	3	3	3	3
	0	-1	0	0	0	0	0	-1	3	3	3	3
	1	-1	0	0	0	1	1	-1	2	3	3	3
	2	-1	0	0	1	2	2	-1	1	2	3	3
	3	-1	0	1	2	3	3	-1	0	1	2	3

	\odot	-1	0	1	2	3	4		\rightarrow	-1	0	1	2	3	4
$\mathcal{H}_{2,5}$	-1	-1	-1	-1	-1	-1	-1	-1	-1	4	4	4	4	4	4
	0	-1	0	0	0	0	0	0	0	-1	4	4	4	4	4
	1	-1	0	0	0	0	1	1	1	-1	3	4	4	4	4
	2	-1	0	0	0	1	2	2	2	-1	2	3	4	4	4
	3	-1	0	0	1	2	3	3	3	-1	1	2	3	4	4
	4	-1	0	1	2	3	4	4	4	-1	0	1	2	3	4

	\odot	-1	0	1	2	3	4	5		\rightarrow	-1	0	1	2	3	4	5
$\mathcal{H}_{2,6}$	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	5	5	5	5	5	5	5
	0	-1	0	0	0	0	0	0	0	0	-1	5	5	5	5	5	5
	1	-1	0	0	0	0	0	1	1	1	-1	4	5	5	5	5	5
	2	-1	0	0	0	0	1	2	2	2	-1	3	4	5	5	5	5
	3	-1	0	0	0	1	2	3	3	3	-1	2	3	4	5	5	5
	4	-1	0	0	1	2	3	4	4	4	-1	1	2	3	4	5	5
	5	-1	0	1	2	3	4	5	5	5	-1	0	1	2	3	4	5

2. Example of the form: $\mathcal{H}_{3,n+1} = \mathcal{L}_3 \cup \mathcal{L}_{n+1}$ for $n \geq 1$.

Denote $H_{3,n+1} = L_3 \cup L_{n+1} = \{-2, -1, 0\} \cup \{0, 1, \dots, n\} = \{-2, -1, 0, 1, 2, \dots, n\}$. For $n = 1, 2$, since elements are from integer numbers then we have the linearly ordered bi-BL-algebras $\mathcal{H}_{3,2} = \mathcal{L}_3 \cup \mathcal{L}_2$, $\mathcal{H}_{3,3} = \mathcal{L}_3 \cup \mathcal{L}_3$, whose tables are:

	\odot	-2	-1	0	1		\rightarrow	-2	-1	0	1
$\mathcal{H}_{3,2}$	-2	-2	-2	-2	-2	-2	-2	1	1	1	1
	-1	-2	-2	-1	-1	-1	-1	-1	1	1	1
	0	-2	-1	0	0	0	0	-2	-1	1	1
	1	-2	-1	0	1	1	1	-2	-1	0	1

	\odot	-2	-1	0	1	2		\rightarrow	-2	-1	0	1	2
$\mathcal{H}_{3,3}$	-2	-2	-2	-2	-2	-2	-2	-2	2	2	2	2	2
	-1	-2	-2	-1	-1	-1	-1	-1	-1	2	2	2	2
	0	-2	-1	0	0	0	0	0	-2	-1	2	2	2
	1	-2	-1	0	0	1	1	1	-2	-1	1	2	2
	2	-2	-1	0	1	2	2	2	-2	-1	0	1	2

Remark 3.8. The examples of the forms $\mathcal{H}_{m+1,n+1}$, for $m, n \geq 1$ are BL-algebras thus are super BL-algebras.

- (2) Examples of the form: Linearly ordered $MV \cup$ linearly ordered BL or linearly ordered $BL \cup$ linearly ordered MV .

Denote $\mathcal{H}_{m+1,n+1,p+1} = \mathcal{L}_{m+1} \cup \mathcal{H}_{n+1,p+1} = \mathcal{L}_{m+1} \cup (\mathcal{L}_{n+1} \cup \mathcal{L}_{p+1}) = (\mathcal{L}_{m+1} \cup \mathcal{L}_{n+1}) \cup \mathcal{L}_{p+1} = \mathcal{H}_{m+1,n+1} \cup \mathcal{L}_{p+1}$, by associativity of \cup .

Example. The set $H_{2,2,2} = L_2 \cup H_{2,2} = \{-1, 0\} \cup \{0, 1, 2\} = H_{2,2} \cup L_2 = \{-1, 0, 1\} \cup \{1, 2\} = \{-1, 0, 1, 2\}$, organized as a lattice in a obvious way and as bi - BL -algebra $\mathcal{H}_{2,2,2} = \mathcal{H}_{2,2} \cup \mathcal{L}_2$ with the following tables:

$\mathcal{H}_{2,2,2}$	\odot	-1	0	1	2	\rightarrow	-1	0	1	2
	-1	-1	-1	-1	-1	-1	2	2	2	2
	0	-1	0	0	0	0	-1	2	2	2
	1	-1	0	1	1	1	-1	0	2	2
	2	-1	0	1	2	2	-1	0	1	2

Remark 3.9. The examples of the forms $\mathcal{H}_{m+1,n+1,p+1}$, for $m, n, p \geq 1$ are BL -algebras thus become a super BL -algebras.

- (3) Examples of the form: Linearly ordered $BL \cup$ linearly ordered BL or equivalent forms.

Denote $\mathcal{H}_{m+1,n+1,p+1,q+1} = \mathcal{H}_{m+1,n+1} \cup \mathcal{H}_{p+1,q+1} = (\mathcal{L}_{m+1} \cup \mathcal{L}_{n+1}) \cup (\mathcal{L}_{p+1} \cup \mathcal{L}_{q+1}) = \mathcal{H}_{m+1,n+1,p+1} \cup \mathcal{L}_{q+1} = \mathcal{L}_{m+1} \cup \mathcal{H}_{n+1,p+1,q+1}$, by associativity of \cup .

Example. The set $H_{2,2,2,2} = H_{2,2} \cup H_{2,2} = H_{2,2,2} \cup L_2 = \{-1, 0, 1, 2\} \cup \{2, 3\} = \{-1, 0, 1, 2, 3\} = L_2 \cup H_{2,2,2} = \{-1, 0\} \cup \{0, 1, 2, 3\}$, organized as a lattice in a obvious way and as bi - BL -algebra $\mathcal{H}_{2,2,2,2} = \mathcal{H}_{2,2,2} \cup \mathcal{L}_2$ with the following tables:

$\mathcal{H}_{2,2,2,2}$	\odot	-1	0	1	2	3	\rightarrow	-1	0	1	1	1
	-1	-1	-1	-1	-1	-1	-1	3	3	3	3	3
	0	-1	0	0	0	0	0	-1	3	3	3	3
	1	-1	0	1	1	1	1	-1	0	3	3	3
	2	-1	0	1	2	2	2	-1	0	1	3	3
	3	-1	0	1	2	3	3	-1	0	1	2	3

Remark 3.10. The examples of the forms $\mathcal{H}_{m+1,n+1,p+1,q+1}$, for $m, n, p, q \geq 1$ are BL -algebras thus become a super BL -algebras.

3.3. Classes of examples of finite, non-linearly ordered bi-BL-algebras

The examples are one of the following forms:

1. Linearly ordered $MV \cup$ non-linearly ordered MV ,
2. Linearly ordered $MV \cup$ non-linearly ordered BL or linearly ordered $BL \cup$ non-linearly ordered MV ,
3. Linearly ordered $BL \cup$ non-linearly ordered BL .

• (1) **Examples of the form: Linearly ordered $MV \cup$ non-linearly ordered MV .**

Denote $\mathcal{H}_{p+1,(n+1) \times (m+1)} = \mathcal{L}_{p+1} \cup \mathcal{L}_{(n+1) \times (m+1)}$, for $p, m, n \geq 1$.

We present two families of examples.

1. Examples of the form: $\mathcal{H}_{2,(n+1) \times (m+1)} = \mathcal{L}_2 \cup \mathcal{L}_{(n+1) \times (m+1)}$ for $n, m \geq 1$.

Denote $H_{2,(n+1) \times (m+1)} = L_2 \cup L_{(n+1) \times (m+1)}$, with $n, m \geq 1$.

We present four examples.

Example 1. The set $H_{2,2 \times 2} = L_2 \cup L_{2 \times 2} = \{-1, 0\} \cup \{0, a, b, 1\} = \{-1, 0, a, b, 1\}$, organized as a lattice as and with operations \rightarrow and \odot in the following tables, is a non-linearly ordered bi-BL-algebra, denoted by $\mathcal{H}_{2,2 \times 2} = \mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}$.

	\odot	-1	0	a	b	1		\rightarrow	-1	0	a	b	1
	-1	-1	-1	-1	-1	-1		-1	1	1	1	1	1
	0	-1	0	0	0	0		0	-1	1	1	1	1
	a	-1	0	a	0	a		a	-1	b	1	b	1
	b	-1	0	0	b	b		b	-1	a	a	1	1
	1	-1	0	a	b	1		1	-1	0	a	b	1

Example 2. The set $H_{2,3 \times 2} = L_2 \cup L_{3 \times 2} = \{-1, 0\} \cup \{0, a, b, c, d, 1\} = \{-1, 0, a, b, c, d, 1\}$, organized as a lattice as and with operations \rightarrow and \odot in the following tables, is a non-linearly ordered BL -algebra, denoted by $\mathcal{H}_{2,3 \times 2} = \mathcal{L}_2 \cup \mathcal{L}_{3 \times 2}$.

$\mathcal{H}_{2,3 \times 2}$	\odot	-1	0	a	b	c	d	1	\rightarrow	-1	0	a	b	c	d	1
	-1	-1	-1	-1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1
	0	-1	0	0	0	0	0	0	0	-1	1	1	1	1	1	1
	a	-1	0	a	0	a	0	a	a	-1	d	1	d	1	d	1
	b	-1	0	0	0	0	b	b	b	-1	c	c	1	1	1	1
	c	-1	0	a	0	a	b	b	c	-1	b	c	d	1	d	1
	d	-1	0	0	b	b	d	d	d	-1	a	a	c	c	1	1
	1	-1	0	a	b	c	d	1	1	-1	0	a	b	c	d	1

Example 3. The set $H_{2,3 \times 3} = L_2 \cup L_{3 \times 3} = \{-1, 0\} \cup \{0, a, b, c, d, e, f, g, 1\} = \{-1, 0, a, b, c, d, e, f, g, 1\}$, organized as a lattice as and with operations \rightarrow and \odot in the following tables, is a non-linearly ordered BL -algebra, denoted by $\mathcal{H}_{2,3 \times 3} = \mathcal{L}_2 \cup \mathcal{L}_{3 \times 3}$.

$\mathcal{H}_{2,3 \times 3}$	\odot	-1	0	a	b	c	d	e	f	g	1
	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
	0	-1	0	0	0	0	0	0	0	0	0
	a	-1	0	0	a	0	0	a	0	0	a
	b	-1	0	a	b	0	a	b	0	a	b
	c	-1	0	0	0	0	0	0	c	c	c
	d	-1	0	0	a	0	0	a	c	c	d
	e	-1	0	a	b	0	a	b	c	d	e
	f	-1	0	0	0	c	c	c	f	f	f
	g	-1	0	0	a	c	c	d	f	f	g
	1	-1	0	a	b	c	d	e	f	g	1
	\rightarrow	-1	0	a	b	c	d	e	f	g	1
	-1	1	1	1	1	1	1	1	1	1	1
	0	-1	1	1	1	1	1	1	1	1	1
	a	-1	g	1	1	g	1	g	1	1	1
	b	-1	f	g	1	f	g	1	f	g	1
	c	-1	e	e	e	1	1	1	1	1	1
	d	-1	d	e	e	g	1	g	1	1	1
	e	-1	c	d	e	f	g	1	f	g	1
	f	-1	b	b	b	e	e	e	1	1	1
	g	-1	a	b	b	d	e	e	g	1	1
	1	-1	0	a	b	c	d	e	f	g	1

Example 4. The set $H_{2,4 \times 2} = L_2 \cup L_{4 \times 2} = \{-1, 0\} \cup \{0, a, b, c, d, e, f, 1\} = \{-1, 0, a, b, c, d, e, f, 1\}$ is a bi - BL -algebra.

2. Examples of the form: $\mathcal{H}_{3,(n+1) \times (m+1)} = \mathcal{L}_3 \cup \mathcal{L}_{(n+1) \times (m+1)}$ for $n, m \geq 1$.

We present here only one example.

The set $H_{3,2 \times 2} = L_3 \cup L_{2 \times 2} = \{-2, -1, 0\} \cup \{0, a, b, 1\} = \{-2, -1, 0, a, b, 1\}$, organized as a lattice as and with operations \rightarrow and \odot in the following tables, is a non-linearly ordered bi-BL-algebra, denoted by $\mathcal{H}_{3,2 \times 2} = \mathcal{L}_3 \cup \mathcal{L}_{2 \times 2}$.

	\odot	-2	-1	0	a	b	1	\rightarrow	-2	-1	0	a	b	1
$\mathcal{H}_{3,2 \times 2}$	-2	-2	-2	-2	-2	-2	-2	-2	1	1	1	1	1	1
	-1	-2	-1	-1	-1	-1	-1	-1	-2	1	1	1	1	1
	0	-2	-1	0	0	0	0	0	-2	-1	1	1	1	1
	a	-2	-1	0	a	0	a	a	-2	-1	b	1	b	1
	b	-2	-1	0	0	b	b	b	-2	-1	a	a	1	1
	1	-2	-1	0	a	b	1	1	-2	-1	0	a	b	1

Remark 3.11. The examples of forms $\mathcal{H}_{p+1,(n+1) \times (m+1)}$, for $p, n, m \geq 1$ are BL-algebras thus are super BL-algebras.

• (2) **Examples of the form: Linearly ordered MV \cup non-linearly ordered BL or linearly ordered BL \cup non-linearly ordered MV.**

Denote for $u, v, n, m \geq 1$, the bi-BL-algebras: $\mathcal{H}_{u+1,v+1,(n+1) \times (m+1)} = \mathcal{L}_{u+1} \cup \mathcal{L}_{v+1} \cup \mathcal{L}_{(n+1) \times (m+1)} = \mathcal{L}_{u+1} \cup \mathcal{H}_{v+1,(n+1) \times (m+1)} = \mathcal{H}_{u+1,v+1} \cup \mathcal{L}_{(n+1) \times (m+1)}$, by the associativity of \cup .

We present two examples.

Example 1. Consider the bi-BL-algebra $\mathcal{H}_{2,2,2 \times 2} = \mathcal{L}_2 \cup \mathcal{H}_{2,2 \times 2} = \mathcal{H}_{2,2} \cup \mathcal{L}_{2 \times 2}$ the underline set, $\{-2, -1, 0, a, b, 1\}$ can be considered either as the union of sets:

$$H_{(2,2),2 \times 2} = [\{-2, 1\} \cup \{-1, 0\}] \cup \{0, a, b, 1\} = [L_2 \cup L_2] \cup L_{2 \times 2}$$

or as the union

$$H_{2,(2,2 \times 2)} = \{-2, -1\} \cup [\{-1, 0\} \cup \{0, a, b, 1\}] = L_2 \cup [L_2 \cup L_{2 \times 2}] = L_2 \cup H_{2,2 \times 2}.$$

It has the following tables:

	\odot	-2	-1	0	a	b	1	\rightarrow	-2	-1	0	a	b	1
$\mathcal{H}_{2,2,2 \times 2}$	-2	-2	-2	-2	-2	-2	-2	-2	1	1	1	1	1	1
	-1	-2	-1	-1	-1	-1	-1	-1	-2	1	1	1	1	1
	0	-2	-1	0	0	0	0	0	-2	-1	1	1	1	1
	a	-2	-1	0	a	0	a	a	-2	-1	b	1	b	1
	b	-2	-1	0	0	b	b	b	-2	-1	a	a	1	1
	1	-2	-1	0	a	b	1	1	-2	-1	0	a	b	1

Example 2. Consider the *bi-BL*-algebra $\mathcal{H}_{2,(2,2)\times 2} = \mathcal{L}_2 \cup \mathcal{H}_{(2,2)\times 2}$. The set $H_{2,(2,2)\times 2} = L_2 \cup H_{(2,2)\times 2} = \{-1, 0\} \cup \{0, a, b, c, d, 1\} = \{-1, 0, a, b, c, d, 1\}$, organized as a lattice and a bounded lattice with the operations \rightarrow and \odot form the following tables is a *bi-BL*-algebra, denoted by $\mathcal{H}_{2,(2,2)\times 2}$.

\odot	-1	0	a	b	c	d	1	\rightarrow	-1	0	a	b	c	d	1
-1	-1	-1	-1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1
0	-1	0	0	0	0	0	0	0	-1	1	1	1	1	1	1
a	-1	0	a	0	a	0	a	a	-1	d	1	d	1	d	1
b	-1	0	0	b	b	b	b	b	-1	a	a	1	1	1	1
c	-1	0	a	b	c	b	c	c	-1	0	a	d	1	d	1
d	-1	0	0	b	b	d	d	d	-1	a	a	c	c	1	1
1	-1	0	a	b	c	d	1	1	-1	0	a	b	c	d	1

Remark 3.12. The examples of forms $\mathcal{H}_{u+1,v+1,(n+1)\times(m+1)}$, for $u, v, n, m \geq 1$ are *BL*-algebras thus become a super *BL*-algebras.

• **(3) Examples of the form: Linearly ordered *BL* \cup non-linearly ordered *BL* or equivalent forms.**

Denote for $u, v, n, m, p \geq 1$, the *bi-BL*-algebras: $\mathcal{H}_{u+1,v+1,(n+1,m+1)\times(p+1)} = \mathcal{H}_{u+1,v+1} \cup \mathcal{L}_{(n+1,m+1)\times(p+1)}$.

Example. Consider the *bi-BL*-algebra $\mathcal{H}_{2,2,(2,2)\times 2} = \mathcal{H}_{2,2} \cup \mathcal{H}_{(2,2)\times 2} = (\mathcal{L}_2 \cup \mathcal{L}_2) \cup \mathcal{H}_{(2,2)\times 2} = \mathcal{L}_2 \cup \mathcal{H}_{2,(2,2)\times 2}$ with the underline set $H_{2,2,(2,2)\times 2} = H_{2,2} \cup H_{(2,2)\times 2} = \{-2, -1, 0\} \cup \{0, a, b, c, d, 1\} = \{-2, -1, 0, a, b, c, d, 1\}$, organized as a lattice, with the operations \rightarrow and \odot in the following tables:

\odot	-2	-1	0	a	b	c	d	1	\rightarrow	-2	-1	0	a	b	c	d	1
-2	-2	-2	-2	-2	-2	-2	-2	-2	-2	1	1	1	1	1	1	1	1
-1	-2	-1	-1	-1	-1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1
0	-2	-1	0	0	0	0	0	0	0	-2	-1	1	1	1	1	1	1
a	-2	-1	0	a	0	a	0	a	a	-2	-1	d	1	d	1	d	1
b	-2	-1	0	0	b	b	b	b	b	-2	-1	a	a	1	1	1	1
c	-2	-1	0	a	b	c	b	c	c	-2	-1	0	a	d	1	d	1
d	-2	-1	0	0	b	b	d	d	d	-2	-1	a	a	c	c	1	1
1	-2	-1	0	a	b	c	c	1	1	-2	-1	0	a	b	c	d	1

Remark 3.13. The examples of forms $\mathcal{H}_{u+1,v+1,(n+1,m+1)\times(p+1)}$, for $u, v, n, m, p \geq 1$ are *BL*-algebras thus become a super *BL*-algebras.

3.3.1. Example of infinite bi-BL-algebras

By [11] we present example of infinite, linearly ordered bi-BL-algebra.

Example. The linearly ordered set(chain) $H_{P(\mathbb{Z}),2} = P(\mathbb{Z}) \cup L_2 = (\mathbb{Z}^- \cup -\infty) \cup L_2 = \{-\infty, \dots, -3, -2, -1, 0\} \cup \{0, 1\} = \{-\infty, \dots, -3, -2, -1, 0, 1\}$ with the operations \rightarrow and \odot defined by the following tables, is a linearly ordered bi-BL-algebra, denoted by $\mathcal{H}_{P(\mathbb{Z}),2} = \mathcal{P}(\mathbb{Z}) \cup \mathcal{L}_2$.

	\odot	$-\infty$	\dots	-3	-2	-1	0	1
	$-\infty$	$-\infty$	$-\infty$	$-\infty$	$-\infty$	$-\infty$	$-\infty$	$-\infty$
	\vdots	\dots	\dots	\vdots	\vdots	\vdots	\vdots	\vdots
$\mathcal{H}_{P(\mathbb{Z}),2}$	-3	$-\infty$	\dots	-6	-5	-4	-3	-3
	-2	$-\infty$	\dots	-5	-4	-3	-2	-2
	-1	$-\infty$	\dots	-4	-3	-2	-1	-1
	0	$-\infty$	\dots	-3	-2	-1	0	0
	1	$-\infty$	\dots	-3	-2	-1	0	1

	\rightarrow	$-\infty$	\dots	-3	-2	-1	0	1
	$-\infty$	1	\dots	1	1	1	1	1
	\vdots	\vdots	\dots	\vdots	\vdots	\vdots	\vdots	\vdots
	-3	$-\infty$	\dots	1	1	1	1	1
	-2	$-\infty$	\dots	-1	1	1	1	1
	-1	$-\infty$	\dots	-2	-1	1	1	1
	0	$-\infty$	\dots	-3	-2	-1	1	1
	1	$-\infty$	\dots	-3	-2	-1	0	1

3.3.2. Classes of finite bi-BL-algebras such that are not super BL-algebras

The examples will be of the form: non-linearly ordered MV/BL-algebra \cup MV/BL-algebra, more precisely of one of the following forms:

- (1) non-linearly ordered MV \cup linearly ordered MV,
- (2) non-linearly ordered MV \cup non-linearly ordered MV,
- (3) non-linearly ordered MV \cup linearly ordered BL,
- (4) non-linearly ordered MV \cup non-linearly ordered BL,
- (5) non-linearly ordered BL \cup linearly ordered MV,
- (6) non-linearly ordered BL \cup non-linearly ordered MV,

(7) non-linearly ordered $BL \cup$ linearly ordered BL ,

(8) non-linearly ordered $BL \cup$ non-linearly ordered BL .

- (1) **Examples of the form: non-linearly ordered $MV \cup$ linearly ordered MV .**

Denote, for $p, q, n \geq 1$

$$\mathcal{D}_{(p+1) \times (q+1), n+1} = \mathcal{L}_{(p+1) \times (q+1)} \cup \mathcal{L}_{n+1}.$$

We present three examples of above form.

Example 1. The *bi-BL*-algebra

$$\mathcal{D}_{2 \times 2, 2} = \mathcal{L}_{2 \times 2} \cup \mathcal{L}_2,$$

with the underline set

$$D_{2 \times 2, 2} = L_{2 \times 2} \cup L_2 = \{0, a, b, c\} \cup \{c, 1\} = \{0, a, b, c, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	a	b	c	1		\rightarrow	0	a	b	c	1
	0	0	0	0	0	0		0	1	1	1	1	1
	a	0	a	0	a	a		a	b	1	b	1	1
$\mathcal{D}_{2 \times 2, 2}$	b	0	0	b	b	b		b	a	a	1	1	1
	c	0	a	b	c	c		c	0	a	b	1	1
	1	0	a	b	c	1		1	0	a	b	c	1

note that $\mathcal{D}_{2 \times 2, 2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$ thus $\mathcal{D}_{2 \times 2, 2}$ is not a super BL -algebra.

Example 2. The bi - BL -algebra

$$\mathcal{D}_{2 \times 2, 3} = \mathcal{L}_{2 \times 2} \cup \mathcal{L}_3,$$

with the underline set

$$\mathcal{D}_{2 \times 2, 3} = L_{2 \times 2} \cup L_3 = \{0, a, b, c\} \cup \{c, d, 1\} = \{0, a, b, c, d, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	a	b	c	d	1		\rightarrow	0	a	b	c	d	1
	0	0	0	0	0	0	0		0	1	1	1	1	1	1
	a	0	a	0	a	a	a		a	b	1	b	1	1	1
$\mathcal{D}_{2 \times 2, 3}$	b	0	0	b	b	b	b		b	a	a	1	1	1	1
	c	0	a	b	c	c	c		c	0	a	b	1	1	1
	d	0	a	b	c	c	d		d	0	a	b	d	1	1
	1	0	a	b	c	d	1		1	0	a	b	c	d	1

note that $\mathcal{D}_{2 \times 2, 3}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$ thus $\mathcal{D}_{2 \times 2, 3}$ is not a super BL -algebra.

Example 3. The bi - BL -algebra

$$\mathcal{D}_{2 \times 3, 2} = \mathcal{L}_{2 \times 3} \cup \mathcal{L}_2,$$

with the underline set

$$\mathcal{D}_{2 \times 3, 2} = L_{2 \times 3} \cup L_2 = \{0, a, b, c, d, n\} \cup \{n, 1\} = \{0, a, b, c, d, n, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	a	b	c	d	n	1		\rightarrow	0	a	b	c	d	n	1
	0	0	0	0	0	0	0	0		0	1	1	1	1	1	1	1
	a	0	0	a	0	0	a	a		a	d	1	1	d	1	1	1
$\mathcal{D}_{2 \times 3, 2}$	b	0	a	b	0	a	b	b		b	c	d	1	c	d	1	1
	c	0	0	0	c	c	c	c		c	b	b	b	1	1	1	1
	d	0	0	a	c	c	d	d		d	a	b	b	d	1	1	1
	n	0	a	b	c	d	n	n		n	0	a	b	c	d	1	1
	1	0	a	b	c	d	n	1		1	0	a	b	c	d	n	1

note that $\mathcal{D}_{2 \times 3, 2}$ is not a BL -algebra, since $(b \rightarrow d) \vee (d \rightarrow b) = d \vee b = n \neq 1$ thus $\mathcal{D}_{2 \times 3, 2}$ is not a super BL -algebra.

• (2) Examples of the form: non-linearly ordered $MV \cup$ non-linearly ordered MV .

For $n, m, u, v \geq 1$, denote,

$$\mathcal{D}_{(n+1) \times (m+1), (u+1) \times (v+1)} = \mathcal{L}_{(n+1) \times (m+1)} \cup \mathcal{L}_{(u+1) \times (v+1)}.$$

Example. The bi - BL -algebra

$$\mathcal{D}_{2 \times 2, 2 \times 2} = \mathcal{L}_{2 \times 2} \cup \mathcal{L}_{2 \times 2},$$

with the underline set

$$D_{2 \times 2, 2 \times 2} = L_{2 \times 2} \cup L_{2 \times 2} = \{0, a, b, n\} \cup \{n, c, d, 1\} = \{0, a, b, n, c, d, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	a	b	n	c	d	1		\rightarrow	0	a	b	n	c	d	1
	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1
	a	0	a	0	a	a	a	a	a	a	b	1	b	1	1	1	1
	b	0	0	b	b	b	b	b	b	b	a	a	1	1	1	1	1
$\mathcal{D}_{2 \times 2, 2 \times 2}$	n	0	a	b	n	n	n	n	n	n	0	a	b	1	1	1	1
	c	0	a	b	n	c	n	c	c	c	0	a	b	d	1	d	1
	d	0	a	b	n	n	d	d	d	d	0	a	b	c	c	1	1
	1	0	a	b	n	c	d	1	1	1	0	a	b	n	c	d	1

note that $D_{2 \times 2, 2 \times 2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = n \neq 1$, thus $D_{2 \times 2, 2 \times 2}$ is not a super BL -algebra.

• (3) Examples of the form: non-linearly ordered $MV \cup$ linearly ordered BL or equivalent forms.

Denote, for $p, q, n, m \geq 1$,

$$\mathcal{D}_{(p+1) \times (q+1), n+1, m+1} = \mathcal{L}_{(p+1) \times (q+1)} \cup \mathcal{H}_{n+1, m+1}.$$

Example. The bi - BL -algebra

$$\mathcal{D}_{2 \times 2, 2, 2} = \mathcal{L}_{2 \times 2} \cup \mathcal{H}_{2, 2} = \mathcal{L}_{2 \times 2} \cup (\mathcal{L}_2 \cup \mathcal{L}_2) = \mathcal{D}_{2 \times 2, 2} \cup \mathcal{L}_2,$$

with the underline set

$$D_{2 \times 2, 2, 2} = L_{2 \times 2} \cup H_{2, 2} = \{0, a, b, c\} \cup \{c, d, 1\} = \{0, a, b, c, d, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	a	b	c	d	1		\rightarrow	0	a	b	c	d	1
	0	0	0	0	0	0	0		0	1	1	1	1	1	1
	a	0	a	0	a	a	a		a	b	1	b	1	1	1
$\mathcal{D}_{2 \times 2, 2, 2}$	b	0	0	b	b	b	b		b	a	a	1	1	1	1
	c	0	a	b	c	c	c		c	0	a	b	1	1	1
	d	0	a	b	c	d	d		d	0	a	b	c	1	1
	1	0	a	b	c	d	1		1	0	a	b	c	d	1

note that $\mathcal{D}_{2 \times 2, 2, 2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$, thus $\mathcal{D}_{2 \times 2, 2, 2}$ is not a super BL -algebra.

- (4) Examples of the form: non-linearly ordered $MV \cup$ non-linearly ordered BL or equivalent forms.

Denote, for $m, n, p, u, v \geq 1$,

$$\mathcal{D}_{(m+1) \times (n+1), p+1, (u+1) \times (v+1)} = \mathcal{L}_{(m+1) \times (n+1)} \cup \mathcal{H}_{p+1, (u+1) \times (v+1)}.$$

Example. The *bi-BL*-algebra:

$$\begin{aligned}\mathcal{D}_{2 \times 2, 2, 2 \times 2} &= \mathcal{L}_{2 \times 2} \cup H_{2, 2 \times 2} = L_{2 \times 2} \cup (L_2 \cup \mathcal{L}_{2 \times 2}) = (\mathcal{L}_{2 \times 2} \cup \mathcal{L}_2) \\ &= \mathcal{D}_{2 \times 2, 2} \cup \mathcal{L}_{2 \times 2},\end{aligned}$$

with the underline set

$$\begin{aligned}D_{2 \times 2, 2, 2 \times 2} &= L_{2 \times 2} \cup H_{2, 2 \times 2} = \{0, a, b, p\} \cup \{p, n\} \cup \{n, c, d, 1\} \\ &= \{0, a, b, p, n, c, d, 1\},\end{aligned}$$

is organized as a lattice as with the following tables:

\odot	0	a	b	p	n	c	d	1	\rightarrow	0	a	b	p	n	c	d	1
0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1
a	0	a	0	a	a	a	a	a	a	b	1	b	1	1	1	1	1
b	0	0	b	b	b	b	b	b	b	a	a	1	1	1	1	1	1
$\mathcal{D}_{2 \times 2, 2, 2 \times 2}$ p	0	a	b	p	p	p	p	p	p	0	a	b	1	1	1	1	1
n	0	a	b	p	n	n	n	n	n	0	a	b	p	1	1	1	1
c	0	a	b	p	n	c	n	c	c	0	a	b	p	d	1	d	1
d	0	a	b	p	n	n	d	d	d	0	a	b	p	c	c	1	1
1	0	a	b	p	n	c	d	1	1	0	a	b	p	n	c	d	1

note that $D_{2 \times 2, 2, 2 \times 2}$ is not a *BL*-algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = p \neq 1$ thus $\mathcal{D}_{2 \times 2, 2, 2 \times 2}$ is not a super *BL*-algebra.

• (5) **Examples of the form: non-linearly ordered *BL* \cup linearly ordered *MV* or equivalent forms.**

We consider here only two examples among all possible examples.

Example 1. The *bi-BL*-algebra

$$\mathcal{D}_{2, 2 \times 2, 2} = \mathcal{H}_{2, 2 \times 2} \cup \mathcal{L}_2 = (\mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}) \cup \mathcal{L}_2 = \mathcal{L}_2 \cup (\mathcal{L}_{2 \times 2} \cup \mathcal{L}_2) = \mathcal{L}_2 \cup \mathcal{D}_{2 \times 2, 2},$$

with the underline set

$$D_{2, 2 \times 2, 2} = H_{2, 2 \times 2} \cup L_2 = \{0, n, a, b, m\} \cup \{m, 1\} = \{0, n, a, b, m, 1\},$$

is organized as a lattice with the following tables:

	\odot	0	n	a	b	m	1		\rightarrow	0	n	a	b	m	1
	0	0	0	0	0	0	0		0	1	1	1	1	1	1
	n	0	n	n	n	n	n		n	0	1	1	1	1	1
$\mathcal{D}_{2,2 \times 2,2}$	a	0	n	a	n	a	a		a	0	b	1	b	1	1
	b	0	n	n	b	b	b		b	0	a	a	1	1	1
	m	0	n	a	b	m	m		m	0	n	a	b	1	1
	1	0	n	a	b	m	1		1	0	n	a	b	m	1

note that $\mathcal{D}_{2,2 \times 2,2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = m \neq 1$ thus $\mathcal{D}_{2,2 \times 2,2}$ is not a super BL -algebra.

Example 2. The bi - BL -algebra

$$\mathcal{D}_{3,2 \times 2,2} = \mathcal{H}_{3,2 \times 2} \cup \mathcal{L}_2 = (\mathcal{L}_3 \cup \mathcal{L}_{2 \times 2}) \cup \mathcal{L}_2 = \mathcal{L}_3 \cup (\mathcal{L}_{2 \times 2} \cup \mathcal{L}_2) = \mathcal{L}_3 \cup \mathcal{D}_{2 \times 2,2},$$

with the underline set

$$D_{3,2 \times 2,2} = H_{3,2 \times 2} \cup L_2 = \{-2, -1, 0, a, b, c\} \cup \{c, 1\} = \{-2, -1, 0, a, b, c, 1\}$$

is organized as a lattice with the following tables:

	\odot	-2	-1	0	a	b	c	1		\rightarrow	-2	-1	0	a	b	c	1
	-2	-2	-2	-2	-2	-2	-2	-2		-2	1	1	1	1	1	1	1
	-1	-2	-2	-1	-1	-1	-1	-1		-1	-1	1	1	1	1	1	1
$\mathcal{D}_{3,2 \times 2,2}$	0	-2	-1	0	0	0	0	0		0	-2	-1	1	1	1	1	1
	a	-2	-1	0	a	0	a	a		a	-2	-1	b	1	b	1	1
	b	-2	-1	0	0	b	b	b		b	-2	-1	a	a	1	1	1
	c	-2	-1	0	a	b	c	c		c	-2	-1	0	a	b	1	1
	1	-2	-1	0	a	b	c	1		1	-2	-1	0	a	b	c	1

note that $\mathcal{D}_{3,2 \times 2,2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$ thus $\mathcal{D}_{3,2 \times 2,2}$ is not a super BL -algebra.

• (6) **Examples of the form: non-linearly ordered $BL \cup$ non-linearly ordered MV or equivalent forms.**

Example. The bi - BL -algebra

$$\begin{aligned} \mathcal{D}_{2,2 \times 2,2 \times 2} &= \mathcal{H}_{2,2 \times 2} \cup \mathcal{L}_{2 \times 2} = (\mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}) \cup \mathcal{L}_{2 \times 2} \\ &= \mathcal{L}_2 \cup (\mathcal{L}_{2 \times 2} \cup \mathcal{L}_{2 \times 2}) = \mathcal{L}_2 \cup \mathcal{D}_{2 \times 2,2 \times 2} \end{aligned}$$

with the support set

$$\begin{aligned} D_{2,2 \times 2,2 \times 2} &= H_{2,2 \times 2} \cup L_{2 \times 2} = \{-1, 0, a, b, n\} \cup \{n, c, d, 1\} \\ &= \{-1, 0, a, b, n, c, d, 1\}, \end{aligned}$$

is organized as a lattice with the following tables:

\odot	-1	0	a	b	n	c	d	1	\rightarrow	-1	0	a	b	n	c	d	1
-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1	1
0	-1	0	0	0	0	0	0	0	0	-1	1	1	1	1	1	1	1
a	-1	0	a	0	a	a	a	a	a	-1	b	1	b	1	1	1	1
b	-1	0	0	b	b	b	b	b	b	-1	a	a	1	1	1	1	1
n	-1	0	a	b	n	n	n	n	n	-1	0	a	b	1	1	1	1
c	-1	0	a	b	n	c	n	c	c	-1	0	a	b	d	1	d	1
d	-1	0	a	b	n	n	d	d	d	-1	0	a	b	c	c	1	1
1	-1	0	a	b	n	c	d	1	1	-1	0	a	b	n	c	d	1

note that $\mathcal{D}_{2,2 \times 2,2 \times 2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = n \neq 1$ thus $\mathcal{D}_{2,2 \times 2,2 \times 2}$ is not super BL -algebra.

• (7) **Examples of the form: non-linearly ordered $BL \cup$ linearly ordered BL or equivalent forms.**

Example. The bi - BL -algebra

$$\mathcal{D}_{2,2 \times 2,2,2} = \mathcal{H}_{2,2 \times 2} \cup \mathcal{H}_{2,2} = (\mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}) \cup (\mathcal{L}_2 \cup \mathcal{L}_2) = \mathcal{L}_2 \cup \mathcal{D}_{2 \times 2,2} \cup \mathcal{L}_2,$$

with the underline set

$$D_{2,2 \times 2,2,2} = H_{2,2 \times 2} \cup H_{2,2} = \{-1, 0, a, b, c\} \cup \{c, d, 1\} = \{-1, 0, a, b, c, d, 1\},$$

is organized as a lattice as with the following tables:

\odot	-1	0	a	b	c	d	1	\rightarrow	-1	0	a	b	c	d	1
-1	-1	-1	-1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1
0	-1	0	0	0	0	0	0	0	-1	1	1	1	1	1	1
a	-1	0	a	0	a	a	a	a	-1	b	1	b	1	1	1
b	-1	0	0	b	b	b	b	b	-1	a	a	1	1	1	1
c	-1	0	a	b	c	c	c	c	-1	0	a	b	1	1	1
d	-1	0	a	b	c	c	d	d	-1	0	a	b	d	1	1
1	-1	0	a	b	c	d	1	1	-1	0	a	b	c	d	1

note that $\mathcal{D}_{2,2 \times 2,2,2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = c \neq 1$ thus $\mathcal{D}_{3,2 \times 2,2}$ is not a super BL -algebra.

- (8) Examples of the form: non-linearly ordered $BL \cup$ non-linearly ordered BL or equivalent forms.

Example. The bi - BL -algebra

$$\begin{aligned} \mathcal{D}_{2,2 \times 2,2,2 \times 2} &= \mathcal{H}_{2,2 \times 2} \cup \mathcal{H}_{2,2 \times 2} = (\mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}) \cup (\mathcal{L}_2 \cup \mathcal{L}_{2 \times 2}) \\ &= \mathcal{L}_2 \cup \mathcal{D}_{2 \times 2,2} \cup \mathcal{L}_{2 \times 2} = \mathcal{D}_{2,2 \times 2,2} \cup \mathcal{L}_{2 \times 2}, \end{aligned}$$

with the underline set

$$\begin{aligned} \mathcal{D}_{2,2 \times 2,2,2 \times 2} &= \mathcal{H}_{2,2 \times 2} \cup \mathcal{H}_{2,2 \times 2} = \{0, m, a, b, p\} \cup \{p, n, c, d, 1\} \\ &= \{0, m, a, b, p, n, c, d, 1\}, \end{aligned}$$

is organized as a lattice as with the following tables:

\odot	0	m	a	b	p	n	c	d	1	\rightarrow	0	m	a	b	p	n	c	d	1
0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1
m	0	m	m	m	m	m	m	m	m	m	0	1	1	1	1	1	1	1	1
a	0	m	a	m	a	a	a	a	a	a	0	b	1	b	1	1	1	1	1
b	0	m	m	b	b	b	b	b	b	b	0	a	a	1	1	1	1	1	1
p	0	m	a	b	p	p	p	p	p	p	0	m	a	b	1	1	1	1	1
n	0	m	a	b	p	n	n	n	n	n	0	m	a	b	p	1	1	1	1
c	0	m	a	b	p	n	c	n	c	c	0	m	a	b	p	d	1	1	1
n	0	m	a	b	p	n	n	d	d	n	0	m	a	b	p	c	c	1	1
1	0	m	a	b	p	n	c	d	1	1	0	m	a	b	p	n	c	d	1

note that $\mathcal{D}_{2,2 \times 2,2,2 \times 2}$ is not a BL -algebra, since $(a \rightarrow b) \vee (b \rightarrow a) = b \vee a = p \neq 1$ thus $\mathcal{D}_{2,2 \times 2,2,2 \times 2}$ is not a super BL -algebra.

3.4. *bi*-Homomorphisms, *bi*-Filters and *bi*-Boolean center

Definition 3.14. Let $L = (L_1 \cup L_2, \wedge, \vee, \odot, \rightarrow, 0, 1)$ and $K = (K_1 \cup K_2, \wedge, \vee, \odot, \rightarrow, 0, 1)$ be two *bi*- BL -algebras. We say a map ϕ from L to K is a *bi*-homomorphism of *bi*- BL -algebras. If $\phi = \phi_1 \cup \phi_2$ where $\phi_1 = \phi|_{L_1}$ from L_1 to K_1 and $\phi_2 = \phi|_{L_2}$ from L_2 to K_2 are BL -homomorphisms.

Definition 3.15. Let $\phi : L \rightarrow K$ be a *bi*-homomorphism, where $L = L_1 \cup L_2$ and $K = K_1 \cup K_2$ are *bi*- BL -algebras the kernel of the *bi*-homomorphism ϕ as $Ker(\phi) = Ker(\phi_1) \cup Ker(\phi_2)$; here $Ker(\phi_1) = \{a_1 \in L_1 \mid \phi_1(a_1) = 1\}$ and $Ker(\phi_2) = \{a_2 \in L_2 \mid \phi_2(a_2) = 1\}$, i.e., $Ker(\phi) = \{a_1 \in L_1, a_2 \in L_2 \mid \phi_1(a_1) = 1, \phi_2(a_2) = 1\}$.

Example 3.16. Let $L = \mathcal{D}_{2 \times 2,2}$ and $K = \mathcal{D}_{2 \times 2,3}$. Define $\phi = \phi_1 \cup \phi_2$ as follow: $\phi_1 : \mathcal{L}_{2 \times 2} \rightarrow \mathcal{L}_{2 \times 2}$ where ϕ_1 is a identity map and $\phi_2 : \mathcal{L}_2 \rightarrow \mathcal{L}_3$ where $\phi_2(c) = c$ and $\phi_2(1) = 1$, then ϕ is a *bi*-homomorphism from L to K and $Ker(\phi_1) = \{c\}$ and $Ker(\phi_2) = \{1\}$, so $Ker(\phi) = \{c, 1\}$.

Definition 3.17. Let $L = L_1 \cup L_2$ be a *bi*- BL -algebra. We say that subset $S = S_1 \cup S_2$ of L is a sub *bi*- BL -algebra of L if $L_1 \cap S = S_1$ and $L_2 \cap S = S_2$ are subalgebra of L_1 and L_2 respectively.

Example 3.18. In the Example 3.2, consider $S_1 = \{0, a, c, 1\}$ and $S_2 = \{0, e, 1\}$, then $S = S_1 \cup S_2 = \{0, a, c, e, 1\}$ is a sub *bi*- BL -algebra of L , since $S \cap L_1 = S_1$ and $S \cap L_2 = S_2$ are subalgebras of L_1 and L_2 respectively.

Definition 3.19. Let $L = L_1 \cup L_2$ be a *bi*- BL -algebras. We say the subset $F = F_1 \cup F_2$ of L is a *bi*-filter of L if F_i is a filter of L_i , where $i = 1, 2$ respectively.

Theorem 3.20. Let $L = L_1 \cup L_2$ and $K = K_1 \cup K_2$ are *bi*- BL -algebras and $\phi : L \rightarrow K$ is a *bi*- BL -algebra homomorphism. Then $Ker(\phi)$ is a *bi*-filter of L .

Example 3.21. In Example 3.2, consider $F_1 = \{a, 1\}$ and $F_2 = \{e, 1\}$. Then $F = F_1 \cup F_2 = \{a, e, 1\}$ is a *bi*-filter of L .

Theorem 3.22. Let $F = F_1 \cup F_2$ be a bi-filter of a bi-BL-algebra $L = L_1 \cup L_2$ such that F_i is a filter of L_i where $i = 1, 2$. Then $\frac{L}{F} := \frac{L_1}{F_1} \cup \frac{L_2}{F_2}$ is a bi-BL-algebra where $\frac{L_i}{F_i} = \{[x]_{F_i} | x \in L_i\}$ and $[x]_{F_i} = \{y \in L_i | x \rightarrow y \in F_i, y \rightarrow x \in F_i\}$, where $x \in L_i$ and $i = 1, 2$.

Definition 3.23. Let $F = F_1 \cup F_2$ be a bi-filter of a super BL-algebra $L = L_1 \cup L_2$. If F is a filter of L , then we say that F is a super filter of L .

Example 3.24. Let $L_1 = \{0, a, c, 1\}$ and $L_2 = \{0, b, c, d, 1\}$. Define \odot and \rightarrow as follow:

L_1	\odot	0	a	c	1
	0	0	0	0	0
	a	0	a	c	a
	c	0	c	c	c
	1	0	a	c	1

L_1	\rightarrow	0	a	c	1
	0	1	1	1	1
	a	0	1	c	1
	c	0	1	1	1
	1	0	a	c	1

L_2	\odot	0	b	c	d	1
	0	0	0	0	0	0
	b	0	b	c	d	b
	c	0	c	c	d	c
	d	0	d	d	0	d
	1	0	b	c	d	1

L_2	\rightarrow	0	b	c	d	1
	0	1	1	1	1	1
	b	0	1	c	d	1
	c	0	1	1	d	1
	d	d	1	1	1	1
	1	0	b	c	d	1

For L , whose tables are the following:

L	\odot	0	a	b	c	d	1
	0	0	0	0	0	0	0
	a	0	a	c	c	d	a
	b	0	c	b	c	d	b
	c	0	c	c	c	d	c
	d	0	d	d	d	0	d
	1	0	a	b	c	d	1

L	\rightarrow	0	a	b	c	d	1
	0	1	1	1	1	1	1
	a	0	1	b	b	d	1
	b	0	a	1	a	d	1
	c	0	1	1	1	d	1
	d	d	1	1	1	1	1
	1	0	a	b	c	d	1

Then L_1 and L_2 are BL -algebras and $L = L_1 \cup L_2$ is a bi - BL -algebra and also L is a super BL -algebra, consider $F_1 = \{a, 1\}$ and $F_2 = \{b, c, 1\}$ are filters of L_1 and L_2 , respectively. Also $F = F_1 \cup F_2 = \{a, b, c, 1\}$ is super filter of L .

Corollary 3.25. *In a super BL -algebra, any super filter is a filter.*

In the following example we show that converse of above corollary is not true.

Example 3.26. In super BL -algebra $\mathcal{H}_{2,2 \times 2}$, we consider $F = \{a, 1\}$. Then F is a filter of $\mathcal{H}_{2,2 \times 2}$ but is not a super filter of $\mathcal{H}_{2,2 \times 2}$, since F is not union each pair filters such that $F = F_1 \cup F_2$ and F_i is filter of L_i , respectively.

Proposition 3.27. *Let $L = L_1 \cup L_2$ be a super BL -algebra and $F = F_1 \cup F_2$ be a super filter of L , then $\frac{L}{F}$ is a BL -algebra.*

If $L = L_1 \cup L_2$ is a super BL -algebra and $F = F_1 \cup F_2$ a super filter of L , then $\frac{L}{F} \neq \frac{\mathcal{L}}{\mathcal{F}}$.

Example 3.28. In Example 3.2 we have:

$$\begin{aligned} [0]_{F_1} &= \frac{0}{F_1} = \{x \in L_1 | x \rightarrow 0 \in F_1, 0 \rightarrow x \in F_1\} = \{0\}, \\ [a]_{F_1} &= \frac{a}{F_1} = \{x \in L_1 | x \rightarrow a \in F_1, a \rightarrow x \in F_1\} = \{a, 1\}, \\ [c]_{F_1} &= \frac{c}{F_1} = \{x \in L_1 | x \rightarrow c \in F_1, c \rightarrow x \in F_1\} = \{c\}, \\ [1]_{F_1} &= \frac{1}{F_1} = \{x \in L_1 | x \rightarrow 1 \in F_1, 1 \rightarrow x \in F_1\} = \{a, 1\}, \end{aligned}$$

Thus $[a]_{F_1} = [1]_{F_1}$, therefore $\frac{L_1}{F_1} = \{[0]_{F_1}, [c]_{F_1}, [1]_{F_1}\}$.

$$\begin{aligned} [0]_{F_2} &= \frac{0}{F_2} = \{x \in L_2 | x \rightarrow 0 \in F_2, 0 \rightarrow x \in F_2\} = \{0\}, \\ [b]_{F_2} &= \frac{b}{F_2} = \{x \in L_2 | x \rightarrow b \in F_2, b \rightarrow x \in F_2\} = \{b, c, 1\}, \\ [c]_{F_2} &= \frac{c}{F_2} = \{x \in L_2 | x \rightarrow c \in F_2, c \rightarrow x \in F_2\} = \{b, c, 1\}, \\ [d]_{F_2} &= \frac{d}{F_2} = \{x \in L_2 | x \rightarrow d \in F_2, d \rightarrow x \in F_2\} = \{d\}, \\ [1]_{F_2} &= \frac{1}{F_2} = \{x \in L_2 | x \rightarrow 1 \in F_2, 1 \rightarrow x \in F_2\} = \{b, c, 1\}, \end{aligned}$$

Thus $[b]_{F_2} = [c]_{F_2} = [1]_{F_2}$, therefore $\frac{L_2}{F_2} = \{[0]_{F_2}, [d]_{F_2}, [1]_{F_2}\}$.

So $\frac{\mathcal{L}}{\mathcal{F}} = \frac{L_1}{F_1} \cup \frac{L_2}{F_2} = \{[0]_{F_1}, [c]_{F_1}, [1]_{F_1}, [0]_{F_2}, [d]_{F_2}, [1]_{F_2}\}$ is a bi - BL -algebra.

In $\frac{L}{F}$ we have:

$$\begin{aligned} [0]_F &= \frac{0}{F} = \{x \in L | x \rightarrow 0 \in F, 0 \rightarrow x \in F\} = \{0\}, \\ [a]_F &= \frac{a}{F} = \{x \in L | x \rightarrow a \in F, a \rightarrow x \in F\} = \{a, b, c, 1\}, \\ [b]_F &= \frac{b}{F} = \{x \in L | x \rightarrow b \in F, b \rightarrow x \in F\} = \{a, b, c, 1\}, \\ [c]_F &= \frac{c}{F} = \{x \in L | x \rightarrow c \in F, c \rightarrow x \in F\} = \{a, b, c, 1\}, \\ [d]_F &= \frac{d}{F} = \{x \in L | x \rightarrow d \in F, d \rightarrow x \in F\} = \{d\}. \end{aligned}$$

So $\frac{L}{F} = \{[0]_F, [d]_F, [1]_F\}$, we show that $\frac{L}{F} \neq \frac{\mathcal{L}}{\mathcal{F}}$, since $[1]_{F_1} \in \frac{\mathcal{L}}{\mathcal{F}}$, but $[1]_{F_1} \notin \frac{L}{F}$.

Definition 3.29. Let $A = A_1 \cup A_2$ be a *bi-BL*-algebra and $\mathcal{B}(A)$ be the Boolean *bi*-algebra associated with the bounded distributive lattice $L(A)$. Elements of $\mathcal{B}(A)$ are called the *bi*-Boolean elements of A and $\mathcal{B}(A) := B(A_1) \cup B(A_2)$, where $B(A_1)$ and $B(A_2)$ are Boolean elements of A_1 and A_2 , respectively.

Example 3.30. In Example 3.1, we have $B(L_1) = B(L_2) = \{0, 1\}$, then $\mathcal{B}(A) = B(L_1) \cup B(L_2) = \{0, 1\}$.

Example 3.31. In Example 3.3, we have $\mathcal{B}(A) = B(L_1) = B(L_2) = \{0, 1\}$ but the Boolean elements of L are $B(L) = \{0, a, d, 1\}$. Thus the *bi*-Boolean elements of a super *BL*-algebra are not equal with Boolean elements of L .

Remark 3.32. Suppose $L = L_1 \cup L_2$ be a chain super *BL*-algebra and also $1_{L_1} = 1_{L_2}$ and $0_{L_1} = 0_{L_2}$, then the *bi*-Boolean elements of L is equal the Boolean elements of L , i.e., $\mathcal{B}(L_1) = B(L_1) = B(L_2) = B(L)$.

Example 3.33. Let $L_1 = \{0, 1, 2, 4\}$ and $L_2 = \{0, 2, 3, 4\}$. Then $L_1 \cup L_2 = \{0, 1, 2, 3, 4\} = \mathcal{L}_5$. Define \odot and \rightarrow as follow:

\mathcal{L}_1	\odot	0	1	2	4	\rightarrow	0	1	2	4
	0	0	0	0	0	0	4	4	4	4
	1	0	0	0	1	1	2	4	4	4
	2	0	0	1	2	2	1	2	4	4
	4	0	1	2	4	4	0	1	2	4
\mathcal{L}_2	\odot	0	2	3	4	\rightarrow	0	2	3	4
	0	0	0	0	0	0	4	4	4	4
	2	0	0	0	2	2	3	4	4	4
	3	0	0	2	3	3	2	3	4	4
	4	0	2	3	4	4	0	2	3	4

L_1 and L_2 are BL -algebras, thus $\mathcal{L}_5 = L_1 \cup L_2$ is a super BL -algebra and we have $B(L_1) = B(L_2) = B(\mathcal{L}_5) = \mathcal{B}(\mathcal{L}_5) = \{0, 1\}$.

Otherwise suppose $L_1 = \{0, 1, 2, 3\}$ and $L_2 = \{0, 2, 3, 4\}$, then $L_1 \cup L_2 = \{0, 1, 2, 3, 4\} = \mathcal{L}_5$. Define \odot and \rightarrow as follow:

\mathcal{L}_1	\odot	0	1	2	3	\rightarrow	0	1	2	3
	0	0	0	0	0	0	3	3	3	3
	1	0	0	0	1	1	2	3	3	3
	2	0	0	1	2	2	1	2	3	3
	3	0	1	2	3	3	0	1	2	3
\mathcal{L}_2	\odot	0	2	3	4	\rightarrow	0	2	3	4
	0	0	0	0	0	0	4	4	4	4
	2	0	0	0	2	2	3	4	4	4
	3	0	0	2	3	3	2	3	4	4
	4	0	2	3	4	4	0	2	3	4

L_1 and L_2 are BL -algebras, thus $\mathcal{L}_5 = L_1 \cup L_2$ is a super BL -algebra and we have $B(L_1) = \{0, 3\}$ and $B(L_2) = \{0, 4\}$, then $\mathcal{B}(\mathcal{L}_5) = \{0, 3, 4\}$ but $B(\mathcal{L}_5) = \{0, 4\}$. So $\mathcal{B}(\mathcal{L}_5) \neq B(\mathcal{L}_5)$.

Definition 3.34. Let $L = L_1 \cup L_2$ be a bi - BL -algebra. We denote by $\mathcal{D}_s(L) = D_s(L_1) \cup D_s(L_2)$ the set of all bi -deductive systems of L where $D_s(L_1)$ and $D_s(L_2)$ are deductive systems of L_1 and L_2 respectively. If L is a super BL -algebra, then $D_s(L)$ is the set of all deductive systems of L .

Example 3.35. Consider $\mathcal{H}_{2,2 \times 2}$. Then $D_s(\mathcal{L}_2) = \{\{0\}, \mathcal{L}_2\}$ and $D_s(\mathcal{L}_{2 \times 2}) = \{\{1\}, \{a, 1\}, \{b, 1\}, \mathcal{L}_{2 \times 2}\}$, thus $\mathcal{D}_s(\mathcal{H}_{2,2 \times 2}) = D_s(\mathcal{L}_2) \cup D_s(\mathcal{L}_{2 \times 2}) = \{\{0\}, \{1\}, \{a, 1\}, \{b, 1\}, \mathcal{L}_2, \mathcal{L}_{2 \times 2}\}$. But $D_s(\mathcal{H}_{2,2 \times 2}) = \{\{1\}, \{a, 1\}, \{b, 1\}, \mathcal{L}_{2 \times 2}, \mathcal{H}_{2,2 \times 2}\}$, thus $\mathcal{D}_s(\mathcal{H}_{2,2 \times 2}) \neq D_s(\mathcal{H}_{2,2 \times 2})$, since $\mathcal{L}_2 \notin D_s(\mathcal{H}_{2,2 \times 2})$.

Remark 3.36. Let L be a bi - BL -algebra. Then $D_s(L)$ together with inclusion relation is not a lattice.

In above example $(\mathcal{D}_s(\mathcal{H}_{2,2 \times 2}), \subseteq)$ is not a lattice, since $\mathcal{L}_2 \cap \{b, 1\} = \{-1, 0\} \cap \{b, 1\} = \emptyset$ and $\emptyset \notin \mathcal{D}_s(\mathcal{H}_{2,2 \times 2})$.

4. CONCLUSION

The union of two subgroups, or two subrings, or two subsemigroups etc. do not form any algebraic structure but all of them find a nice *bialgebraic* structure as *bigroups*, *birings*, *bisemigroups* etc. Except for this *bialgebraic* structure these would remain only as sets without any nice algebraic structure on them. Further when these *bialgebraic* structures are defined on them they enjoy not only the inherited qualities of the algebraic structure from which they are taken but also several distinct algebraic properties that are not present in algebraic structures.

We introduced the notion of a *bi-BL*-algebra and study it in detail. After that the notions of a *bi-filter*, *bi-deductive system* and *bi-Boolean center* of a *bi-BL*-algebra are introduced. We have also presented classes of *bi-BL*-algebras and we stated relation between *bi-filters* and quotient *bi-BL*-algebra. Finally we show that the set of all deductive systems of a *bi-BL*-algebra together with inclusion relation is not a lattice.

Acknowledgments

The authors wish to thank the reviewers for their excellent suggestions that have been incorporated into this paper.

REFERENCES

- [1] A. Borumand Saeid, A. Ahadpanah and L. Torkzadeh, *Smarandache BL-algebra*, J. Applied Logic **8** (2010), 253–261. doi:10.1016/j.jal.2010.06.001
- [2] A. Borumand Saeid and S. Motamed, *Normal filters in BL-algebras*, World Applied Sci. J. **7** (Special Issue Appl. Math.), (2009), 70–76.
- [3] A. Borumand Saeid and S. Motamed, *Some Results in BL-algebras*, Math. Logic Quat **55** (6) (2009), 649–658. doi:10.1002/malq.200910025
- [4] D. Busneag and D. Piciu, *On the lattice of deductive systems of a BL-algebra*, Central Eur. J Math. **1** (2) (2003), 221–238. doi:10.2478/BF02476010
- [5] R. Cingoli, I.M.L. D’Ottaviano and D. Mundici, *Algebraic Foundations of Many-valued Reasoning*, Kluwer Academic publ., Dordrecht, 2000. doi:10.1007/978-94-015-9480-6
- [6] R. Cignoli, F. Esteva, L. Godo and A. Torrens, *Basic fuzzy logic is the logic of continuous t-norm and their residua*, Soft Comput **4** (2000), 106–112. doi:10.1007/s005000000044
- [7] A. Di Nola, G. Georgescu and A. Iorgulescu, *Pseudo BL-algebra: Part I*, Mult val logic **8** (5-6) (2002), 673–714.
- [8] A. Di Nola and L. Leustean, *Compact representations of BL-algebras*, Arch-Math. Logic **42** (2003), 737–761. doi:10.1007/s00153-003-0178-y

- [9] P. Hajek, *Metamathematics of Fuzzy Logic*, Kluwer Academic Publishers, Dordrecht, 1998. <http://dx.doi.org/10.1007/978-94-011-5300-3>
- [10] M. Haveski, A. Borumand Saeid and E. Eslami, *Some types of filters in BL-algebras*, *Soft Computing* **10** (2006), 657–664. doi:10.1007/s00500-005-0534-4
- [11] A. Iorgulescu, *Algebras of Logic as BCK-algebras*, Academy of Economic Studies, Bucharest, Editura 2008.
- [12] A. Iorgulescu, *Classes of BCK-algebra-part III*, Preprint series of the Institute of Mathematics of the Romanian Academy, preprint nr. 3/2004 (2004), 1–37.
- [13] M. Kondo and W.A. Dudck, *Filter theory of BL-algebras*, *Soft Computing* **12** (2007), 419–423.
- [14] R. Padilla, *Smarandache algebraic structures*, *Bull. Pure Appl. Sci., Delhi* **17** (1) (1998), 119–121.
- [15] D. Piciu, *Algebras of Fuzzy Logic*, Ed. Universitaria Craiova, 2007.
- [16] E. Turunen, *BL-algebras of basic fuzzy logic*, *Mathware and soft computing* **6** (1999), 49–61.
- [17] E. Turunen, *Boolean deductive systems of BL-algebras*, *Arch Math. Logic* **40** (2001), 467–473. doi:10.1007/s001530100088
- [18] E. Turunen, *Mathematics behind fuzzy logic*, Physica-Verlag, 1999.
- [19] W.B. Vasantha Kandasamy, *Bialgebraic structures and Smaranche bialgebraic structures*, American Research Press, 2003.

Received 14 August 2011
Revised 21 September 2011