

## ON PSEUDO $BE$ -ALGEBRAS

RAJAB ALI BORZOOEI

*Department of Mathematics*  
*Shahid Beheshti University of Tehran, Tehran, Iran*

**e-mail:** borzooei@sbu.ac.ir

ARSHAM BORUMAND SAEID

*Department of Mathematics*  
*Shahid Bahonar University of Kerman, Kerman, Iran*

**e-mail:** arsham@uk.ac.ir

AKBAR REZAEI

*Department of Mathematics, Payame Noor University*  
*P.O. Box. 19395-3697, Tehran, Iran*

**e-mail:** Rezaei@pnu.ac.ir

AKEFE RADFAR

*Department of Mathematics, Payame Noor University*  
*P.O. Box. 19395-3697, Tehran, Iran*

**e-mail:** Radfaratefe@gmail.com

AND

REZA AMERI

*Department of Mathematics, Tehran University*  
*Tehran, Iran*

**e-mail:** rameri@ut.ac.ir

### Abstract

In this paper, we introduce the notion of pseudo  $BE$ -algebra which is a generalization of  $BE$ -algebra. We define the concepts of pseudo subalgebras and pseudo filters and prove that, under some conditions, pseudo subalgebra can be a pseudo filter. We prove that every homomorphic image and pre-image of a pseudo filter is also a pseudo filter. Furthermore, the notion

of pseudo upper sets in pseudo  $BE$ -algebras introduced and is proved that every pseudo filter is an union of pseudo upper sets.

**Keywords:**  $BE$ -algebra, Pseudo  $BE$ -algebra, pseudo filter, pseudo upper set.

**2010 Mathematics Subject Classification:** 06F35, 03G25, 03B52.

## 1. INTRODUCTION

In 1966, Y. Imai and K. Iseki [8] introduced two classes of abstract algebras:  $BCK$ -algebras and  $BCI$ -algebras. It is known that the class of  $BCK$ -algebras is a proper subclass of the class of  $BCI$ -algebras. Some recent researchers led to generalizations of the notion of pseudo structure on some types of algebras. G. Georgescu and A. Iorgulescu [5], and independently J. Rachunek [13], introduced pseudo- $MV$  algebras which are a non-commutative generalization of  $MV$ -algebras. After pseudo- $MV$  algebras, the pseudo- $BL$  algebras [6], and the pseudo- $BCK$  algebras as an extended notion of  $BCK$ -algebras by G. Georgescu and A. Iorgulescu [7], were introduced and studied. A. Walendziak give a system of axioms defining pseudo- $BCK$  algebras [17]. In [9], Y.B. Jun and *et al.*, introduced the concepts of pseudo-atoms, pseudo- $BCI$  ideals and pseudo- $BCI$  homomorphisms in pseudo- $BCI$  algebras and characterizations of a pseudo- $BCI$  ideal, and provide conditions for a subset to be a pseudo- $BCI$  ideal. Y.H. Kim and K.S. So [11], discuss on minimal elements in pseudo- $BCI$  algebras.

The notion of  $BE$ -algebras was introduced by H.S. Kim and Y.H. Kim [10]. Using the notion of upper sets they gave an equivalent condition of the filter in  $BE$ -algebras.  $BE$ -algebras was deeply studied by S.S. Ahn and *et al.*, [1, 2, 3], Walendziak [16], A. Rezaei and *et al.*, [4, 14, 15]. Also, B.L. Meng [12], give a procedure by which generate a filter by a subset in a transitive  $BE$ -algebra, and give some characterizations of Noetherian and Artinian  $BE$ -algebras.

In the present paper we continue study in  $BE$ -algebras and introduce the notion of pseudo  $BE$ -algebras and pseudo filter in pseudo  $BE$ -algebras, also we discuss on the relationship between the pseudo sub algebras with pseudo filters in pseudo  $BE$ -algebras. We show that every pseudo filter is a pseudo subalgebra and give an example such that the converse is not true in general. The notion of pseudo upper set in pseudo  $BE$ -algebras is introduced and we show that every pseudo filter  $F$  of  $X$  is a union of pseudo upper sets (i.e.,  $F = \bigcup_{x \in F} A(x, 1)$ ).

Now, we review the definitions and some elementary aspects that are necessary for this paper.

Recall that a  $BE$ -algebra is an algebra  $(X; *, 1)$  of type  $(2, 0)$  satisfying the following axioms: ([10])

- (BE1)  $x * x = 1$ ,
- (BE2)  $x * 1 = 1$ ,
- (BE3)  $1 * x = x$ ,
- (BE4)  $x * (y * z) = y * (x * z)$ , for all  $x, y, z \in X$ .

A binary relation " $\leq$ " on  $BE$ -algebra  $X$  is defined by  $x \leq y$  if and only if  $x * y = 1$ .

A  $BE$ -algebra  $X$  has the following properties:

- (i)  $x * (y * x) = 1$ ,
- (ii)  $y * ((y * x) * x) = 1$ , for all  $x, y \in X$ .

A non-empty subset  $S$  of a  $BE$ -algebra  $X$  is called a subalgebra of  $X$  if  $x * y \in S$  whenever  $x, y \in S$ .

A mapping  $f : X \rightarrow Y$  of  $BE$ -algebras is called a homomorphism if  $f(x * y) = f(x) * f(y)$ , for all  $x, y \in X$ .

A non-empty subset  $F$  of  $BE$ -algebra  $X$  is called a filter of  $X$  if (1)  $1 \in F$ , (2)  $x \in F$  and  $x * y \in F$  implies  $y \in F$ .

Let  $A$  be a non-empty subset of  $BE$ -algebra  $X$ , then the set

$$\langle A \rangle = \bigcap \{G \in F(X) \mid A \subseteq G\}$$

is called the filter generated by  $A$ , written  $\langle A \rangle$ . If  $A = \{a\}$ , we will denote  $\langle \{a\} \rangle$ , briefly by  $\langle a \rangle$ , and we call it a principal filter of  $BE$ -algebra  $X$ . For  $F \in F(X)$  and  $a \in X$ , we denote by  $F_a$  the filter generated by  $F \cup \{a\}$ , where  $F(X)$  is the set of all filters of  $X$ .

## 2. PSEUDO $BE$ -ALGEBRAS

**Definition.** An algebra  $(X; *, \diamond, 1)$  of type  $(2, 2, 0)$  is called a pseudo  $BE$ -algebra if satisfies in the following axioms:

- (pBE1)  $x * x = 1$  and  $x \diamond x = 1$ ,
- (pBE2)  $x * 1 = 1$  and  $x \diamond 1 = 1$ ,
- (pBE3)  $1 * x = x$  and  $1 \diamond x = x$ ,
- (pBE4)  $x * (y \diamond z) = y \diamond (x * z)$ ,
- (pBE5)  $x * y = 1 \Leftrightarrow x \diamond y = 1$ ,

for all  $x, y, z \in X$ .

In a pseudo  $BE$ -algebra, we can introduce a binary relation " $\leq$ " by  $x \leq y \Leftrightarrow x * y = 1 \Leftrightarrow x \diamond y = 1$ , for all  $x, y \in X$ .

**Remark.** If  $X$  is a pseudo  $BE$ -algebra satisfying  $x * y = x \diamond y$ , for all  $x, y \in X$ , then  $X$  is a  $BE$ -algebra.

**Example 1.** (i) Let  $X = \{1, a, b, c\}$ . Define the operations " $*$ " and " $\diamond$ " as follow:

$*$	1	a	b	c
1	1	a	b	c
a	1	1	1	1
b	1	a	1	c
c	1	b	1	1

$\diamond$	1	a	b	c
1	1	a	b	c
a	1	1	1	1
b	1	c	1	c
c	1	c	1	1

Then  $(X; *, \diamond, 1)$  is a pseudo  $BE$ -algebra. We can see that  $(X; *, 1)$  and  $(X; \diamond, 1)$  are  $BE$ -algebras.

(ii) Let  $X = \{1, a, b, c\}$ . Define the operations " $*$ " and " $\diamond$ " on  $X$  as follow:

$*$	1	a	b	c
1	1	a	b	c
a	1	1	1	b
b	1	a	1	c
c	1	1	1	1

$\diamond$	1	a	b	c
1	1	a	b	c
a	1	1	1	a
b	1	a	1	a
c	1	1	1	1

Then  $(X; *, \diamond, 1)$  is a pseudo  $BE$ -algebra. Also, we can see that

$$a * (b * c) = a * c = b \neq b * (a * c) = b * b = 1.$$

Thus  $(X; *, 1)$  is not a  $BE$ -algebra.

(iii) Let  $X = (-\infty, -3] \cup \{-2, -1, 0, 1\}$ . Define the operations " $*$ " and " $\diamond$ " on  $X$  as follows:

$$x * y = \begin{cases} 1 & \text{if } x \leq y \\ 0 & \text{if } (x = 0 \text{ and } y = -2) \text{ or } (x = -1 \text{ and } y = -2) \\ y & \text{otherwise} \end{cases}$$

and

$$x \diamond y = \begin{cases} 1 & \text{if } x \leq y \\ 0 & \text{if } x = -1 \text{ and } y = -2 \\ y & \text{otherwise.} \end{cases}$$

Then  $(X; *, \diamond, 1)$  is a pseudo  $BE$ -algebra.

**Proposition 2.** *In a pseudo  $BE$ -algebra  $X$ , the following holds:*

- (1)  $x * (y \diamond x) = 1, x \diamond (y * x) = 1,$
- (2)  $x \diamond (y \diamond x) = 1, x * (y * x) = 1,$
- (3)  $x \diamond ((x \diamond y) * y) = 1, x * ((x * y) \diamond y) = 1,$
- (4)  $x * ((x \diamond y) * y) = 1, x \diamond ((x * y) \diamond y) = 1,$
- (5) *if  $x \leq y * z$ , then  $y \leq x \diamond z$ ,*
- (6) *if  $x \leq y \diamond z$ , then  $y \leq x * z$ ,*
- (7)  $1 \leq x$ , *implies  $x = 1$ ,*
- (8) *if  $x \leq y$ , then  $x \leq z * y$  and  $x \leq z \diamond y$ ,*
- (9) *if  $x * y = z$ , then  $y * z = y \diamond z = 1$  (If  $x \diamond y = z$ , then  $y * z = y \diamond z = 1$ ),*
- (10) *if  $x * y = x$  and  $x \neq 1$ , then  $x \diamond y \neq y$ ,*
- (11) *if  $x * y = y$  and  $x \neq 1$ , then  $x \diamond y \neq x$ ,*
- (12) *if  $x * y = x$  and  $x \diamond y = z$ , then  $x * z = x \diamond z = 1$ , and*

$$x * (y * z) = (x * y) * (x * z) = x \diamond (y * z) = (x \diamond y) * (x \diamond z) = 1,$$

- (13) *if  $x * y = y$  and  $x \diamond y = z$ , then  $x * z = z$  and*

$$x * (y * z) = (x * y) * (x * z) = x * (y \diamond z) = (x * y) \diamond (x * z) = 1,$$

- (14) *if  $x * y = z$  and  $x \diamond y = t$ , then  $x \diamond z = x * t$ ,*

for all  $x, y, z \in X$ .

**Proof.** (1) Let  $x, y \in X$ . By  $(pBE4)$ ,  $(pBE1)$  and  $(pBE2)$  we have

$$x * (y \diamond x) = y \diamond (x * x) = y \diamond 1 = 1$$

and

$$x \diamond (y * x) = y * (x \diamond x) = y * 1 = 1.$$

(2) By  $(pBE5)$  and (1) the proof is clear.

(3) Let  $x, y \in X$ . By  $(pBE4)$  and  $(pBE1)$  we have

$$x \diamond ((x \diamond y) * y) = (x \diamond y) * (x \diamond y) = 1$$

and

$$x * ((x * y) \diamond y) = (x * y) \diamond (x * y) = 1.$$

(4) By  $(pBE5)$  and (3) the proof is clear.

(5) Let  $x \leq y * z$  and  $x, y, z \in X$ . Then by (pBE5) and (pBE4) we have

$$x \diamond (y * z) = y * (x \diamond z) = 1.$$

Hence  $y \leq x \diamond z$ .

(6) The proof is similar to (5).

(7) Since  $1 \leq x$ , we have  $1 * x = 1 \diamond x = 1$ . Now, by (pBE3),  $x = 1 * x = 1 \diamond x = 1$ .

(8) Let  $z \in X$ . Since  $x * y = x \diamond y = 1$ , by using (pBE4), (pBE2) and (pBE5) we have

$$x \diamond (z * y) = z * (x \diamond y) = z * 1 = 1 \text{ and so } x * (z * y) = 1.$$

Thus  $x \leq z * y$ . By a similar way,  $x \leq z \diamond y$ .

(9) Let  $x * y = z$ . By (pBE4) and (pBE1),

$$y \diamond z = y \diamond (x * y) = x * (y \diamond y) = x * 1 = 1.$$

Now, by (pBE5), we get that  $y * z = 1$ .

(10) Let  $x * y = x$  and  $x \neq 1$ . If  $x \diamond y = y$ , then

$$x * (x \diamond y) = x * y = x, \quad x \diamond (x * y) = x \diamond x = 1,$$

and so by (pBE4),  $x = 1$ , which is a contradiction.

(11) Let  $x * y = y$  and  $x \neq 1$ . If  $x \diamond y = x$ , then

$$x * (x \diamond y) = x * x = 1, \quad x \diamond (x * y) = x \diamond y = x,$$

and so by (pBE4),  $x = 1$ , which is a contradiction.

(12) Let  $x * y = x$  and  $x \diamond y = z$ . Then

$$x * (x \diamond y) = x * z, \quad x \diamond (x * y) = x \diamond x = 1.$$

Hence by (pBE4), we get that  $x * z = 1$ . Since  $x \diamond y = z$ , by (9),  $y * z = y \diamond z = 1$ . By (pBE5),  $x * z = x \diamond z = 1$ . Then

$$x * (y * z) = x * 1 = 1, \quad (x * y) * (x * z) = x * 1 = 1.$$

Hence by (pBE5),  $x \diamond (y * z) = 1$ . Moreover,  $(x \diamond y) * (x \diamond z) = z * 1 = 1$ .

(13) Let  $x * y = y$  and  $x \diamond y = z$ . By (9),  $y * z = y \diamond z = 1$ . Then

$$x * (x \diamond y) = x * z, \quad x \diamond (x * y) = x \diamond y = z.$$

Hence by  $(pBE4)$ ,  $x * z = z$ . Then

$$x * (y \diamond z) = x * 1 = 1, \quad x * (y * z) = x * 1 = 1, \quad (x * y) * (x * z) = y * z = 1,$$

and so by  $(pBE5)$ ,  $(x * y) \diamond (x * z) = 1$ .

(14) If  $x * y = z$  and  $x \diamond y = t$ , then  $x \diamond z = x \diamond (x * y) = x * (x \diamond y) = x * t$ . ■

From now on  $X$  is a pseudo  $BE$ -algebra, unless otherwise is stated.

**Definition.** A subalgebra of  $X$  is a non-empty subset  $S$  of  $X$  which satisfies  $x * y \in S$  and  $x \diamond y \in S$ , for all  $x, y \in S$ .

**Example 3.** Let  $X = \{1, a, b, c, d\}$ . Define operations " $*$ " and " $\diamond$ " on  $X$  as follows:

$*$	1	a	b	c	d
1	1	a	b	c	d
a	1	1	a	1	a
b	1	1	1	1	a
c	1	a	a	1	a
d	1	1	1	1	1

$\diamond$	1	a	b	c	d
1	1	a	b	c	d
a	1	1	c	1	c
b	1	1	1	1	c
c	1	a	b	1	d
d	1	1	1	1	1

Then  $(X, *, \diamond)$  is a pseudo  $BE$ -algebra and  $\{1, a, b, c\}$  is a sub-algebra of  $X$ .

**Theorem 4.** If  $\{S_i\}_{i \in I}$  is a family of subalgebras of  $X$ , then  $\bigcap_{i \in I} S_i$  is a subalgebra of  $X$ , too.

**Definition.** A non-empty subset  $F$  of  $X$  is called a pseudo filter of  $X$  if it satisfies in the following axioms:

$$(pF1) \quad 1 \in F,$$

$$(pF2) \quad x \in F \text{ and } x * y \in F \text{ imply } y \in F.$$

A pseudo filter  $F$  is proper if and only if  $F \neq X$ .

A maximal pseudo filter is a proper filter such that it is not included in any other proper pseudo filter.

**Proposition 5.** Let  $F \subseteq X$  and  $1 \in F$ .  $F$  is a pseudo filter if and only if  $x \in F$  and  $x \diamond y \in F$  imply  $y \in F$ , for all  $x, y \in X$  ( $pF3$ ).

**Proof.** Let  $x \in F$  and  $x \diamond y \in F$ . By Proposition 2 (4), we have  $x * ((x \diamond y) * y) = 1$ . By  $(pF1)$ ,  $x * ((x \diamond y) * y) \in F$ . By assumption,  $(x \diamond y) * y \in F$ . Hence  $y \in F$ .

Conversely, let  $x \in F$  and  $x * y \in F$ . By Proposition 2 (4), we have  $x \diamond ((x * y) \diamond y) = 1$ . By (pF1),  $x \diamond ((x * y) \diamond y) \in F$ . From (pF3) it follows that  $(x * y) \diamond y \in F$ . Hence  $y \in F$ . ■

**Example 6.** (i)  $\{1\}$  and  $X$  are pseudo filters of  $X$ , in which we call trivial pseudo filters.

(ii) In Example 1(ii),  $F = \{1, b\}$  is a pseudo filter, but  $G = \{1, a\}$  is not a pseudo filter, because  $a * c = 1 \in G$  and  $a \in G$  but  $c \notin G$ .

**Proposition 7.** *Let  $F$  be a pseudo filter of  $X$  and  $y \in X$ . If  $x \leq y$  and  $x \in F$ , then  $y \in F$ .*

**Theorem 8.** *Any pseudo filter of  $X$  is a pseudo subalgebra.*

**Proof.** Assume that  $F$  is a pseudo filter of  $X$  and  $x, y \in F$ . Since  $x \leq y * x$ , and by Proposition 7 we get that  $y * x, y \diamond x \in F$ . Hence  $F$  is a pseudo subalgebra of  $X$ . ■

**Note.** The converse of Theorem 8, is not true in general. Because in Example 1(ii),  $S = \{1, a\}$  is a pseudo subalgebra, but it is not a pseudo filter.

**Theorem 9.** *If  $\{F_i\}_{i \in I}$  is a family of pseudo filters of  $X$ , then  $\bigcap_{i \in I} F_i$  is a pseudo filter of  $X$ , too.*

**Proposition 10.** *If  $F$  is a pseudo filter of  $X$ , then*

- (i)  $\forall x, y, z \in X, x, y \in F, x \leq y * z \Rightarrow z \in F$ ,
- (ii)  $\forall x, y, z \in X, x, y \in F, x \leq y \diamond z \Rightarrow z \in F$ .

**Proof.** Assume that  $F$  is a pseudo filter of  $X$  and  $x, y, z \in X$  such that  $x, y \in F$  and  $x \leq z * y$ . Then  $x \diamond (y * z) = 1 \in F$  and so  $y * z \in F$ . Since  $y \in F$ ,  $z \in F$ .

Now, let  $x, y, z \in X$  such that  $x, y \in F$  and  $x \leq y \diamond z$ . Then  $x * (y \diamond z) = 1 \in F$ , and so  $y \diamond z \in F$ . Since  $y \in F$ ,  $z \in F$ . ■

**Theorem 11.** *Let  $F$  be a subset of  $X$  containing 1. Then  $F$  is pseudo filter of  $X$  if and only if  $x \leq y * z$  imply  $z \in F$ , for all  $x, y \in F$  and  $x \in X$ .*

**Proof.** ( $\Rightarrow$ ). By Proposition 10, the proof is clear.

( $\Leftarrow$ ). By assumption,  $1 \in F$ . Let  $x, x \diamond y \in F$  and  $y \in X$ . Proposition 2(3) implies that  $x \diamond ((x \diamond y) * y) = 1$  and hence  $x \leq (x \diamond y) * y$ . Since  $x, x \diamond y \in F$ , we conclude that  $y \in F$ . By Proposition 5, we get that  $F$  is a pseudo filter. ■



For element  $a$  of  $X$ , put

$$H(a) := \{x \in X : a \leq x\},$$

which is called the terminal section of an element  $a$ . Since  $1, a \in H(a)$ , we can see that  $H(a)$  is not an empty set.

**Theorem 12.** *For any element  $a$  of  $X$ , the terminal section  $H(a)$  is a pseudo filter if and only if the following implications hold:*

- (i)  $\forall x, y, z \in X, \quad z \leq x * y, \text{ and } z \leq x \Rightarrow z \leq y;$
- (ii)  $\forall x, y, z \in X, \quad z \leq x \diamond y, \text{ and } z \leq x \Rightarrow z \leq y.$

**Proof.** Assume that for each  $a \in X$ ,  $H(a)$  is a pseudo filter of  $X$ . Let  $x, y, z \in X$  be such that  $z \leq x * y$ ,  $z \leq x \diamond y$ , and  $z \leq x$ . Then  $x * y \in H(z)$ ,  $x \diamond y \in H(z)$  and  $x \in H(z)$ . Since  $H(z)$  is pseudo filter,  $y \in H(z)$ . Therefore,  $z \leq y$ .

Conversely, consider  $H(z)$ , for  $z \in X$ . Obviously  $1 \in H(z)$ . Let  $x * y \in H(z)$ , for all  $x \in H(z)$ , i.e.,  $z \leq x * y$ , and  $z \leq x$ . Then from hypothesis,  $z \leq y$  i.e.,  $y \in H(z)$ . Hence  $H(z)$  is a pseudo filter of  $X$ , for all  $z \in X$ . ■

In the following, we provide some conditions for a subalgebra to be a pseudo filter.

**Theorem 13.** *Let  $F$  be a subalgebra of  $X$ . Then  $F$  is a pseudo filter of  $X$  if and only if for all  $x, y \in X$ ,*

$$x \in F, y \in X \setminus F \Rightarrow x * y \in X \setminus F \text{ and } x \diamond y \in X \setminus F.$$

**Proof.** Assume that  $F$  is a pseudo filter of  $X$  and  $x, y \in X$ , such that  $x \in F$  and  $y \in X \setminus F$ . If  $x * y \notin X \setminus F$ , then  $x * y \in F$ , i.e.,  $y \in F$ , which is a contradiction. Hence  $x * y \in X \setminus F$ . Now, if  $x \diamond y \notin X \setminus F$ , then  $x \diamond y \in F$ , i.e.,  $y \in F$ , which is a contradiction. Hence  $x \diamond y \in X \setminus F$ .

Conversely, assume that the hypothesis is valid. Since  $F$  is a subalgebra, then  $1 \in F$ . For every  $x \in F$ , let  $x * y \in F$ . If  $y \notin F$ , then  $x * y \in X \setminus F$  by assumption, which is a contradiction. Hence  $y \in F$ . Therefore,  $F$  is a pseudo filter. ■

**Definition.** A mapping  $f : X \rightarrow Y$  of pseudo  $BE$ -algebras is called a pseudo homomorphism if  $f(x * y) = f(x) * f(y)$  and  $f(x \diamond y) = f(x) \diamond f(y)$ , for all  $x, y \in X$ .

Note that if  $f : X \rightarrow Y$  is a pseudo homomorphism, then  $f(1_X) = 1_Y$ .

**Theorem 14.** *Let  $f : X \rightarrow Y$  be a pseudo homomorphism. Then*

- (i) *if  $F$  is a pseudo filter of  $Y$ , then  $f^{-1}(F)$  is a pseudo filter of  $X$ ,*

- (ii) if  $f$  is surjective and  $G$  is a pseudo filter of  $X$  containing  $\ker(f)$ , then  $f(G)$  is a pseudo filter of  $Y$ .

**Proof.** (i) Assume that  $F$  is a pseudo filter of  $Y$ . Obviously  $1_X \in f^{-1}(F)$ . Let  $x, x * y \in f^{-1}(F)$ . It follows that  $f(x) * f(y) = f(x * y) \in F$ . Then  $f(y) \in F$ , since  $f(x) \in F$  and  $F$  is a pseudo filter of  $Y$ . Therefore,  $y \in f^{-1}(F)$  and hence  $f^{-1}(F)$  is a pseudo filter of  $X$ .

(ii) Obviously  $1_Y \in f(G)$ . Let  $x, x * y \in f(G)$  and  $y \in Y$ . Since  $f$  is surjective, there exists  $c \in X$  such that  $f(c) = y$ . Let  $a, b \in G$  be such that  $f(a) = x$  and  $f(b) = x * y$ . We have

$$f(b) = x * y = f(a) * f(c) = f(a * c).$$

Then  $f(b * (a * c)) = 1_Y$  and hence  $b * (a * c) \in \ker(f) \subseteq G$ . Therefore,  $b * (a * c) \in G$ . Since  $b \in G$ , we see that  $a * c \in G$ , and so  $c \in G$ . Hence  $y = f(c) \in f(G)$ . Consequently,  $f(G)$  is a pseudo filter of  $Y$ . ■

In the following example we show that if  $f$  is not surjective in Theorem 14(ii), then we can not implies that  $f(G)$  is a pseudo filter of  $Y$ .

**Example 15.** Let  $Y$  be pseudo  $BE$ -algebra in Example 3(i), and  $X = \{1, a, b, c\}$ . Define the operations " $*$ " and " $\diamond$ " on  $X$  as follows:

$*$	1	a	b	c
1	1	a	b	c
a	1	1	a	1
b	1	1	1	1
c	1	a	a	1

$\diamond$	1	a	b	c
1	1	a	b	c
a	1	1	c	1
b	1	1	1	1
c	1	a	b	1

Then  $(X; *, \diamond, 1)$  is a pseudo  $BE$ -algebra. Now, if we consider  $f : X \rightarrow Y$  as the identity map, then  $f$  is a pseudo homomorphism and  $f(X) = X$ . Can easily seen that  $X = \{1, a, b, c\}$  is a trivial pseudo filter of  $X$ , but  $f(X)$  is not a pseudo filter of  $Y$ , because

$$a * d = a \in f(X), \quad a \in f(X) \text{ but } d \notin f(X).$$

**Corollary 16.** Let  $f : X \rightarrow Y$  be a pseudo homomorphism. Then

$$\ker(f) := \{x \in X | f(x) = 1_Y\}$$

is a pseudo filter of  $X$ .

**Proof.** Obviously,  $1_X \in \ker(f)$ . Hence  $\emptyset \neq \ker(f)$ . Let  $x, x * y \in \ker(f)$ . Then  $f(x) = f(x * y) = 1_Y$ . Therefore,  $1_Y = f(x * y) = f(x) * f(y) = 1_Y * f(y) = f(y)$  and  $1_Y = f(x \diamond y) = f(x) \diamond f(y) = 1_Y \diamond f(y) = f(y)$ . Thus  $y \in \ker(f)$ . ■

3. PSEUDO UPPER SET IN  $BE$  ALGEBRAS

Let  $x, y \in X$ . Define

$$A(x, y) := \{z \in X : x * (y \diamond z) = 1\}.$$

We call  $A(x, y)$  an pseudo upper set of  $x$  and  $y$ .

**Note.** It is easy to see that,  $1, x, y \in A(x, y)$ . The set  $A(x, y)$ , where  $x, y \in X$ , is not a pseudo filter of  $X$ , in general. Also, by using  $(pBE4)$  and  $(pBE5)$  we have

$$\begin{aligned} A(x, y) &:= \{z \in X : x * (y \diamond z) = 1\} = \{z \in X : y \diamond (x * z) = 1\} \\ &= \{z \in X : y * (x * z) = 1\} \\ &= \{z \in X : y \diamond (x \diamond z) = 1\} \\ &= \{z \in X : y * (x \diamond z) = 1\} = A(y, x). \end{aligned}$$

**Example 17.** Let  $X$  be a pseudo  $BE$ -algebra in Example 1(ii). It is easy to check that  $A(b, c) = \{1, b, c\}$ . Now, we can see that  $c * a = b \in A(b, c)$ ,  $c \diamond a = c \in A(b, c)$  and  $c \in A(b, c)$ , but  $a \notin A(b, c)$ . Thus  $A(b, c)$  is not a pseudo filter of  $X$ .

**Proposition 18.** (i)  $A(x, 1) \subseteq A(x, y)$ ,

(ii) if  $A(x, 1)$  is a pseudo filter of  $X$  and  $y \in A(x, 1)$ , then  $A(x, y) \subseteq A(x, 1)$ .

(iii) if there is  $y \in X$ , such that  $y * z = 1$ , for all  $z \in X$ , then  $A(x, y) = X = A(y, x)$ , for all  $x, y \in X$ .

**Proof.** (i) Let  $z \in A(x, 1)$ . Then  $x * (1 \diamond z) = x * z = 1$ . Hence  $x * (y \diamond z) = y \diamond (x * z) = y \diamond 1 = 1$ . Thus  $z \in A(x, y)$  and so  $A(x, 1) \subseteq A(x, y)$ .

(ii) Let  $A(x, 1)$  be a pseudo filter,  $y \in A(x, 1)$  and  $z \in A(x, y)$ . Then  $x * (y \diamond z) = 1$ . Hence  $y \diamond z \in A(x, 1)$ . Now, since  $A(x, 1)$  is a pseudo filter and  $y \in A(x, 1)$ ,  $z \in A(x, 1)$ . Thus  $A(x, y) \subseteq A(x, 1)$ .

(iii) Obviously  $A(x, y) \subseteq X$ , for all  $x \in X$ . Let  $z \in X$ . Then by assumption there is an  $y \in X$  such that  $y * z = 1$ . Hence we have  $1 = x \diamond 1 = x \diamond (y * z)$ . Therefore,  $z \in A(x, y)$ . Thus  $X \subseteq A(x, y)$  and so  $X = A(x, y)$ . ■

**Proposition 19.** Let  $x \in X$ . Then  $A(x, 1) = \bigcap_{y \in X} A(x, y)$ .

**Proof.** By Proposition 18(i), we have  $A(x, 1) \subseteq \bigcap_{y \in X} A(x, y)$ . Let  $z \in \bigcap_{y \in X} A(x, y)$ . Then  $z \in A(x, y)$ , for all  $y \in X$ , and so  $z \in A(x, 1)$ . Hence  $\bigcap_{y \in X} A(x, y) \subseteq A(x, 1)$ . ■

**Theorem 20.** Let  $F$  be a non-empty subset of  $X$ . Then  $F$  is a pseudo filter of  $X$  if and only if  $A(x, y) \subseteq F$ , for all  $x, y \in F$ .

**Proof.** Let  $F$  be a pseudo filter of  $X$  and  $x, y \in F$ . If  $z \in A(x, y)$ , then  $x * (y \diamond z) = 1 \in F$ . Since  $F$  is a pseudo filter and  $x, y \in F$ , by using  $(pF2)$ ,  $y \diamond z \in F$  and so by Proposition 5,  $z \in F$ . Hence  $A(x, y) \subseteq F$ .

Conversely, suppose that  $A(x, y) \subseteq F$ , for all  $x, y \in F$ . Since  $x * (y \diamond 1) = x * 1 = 1$ , we have  $1 \in A(x, y) \subseteq F$ . Let  $a, a * b \in F$ . Since  $1 = (a * b) \diamond (a * b) = a * ((a * b) \diamond b)$ , we have  $b \in A(a, a * b) \subseteq F$ . Hence  $b \in F$ . Consequently,  $F$  is a pseudo filter of  $X$ . ■

**Theorem 21.** *If  $F$  is a pseudo filter of  $X$ , then*

$$F = \bigcup_{x, y \in F} A(x, y).$$

**Proof.** Let  $F$  be a pseudo filter of  $X$  and  $z \in F$ . Since  $z * (1 \diamond z) = z * z = 1$ , we have  $z \in A(z, 1)$ . Hence

$$F \subseteq \bigcup_{z \in F} A(z, 1) \subseteq \bigcup_{x, y \in F} A(x, y)$$

If  $z \in \bigcup_{x, y \in F} A(x, y)$ , then there exist  $a, b \in F$  such that  $z \in A(a, b)$ . Hence by Theorem 20, we have  $z \in F$ . This means that  $\bigcup_{x, y \in F} A(x, y) \subseteq F$ . ■

**Theorem 22.** *If  $F$  is a pseudo filter of  $X$ , then*

$$F = \bigcup_{x \in F} A(x, 1).$$

**Proof.** Let  $F$  be a pseudo filter of  $X$  and  $z \in F$ . Since  $z * (1 \diamond z) = z * z = 1$ , we have  $z \in A(z, 1)$ . Hence

$$F \subseteq \bigcup_{z \in F} A(z, 1).$$

If  $z \in \bigcup_{x \in F} A(x, 1)$ , then there exists an  $a \in F$  such that  $z \in A(a, 1)$ , which means that  $a * z = a * (1 \diamond z) = 1 \in F$ . Since  $F$  is a pseudo filter of  $X$  and  $a \in F$ , we see that  $z \in F$ . This means that  $\bigcup_{x \in F} A(x, 1) \subseteq F$ . ■

#### 4. CONCLUSION AND FUTURE WORK

In this paper we generalized the notion of  $BE$ -algebras and introduced the notion of pseudo  $BE$ -algebras, pseudo subalgebras, pseudo filters and pseudo upper set. Furthermore, we prove that every pseudo filter is an union of pseudo upper sets.

We believe that such results are very useful for study in this structure and applied for other algebraic structures. Since quotient algebras are a basic tool for exploring the structures of pseudo  $BE$ -algebras and there are close contacts

among pseudo filters, pseudo congruence and quotient algebras. In the future work, we are going to consider the notion of the pseudo filters on distributive pseudo  $BE$ -algebras to get a quotient algebras induced by this filters. Also, we define pseudo congruence relations on pseudo  $BE$ -algebras and give the construction of quotient algebras. We try to define pseudo commutative  $BE$ -algebras and other types of pseudo filters in pseudo  $BE$ -algebras and discuss on relationship between them.

### Acknowledgments

The authors wish to thank the reviewers for their excellent suggestions that have been incorporated into this paper.

### REFERENCES

- [1] S.S. Ahn and Y.H. So, *On ideals and upper sets in  $BE$ -algebras*, Sci. Math. Jpn. **68** (2) (2008) 279–285.
- [2] S.S. Ahn and K.S. So, *On generalized upper sets in  $BE$ -algebras*, Bull. Korean Math. Soc. **46** (2) (2009) 281–287. doi:10.4134/BKMS.2009.46.2.281
- [3] S.S. Ahn, Y.H. Kim and J.M. Ko, *Filters in commutative  $BE$ -algebras*, Bull. Korean Math. Soc. **27** (2) (2012) 233–242. doi:10.4134/CKMS.2012.27.2.233
- [4] A. Borumand Saeid, A. Rezaei and R.A. Borzoei, *Some types of filters in  $BE$ -algebras*, (submitted).
- [5] G. Georgescu and A. Iorgulescu, *Pseudo-MV algebras: a noncommutative extension of MV algebras*, Information Technology (Bucharest, 1999), 961–968, Inforec, Bucharest.
- [6] G. Georgescu and A. Iorgulescu, *Pseudo-BL algebras: a noncommutative extension of BL algebras*, in Abstracts of the Fifth International Conference FSTA 2000, Slovakia, February 2000, 90–92.
- [7] G. Georgescu and A. Iorgulescu, *Pseudo-BCK algebras: an extension of BCK-algebras*, Combinatorics, computability and logic, 97–114, Springer Ser. Discrete Math. Theor. Comput. Sci., Springer, London, 2001.
- [8] Y. Imai and K. Iseki, *On axiom systems of propositional Calculi*, XIV proc. Jpn. Academy **42** (1966) 19–22.
- [9] Y.B. Jun, H.S. Kim and J. Neggers, *On pseudo-BCI ideals of pseudo-BCI algebras*, Math. Rec. **58** (2006) 39–46.
- [10] H.S. Kim and Y.H. Kim, *On  $BE$ -algebras*, Sci. Math. Jpn. **66** (1) (2007) 113–117.
- [11] Y.H. Kim and K.S. So, *On minimality in pseudo-BCI algebras*, Commun. Korean Math. Soc. **27** (1) (2012) 7–13. doi:10.4134/CKMS.2012.27.1.007
- [12] B.L. Meng, *On filters in  $BE$ -algebras*, Sci. Math. Jpn. **71** (2010), 201–207.

- [13] J. Rachunek, *A non commutative generalization of MV-algebras*, Czechoslovak Math. J. **52** (127) (2002) 255–273.
- [14] A. Rezaei and A. Borumand Saeid, *Some results in BE-algebras*, Analele Universitatii Oradea Fasc. Matematica, Tom XIX (2012), 33–44.
- [15] A. Rezaei and A. Borumand Saeid, *Commutative ideals in BE-algebras*, Kyungpook Math. J. **52** (2012) 483–494. doi:10.5666/KMJ.2012.52.4.483
- [16] A. Walendziak, *On commutative BE-algebras*, Sci. Math. Jpn. **69** (2) (2008) 585–588.
- [17] A. Walendziak, *On axiom systems of pseudo-BCK algebras*, Bull. Malays. Math. Sci. Soc. **34** (2) (2011) 287–293.

Received 22 February 2013

Revised 11 May 2013