

LIE IDEALS IN PRIME Γ -RINGS WITH DERIVATIONS

NISHTEMAN N. SULIMAN

*Salahaddin University,
Department of Mathematics,
Erbil, Iraq*

e-mail: vananesh@gmail.com

AND

ABDUL-RAHMAN H. MAJEED

*Baghdad University,
Department of Mathematics,
Baghdad, Iraq*

e-mail: Ahmajeed@yahoo.com

Abstract

Let M be a 2 and 3-torsion free prime Γ -ring, d a nonzero derivation on M and U a nonzero Lie ideal of M . In this paper it is proved that U is a central Lie ideal of M if d satisfies one of the following

- (i) $d(U) \subset Z$,
- (ii) $d(U) \subset U$ and $d^2(U) = 0$,
- (iii) $d(U) \subset U$, $d^2(U) \subset Z$.

Keywords: prime Γ -rings, Lie ideals, derivations.

2010 Mathematics Subject Classification: Primary: 16Y99; Secondary: 16W25, 16N60.

1. INTRODUCTION

The concept of a Γ -ring was first introduced by Nobusawa [5], and generalized by Barnes [1] as follows: A Γ -ring is a pair (M, Γ) where M and Γ are additive abelian groups for which there exists a map from $M \times \Gamma \times M$ to M (the image of (x, α, y) was denoted by $x\alpha y$) such that

- (i) $(x + y)\alpha z = x\alpha z + y\alpha z,$
 $x(\alpha + \beta)y = x\alpha y + x\beta y,$
 $x\alpha(y + z) = x\alpha y + x\alpha z,$
- (ii) $(x\alpha y)\beta z = x\alpha(y\beta z),$

for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$, then M is called a Γ -ring.

Recall that a Γ -ring M is called prime if for any two elements $x, y \in M$, $x\Gamma M\Gamma y = 0$ implies either $x = 0$ or $y = 0$, and M is called semiprime if $x\Gamma M\Gamma x = 0$ with $x \in M$ implies $x = 0$. Note that every prime Γ -ring is obviously semiprime. An additive mapping $d: M \rightarrow M$ is called a derivation if $d(x\alpha y) = d(x)\alpha y + x\alpha d(y)$ for all $x, y \in M$ and $\alpha \in \Gamma$. An additive subgroup I of M is called a left (right) ideal of M if $M\Gamma I \subset I$ ($I\Gamma M \subset I$). If I is both left and right ideal of M , then we say I is an ideal of M . The set $Z = \{x \in M; x\alpha y = y\alpha x \text{ for all } x, y \in M \text{ and } \alpha \in \Gamma\}$ is called the center of M . An additive subgroup U of M is said to be a Lie ideal of M if $[u, x]_\alpha \in U$, for all $u \in U, x \in M$ and $\alpha \in \Gamma$. M is n -torsion free if $nx = 0$, for $x \in M$ implies $x = 0$, where n is an integer. The commutator $x\alpha y - y\alpha x$ will be denoted by $[x, y]_\alpha$. We will use for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$, the basic commutator identities:

$$\begin{aligned} [x\alpha y, z]_\beta &= x\alpha[y, z]_\beta + [x, z]_\beta\alpha y + x[\alpha, \beta]_zy, \text{ and} \\ [x, y\alpha z]_\beta &= y\alpha[x, z]_\beta + [x, y]_\beta\alpha z + y[\beta, \alpha]_xz. \end{aligned}$$

Throughout this paper, We consider the following assumption $x\alpha y\beta z = x\beta y\alpha z$, for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$ and it will be represented by property (*) is a central.

According to the assumption property (*), the above two identities reduced to

$$\begin{aligned} [x\alpha y, z]_\beta &= x\alpha[y, z]_\beta + [x, z]_\beta\alpha y, \text{ and} \\ [x, y\alpha z]_\beta &= y\alpha[x, z]_\beta + [x, y]_\beta\alpha z. \end{aligned}$$

The relationship between the derivations and Lie ideals of a prime ring has been investigated by a number of authors (see [2, 3] and [4]). In [2], Bergen, Herstein and Kerr showed that if U is a nonzero Lie ideal of a 2-torsion free prime ring R and d a nonzero derivation of R such that $d^2(U) = 0$ or $d^2(U) \subset Z$ then U is central. Our aim in this paper is generalized the above results in prime Γ -rings with Lie ideals.

2. THE RESULTS

For proving the main results, we have needed some important lemmas. So we start as follows:

Remark 1. Let M be 2-torsion free prime Γ -ring and d a derivation of M . Then for all $x, y \in M$ and $\alpha \in \Gamma$, we have the followings:

- (i) If $d^2 = 0$ on M , then $d = 0$,
- (ii) $d([x, y]_\alpha) = [d(x), y]_\alpha + [x, d(y)]_\alpha$,
- (iii) $d^2(x\alpha y) = d^2(x)\alpha y + 2d(x)\alpha d(y) + x\alpha d^2(y)$,
- (iv) $d^3(x\alpha y) = d^3(x)\alpha y + 3d^2(x)\alpha d(y) + 3d(x)\alpha d^2(y) + x\alpha d^3(y)$.

Lemma 2 ([6], Lemma 1). *Let M be 2-torsion free prime Γ -ring and Z the center of M . Then the following are satisfied:*

- (i) *If $x \in Z$, and $x\Gamma y = 0$, then either $x = 0$ or $y = 0$.*
- (ii) *If $x \in Z$, and $x\Gamma y \subset Z$, then either $x = 0$ or $y \in Z$.*

Lemma 3 ([3], Lemma 2). *Let $0 \neq U$ be a Lie ideal of a 2-torsion free prime Γ -ring M and $U \not\subseteq Z$. If for $a, b \in M$ such that $a\Gamma U\Gamma b = 0$, then $a = 0$ or $b = 0$.*

Lemma 4. *Let U be a nonzero Lie ideal of prime Γ -ring M . If $[M, U]_\alpha \subset Z$, then $U \subset Z$.*

Proof. For all $x \in M, u \in U$ and $\alpha \in \Gamma$, we have $[x, u]_\alpha \in [M, U]_\alpha$. Replacing x by $x\beta u$, we get

$$[x\beta u, u]_\alpha = [x, u]_\alpha \beta u \in Z, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Since $[x, u]_\alpha \in Z$, then by Lemma 2(ii) we obtain $[x, u]_\alpha = 0$ or $u \in Z$, then the result required. \blacksquare

Lemma 5. *Let $0 \neq U$ be a Lie ideal of 2-torsion free prime Γ -ring M satisfying property (*). If $[U, U]_\Gamma = 0$, then $U \subset Z$ (If U is a commutative Lie ideal, then U is central).*

Proof. For all $x \in M, u \in U$ and $\alpha \in \Gamma$, we have $[u, x]_\alpha \in U$. Hence by hypothesis we have

$$[u, [u, x]_\alpha]_\beta = 0, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Equivalently

$$(1) \quad u\beta[u, x]_\alpha = [u, x]_\alpha \beta u, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Replacing x by $x\alpha y$, for $y \in M$ and $\alpha \in \Gamma$, we get

$$(2) \quad u\beta x\alpha[u, y]_\alpha + u\beta[u, x]_\alpha \alpha y = x\alpha[u, y]_\alpha \beta u + [u, x]_\alpha \alpha y \beta u.$$

Using (1) for $u\beta[u, x]_\alpha = [u, x]_\alpha\beta u$ and $[u, y]_\alpha\beta u = u\beta[u, y]_\alpha$ in (2) we obtain

$$u\beta x\alpha[u, y]_\alpha + [u, x]_\alpha\beta u\alpha y = x\alpha u\beta[u, y]_\alpha + [u, x]_\alpha\alpha y\beta u.$$

Using property (*) we get $2[u, x]_\alpha\beta[u, y]_\alpha = 0$. Since M is 2-torsion free, this leads to

$$[u, x]_\alpha\beta[u, y]_\alpha = 0, \text{ for all } x, y \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Replacing y by $y\gamma x$, we find that

$$[u, x]_\alpha\beta y\gamma[u, x]_\alpha = 0, \text{ for all } x, y \in M, u \in U \text{ and } \alpha, \beta, \gamma \in \Gamma.$$

Thus $[u, x]_\Gamma\Gamma M\Gamma[u, x]_\Gamma = 0$, for all $x \in M, u \in U$. By primeness of M , we conclude $[u, x]_\Gamma = 0$, yields $U \subset Z$. ■

Lemma 6. *Let U be a nonzero Lie ideal of 2-torsion free prime Γ -ring M and d a nonzero derivation of M . If $a \in U$ such that $[a, d(x)]_\alpha = 0$, for all $x \in M$ and $\alpha \in \Gamma$, then $a \in Z$.*

Proof. By hypothesis we have $[a, d(x)]_\alpha = 0$, for all $x \in M$ and $\alpha \in \Gamma$.

Replacing x by $x\beta y$, we get

$$\begin{aligned} 0 &= [a, d(x\beta y)]_\alpha \\ &= [a, d(x)]_\alpha\beta y + d(x)\beta[a, y]_\alpha + x\beta[a, d(y)]_\alpha + [a, x]_\alpha\beta d(y) \\ &= d(x)\beta[a, y]_\alpha a + [a, x]_\alpha\beta d(y). \end{aligned}$$

Replacing x by $d(x)$, we obtain

$$d^2(x)\beta[a, y]_\alpha = 0, \text{ for all } x, y \in M \text{ and } \alpha, \beta \in \Gamma.$$

Replacing y by $z\gamma y$, we get

$$d^2(x)\beta z\gamma[a, y]_\alpha = 0, \text{ for all } x, y, z \in M \text{ and } \alpha, \beta, \gamma \in \Gamma.$$

By primeness we get $d^2(x) = 0$ or $[a, y]_\alpha = 0$, since $d \neq 0$, therefore $a \in Z$. ■

Theorem 7. *Let U be a nonzero Lie ideals of a 2-torsion free prime Γ -ring M and d a nonzero derivation of M . If $d(U) \subset Z$, then $U \subset Z$.*

Proof. suppose that $U \not\subseteq Z$, then by Lemma 5 we have $V = [U, U] \not\subseteq Z$. Let $u, w \in U$, hence from

$$d([u, w]_\alpha) = [d(u), w]_\alpha + [u, d(w)]_\alpha = 0.$$

Since $d(u), d(w) \in Z$. It follows that $d(V) = 0$.

Let $v \in V, m \in M$ and $\alpha \in \Gamma$, since $d(v) = 0$ and $d([v, m]_\alpha) = 0$, we get

$$[v, d(m)]_\alpha = 0, \text{ for all } v \in V, m \in M \text{ and } \alpha \in \Gamma.$$

Therefore by Lemma 6 we get $v \in Z$, contradiction. Accordingly, $U \subset Z$. ■

Lemma 8. Let $U \not\subseteq Z$ be a Lie ideal of 2-torsion free prime Γ -ring M and d a nonzero derivation of M . If $a \in M$ and $a\Gamma d(U) = 0$ ($d(U)\Gamma a = 0$), then $a = 0$.

Proof. For all $u \in U, x \in M$ and $\alpha \in \Gamma$ we have $[u, x]_\beta \gamma u \in U$. By hypothesis we have

$$\begin{aligned} 0 &= a\alpha d([u, x]_\beta \gamma u) \\ &= a\alpha [u, x]_\beta \gamma d(u), \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta, \gamma \in \Gamma. \end{aligned}$$

Replacing x by $d(v)\lambda x$, we get

$$a\alpha u\beta d(v)\lambda x\gamma d(u) = 0, \text{ for all } u, v \in U, x \in M \text{ and } \alpha, \beta, \gamma, \lambda \in \Gamma.$$

By primeness we obtain $a\alpha u\beta d(v) = 0$ or $d(u) = 0$.

Now let $K = \{u \in U \mid a\alpha u\beta d(v) = 0\}$ and $L = \{u \in U \mid d(u) = 0\}$. Since K and L are additive subgroups of U and $U = K \cup L$, but a group can't be union of its two proper subgroups and hence $U = K$ or $U = L$.

According to Theorem 7, $d(U) \neq 0$, which proves that $U = K$. Hence we get $a\Gamma U\Gamma d(v) = 0$, for all $v \in U$. By Lemma 3 we get $a = 0$ or $d(v) = 0$, again by Theorem 7 $d(U) \neq 0$, therefore $a = 0$. ■

Theorem 9. Let M be a 2-torsion free prime Γ -ring, U be a nonzero Lie ideal of M and d be a nonzero derivation of M . If $d^2(U) = 0$ and $d(U) \subset U$, then $U \subset Z$.

Proof. Suppose that $U \not\subseteq Z$, for all $x \in M, u \in U$ and $\alpha \in \Gamma$ we have $[x, u]_\alpha \in U$. Since $d^2(U) = 0$, then by using Remark 1 we get

$$\begin{aligned} 0 &= d^2([x\beta u, u]_\alpha) \\ &= d^2([x, u]_\alpha)\beta u + 2d([x, u]_\alpha)\beta d(u) + [x, u]_\alpha\beta d^2(u). \end{aligned}$$

Since M is 2-torsion free and $d^2(U) = 0$, then we get

$$d([x, u]_\alpha)\beta d(u) = 0, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Replacing u by $u + d(u)$, we get $d([x, d(u)]_\alpha)\beta d(u) = 0$, so that

$$[d(x), d(u)]_\alpha\beta d(u) = 0, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

According to Lemma 8 we get $[d(x), d(u)]_\alpha = 0$ for all $x \in M, w \in U$ and $\alpha \in \Gamma$, therefor by Lemma 6 we conclude that $d(U) \subset Z$, which is contradicts Theorem 7, this prove the theorem. \blacksquare

Lemma 10. *Let M be a 2 and 3-torsion free prime Γ -ring, U be a nonzero Lie ideal of M and d be a nonzero derivation of M . If $d(U) \subset U$, $d^2(U) \subset Z$ and $d^3(U) = 0$ then $U \subset Z$.*

Proof. For all $x \in M, u \in U$ and $\alpha \in \Gamma$ we have $[x, u]_\alpha \in U$. Since $d^3(U) = 0$, then we obtain $d^3([x, u]_\alpha) = 0$. Replacing x by $x\beta u$ and using Remark 1(iv) we get

$$\begin{aligned} 0 &= d^3([x\beta u, u]_\alpha) \\ &= 3d^2([x, u]_\alpha)\beta d(u) + 3d([x, u]_\alpha)\beta d^2(u). \end{aligned}$$

Since M is 3-torsion free, then we get

$$d^2([x, u]_\alpha)\beta d(u) + d([x, u]_\alpha)\beta d^3(u) = 0, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Replacing u by $d(u)$ and using $d^2(U) = 0$ we obtain

$$d^2([x, d(u)]_\alpha)\beta d^2(u) = 0.$$

Since $d^2(U) \subset Z$, then by Lemma 2(i) we get

$$(3) \quad d^2([x, d(u)]_\alpha) = 0 \quad \text{or} \quad d^2(u) = 0.$$

If $d^2([x, d(u)]_\alpha) = 0$, then replacing x by $x\beta d(u)$ we obtain

$$\begin{aligned} 0 &= d^2([x\beta d(u), d(u)]_\alpha) \\ &= d^2([x, d(u)]_\alpha)\beta d(u) \\ &= d^2([x, d(u)]_\alpha)\beta d(u) + 2d([x, d(u)]_\alpha)\beta d^2(u) + [x, d(u)]_\alpha\beta d^3(u). \end{aligned}$$

Since $d^3(U) = 0$, M is a 2-torsion free and by relation (3), then the last equation reduced to

$$d([x, d(u)]_\alpha \beta d^2(u)) = 0, \text{ for all } x \in M, u \in U \text{ and } \alpha, \beta \in \Gamma.$$

Since $d^2(U) \subset Z$, then by Lemma 2(i) we get $d([x, d(u)]_\alpha) = 0$ or $d^2(u) = 0$.

If $d([x, d(u)]_\alpha) = 0$, then replacing x by $x\gamma d(u)$, we obtain

$$\begin{aligned} 0 &= d([x\gamma d(u), d(u)]_\alpha) \\ &= d([x, d(u)]_\alpha \gamma d(u)) \\ &= d([x, d(u)]_\alpha) \gamma d(u) + [x, d(u)]_\alpha \gamma d^2(u) \\ &= [x, d(u)]_\alpha \beta d^2(u). \end{aligned}$$

Since $d^2(U) \subset Z$ and $d(U) \subset U$, then by Lemma 2(i) we get $[x, d(u)]_\alpha = 0$ or $d^2(u) = 0$. If $[x, d(u)]_\alpha = 0$, then we have $d(u) \subset Z$. Hence from relation (3) we have either $d(u) \subset Z$ or $d^2(u) = 0$.

Now let $K = \{u \in U \mid d(u) \subset Z\}$ and $L = \{u \in U \mid d^2(u) = 0\}$. Since K and L are additive subgroups of U and $U = K \cup L$, but a group can't be union of its two proper subgroups and hence $U = K$ or $U = L$. If $U = K$, that is $d(u) \subset Z$, then by Theorem 7 we get $U \subset Z$, or $U = L$, that is $d^2(u) = 0$, hence by Theorem 9 we get $U \subset Z$. ■

Theorem 11. *Let M be a 2 and 3-torsion free prime Γ -ring, U be a nonzero Lie ideal of M and d be a nonzero derivation of M . If $d(U) \subset U$ and $d^2(U) \subset Z$, then $U \subset Z$.*

Proof. For all $x \in M$, $u \in U$ and $\alpha \in \Gamma$ we have

$$(4) \quad d^2([x, u]_\alpha) \in Z.$$

Replacing x by $x\beta d^2(v)$, where $v \in U$ and $\beta \in \Gamma$, and using $d^2(U) \subset Z$, we get

$$(5) \quad 2d([x, u]_\alpha) \beta d^3(v) + [x, u]_\alpha \beta d^4(v) \subset Z, \text{ for all } u, v \in U, x \in M \text{ and } \alpha, \beta \in \Gamma.$$

Replacing x by $x\gamma d^2(w)$ in relation (5), where $w \in U$ and $\gamma \in \Gamma$, and using $d^2(U) \subset Z$ and M is 2-torsion free, then the relation (5) reduced to

$$[x, u]_\alpha \gamma d^3(w) \beta d^3(v) \in Z, \text{ for all } v, u, w \in U, x \in M \text{ and } \alpha, \beta, \gamma \in \Gamma.$$

Since $d^2(U) \subset Z$ and $d(U) \subset U$, then $d^3(U) \subset Z$ and thus by Lemma 2(ii) we have $d^3(U) = 0$ or $[x, u]_\alpha \subset Z$. Therefore if $d^3(U) = 0$, hence by Lemma 10 yields $U \subset Z$. If $[M, U]_\alpha \subset Z$, then by Lemma 4 we get $U \subset Z$. ■

REFERENCES

- [1] W. E. Barnes, *On the Γ -Rings of Nobusawa*, Pacific J. Math. **18** (1966) 411–422. doi:10.2140/pjm.1966.18.411
- [2] J. Bergan, I. N. Herstein, and W. Kerr, *Lie Ideals and Derivations of Prime Rings*, J. Algebra **71** (1981) 259–267. doi:10.1016/0021-8693(81)90120-4
- [3] A. K. Halder and A. C. Paul, *Jordan Left Derivations on Lie Ideals of Prime Γ -Rings*, Punjab Univ. J. of Math. (2011) 1–7.
- [4] P.H.Lee and T.K.Lee, *Lie Ideals of Prime Rings with Derivations*, Bull. Inst. Math. Acad. Scin. **11** (1983) 75–80.
- [5] N. Nobusawa, *On a Generalization of the Ring Theory*, Osaka J. Math. **1** (1964) 81–89.
- [6] M. Soyuturk, *The Commutativity in Prime Gamma Rings with Derivation*, Tr. J. Math. **18** (1994) 149–155.

Received 24 October 2012

Revised 3 January 2013