

## JORDAN NUMBERS, STIRLING NUMBERS AND SUMS OF POWERS

ROMAN WITUŁA, KONRAD KACZMAREK, PIOTR LORENC,

EDYTA HETMANIOK AND MARIUSZ PLESZCZYŃSKI

*Institute of Mathematics*  
*Silesian University of Technology*  
*Kaszubska 23, 44-100 Gliwice, Poland*

**e-mail:** roman.witula,konrad.kaczmarek,edyta.hetmaniok@polsl.pl

### Abstract

In the paper a new combinatorical interpretation of the Jordan numbers is presented. Binomial type formulae connecting both kinds of numbers mentioned in the title are given. The decomposition of the product of polynomial of variable  $n$  into the sums of  $k$ th powers of consecutive integers from 1 to  $n$  is also studied.

**Keywords:** Bernoulli numbers, binomial coefficients, Jordan numbers, Stirling numbers, Živković numbers.

**2010 Mathematics Subject Classification:** 11B68, 11B73, 11B83.

### 1. INTRODUCTION

This paper was inspired by the wish to generalize the elementary formulae (formulae (2)–(7) given below) generated by the authors in the course of discussion on D. Knuth's excellent publication [9] (devoted to the analysis of relations between sums of the powers of consecutive positive integers, inspired by Faulhaber's "old" and "deeper" results). It led us first to find the general connection of type (8) (known form) and (9) (probably new form). In the course of discussing these relations a very interesting and original problem has appeared, which consisted in deriving a formula of type (11), meaning a decomposition of the product  $n^k \sum_{l=1}^n l^r$  into the sum of type  $\sum_{s=1}^{k+r} a_s(k, r) \sum_{l=1}^n l^s$ . Section 3 is devoted to this problem. Section 4 presents the general formula (26) (known formula) giving the decomposition of the Stirling numbers of second kind in the linear combination of binomial coefficients by using the Jordan numbers. Finally, in Section 5 the new combinatoric interpretation of the Jordan numbers is presented.

## 2. STIRLING NUMBERS

It should be reminded that Stirling numbers of the second kind or the partition numbers  $S(n, k)$ ,  $k, n \in \mathbb{N}$ ,  $k \leq n$ , satisfy the triangular recurrence relation [12, 13, 14]:

$$(1) \quad \begin{aligned} S(n, k) &= S(n-1, k-1) + k S(n-1, k), \\ S(k, k) &= 1, \quad S(k, 0) = 0. \end{aligned}$$

Moreover, we adopt here Donald Knuth's notation [6, 9]:

$$\sum f(n) := \sum_{l=1}^n f(l)$$

for every function  $f : \mathbb{N} \rightarrow \mathbb{C}$ . For example we have

$$\sum n^k := \sum_{l=1}^n l^k, \quad k, n \in \mathbb{N}.$$

We shall start our deliberations by presenting the above-mentioned basic identities which can be easily verified by direct calculations. So, the following identities hold:

$$(2) \quad \sum n = S(n+1, n),$$

$$(3) \quad \begin{aligned} \sum n^3 + \sum n^2 &= \sum n^2(n+1) = \frac{1}{2}(3n+1) \binom{n+2}{3} \\ &= \frac{1}{2}(3(n-1)+4) \binom{n+2}{3} \\ &= 6 \binom{n+2}{4} + 2 \binom{n+2}{3} = 2S(n+2, n), \end{aligned}$$

$$(4) \quad \begin{aligned} 3 \sum n^5 + 10 \sum n^4 + 9 \sum n^3 + 2 \sum n^2 &= \sum n^2(n+1)(n+2)(3n+1) \\ &= 24 \binom{n+3}{4} \binom{n+1}{2} = 24S(n+3, n), \end{aligned}$$

$$(5) \quad S(n+3, n) = 15 \binom{n+3}{6} + 10 \binom{n+3}{5} + \binom{n+3}{4},$$

$$\begin{aligned} \sum n^7 + 7 \sum n^6 + 17 \sum n^5 + 17 \sum n^4 + 6 \sum n^3 &= \sum n^3(n+1)^2(n+2)(n+3) \\ (6) \qquad \qquad \qquad &= 48 S(n+4, n) = (15n^3 + 30n^2 + 5n - 2) \binom{n+4}{5}, \end{aligned}$$

$$(7) \quad S(n+4, n) = 105 \binom{n+4}{8} + 105 \binom{n+4}{7} + 25 \binom{n+4}{6} + \binom{n+4}{5}.$$

Stirling numbers of the second kind are presented in equations (3), (5) and (7) which is not surprising because the following inversion formulae hold.

**Proposition 1.** *We have*

$$(8) \qquad \qquad \qquad \sum n^r = \sum_{l=0}^r S(r, l) l! \binom{n+1}{l+1}$$

$$(9) \qquad \qquad \qquad = \sum_{l=1}^{r+1} S(r+1, l) (l-1)! \binom{n}{l}.$$

**Proof.** Formula (8) is drawn from [9], whereas the second formula seems to be new, yet it may be easily derived from the first one:

$$\begin{aligned} \sum_{l=0}^r S(r, l) l! \binom{n+1}{l+1} &= \sum_{l=0}^r S(r, l) l! \left( \binom{n}{l+1} + \binom{n}{l} \right) \\ &= S(r, 0) + \sum_{l=0}^{r-1} \binom{n}{l+1} \left( S(r, l) l! + S(r, l+1) (l+1)! \right) + S(r, r) r! \binom{n}{r+1} \\ &= \sum_{l=0}^{r-1} \binom{n}{l+1} l! \left( S(r, l) + S(r, l+1) (l+1) \right) + S(r+1, r+1) r! \binom{n}{r+1} \\ &= \sum_{l=1}^{r+1} \binom{n}{l} (l-1)! S(r+1, l). \end{aligned}$$

■

### 3. DECOMPOSITION OF PRODUCTS $n^k \sum n^m$

We note that from (9) we obtain

$$\begin{aligned} (n+1) \sum n^r &= \sum_{l=1}^{r+1} S(r+1, l) (l-1)! (l+1) \binom{n+1}{l+1} \\ (10) \qquad \qquad &\stackrel{\text{by (8)}}{=} \sum n^{r+1} + \sum_{l=1}^{r+1} S(r+1, l) (l-1)! \binom{n+1}{l+1}, \end{aligned}$$

which, again by (8), suggests a formula of following form

$$(11) \quad n^k \sum n^r = \sum_{s=1}^{k+r} a_s(k, r) \sum n^{k+r+1-s}.$$

Since we have (see [6]):

$$(12) \quad \begin{aligned} (r+1) \sum n^r &= \sum_{l=0}^r (-1)^l \binom{r+1}{l} B_l n^{r+1-l} \iff \\ n^{r+1} &= (r+1) \sum n^r + (r+1) B_1 n^r - \sum_{k=1}^{\lfloor r/2 \rfloor} \binom{r+1}{2k} B_{2k} n^{r+1-2k}, \end{aligned}$$

where  $B_l$  are the Bernoulli's numbers, we obtain indeed formulae of the type (11). From (12) the following formulae can be generated.

**Theorem 2.** *We have*

$$(13) \quad (r+1)n \sum n^r = (r+2) \sum n^{r+1} + \sum_{s=1}^r \binom{r+1}{s} B_s \sum n^{r-s+1},$$

$$(14) \quad \begin{aligned} (r+1)n^2 \sum n^r &= (r+2)n \sum n^{r+1} + \sum_{s=1}^r \binom{r+1}{s} B_s n \sum n^{r-s+1} \\ &= (r+3) \sum n^{r+2} + \sum_{p=1}^{r+1} \binom{r+2}{p} B_p \sum n^{r-p+2} \\ &+ \sum_{s=1}^r \binom{r+1}{s} \frac{B_s}{r-s+2} \left( (r-s+3) \sum n^{r-s+2} \right. \\ &+ \left. \sum_{q=1}^{r-s+1} \binom{r-s+2}{q} B_q \sum n^{r-s-q+2} \right) \\ &= (r+3) \sum n^{r+2} + \sum_{p=1}^r \frac{2r-p+5}{r+2} \binom{r+2}{p} B_p \sum n^{r-p+2} \\ &+ (r+2) B_{r+1} \sum n + \sum_{s=1}^r \sum_{q=1}^{r-s+1} \binom{r+1}{s} \binom{r-s+2}{q} \frac{B_s B_q}{r-s+2} \sum n^{r-s-q+2}. \end{aligned}$$

Hence, after the appropriate regrouping, we get

$$\begin{aligned}
 (15) \quad & (r+1)n^2 \sum n^r = (r+3) \sum n^{r+2} - (r+2) \sum n^{r+1} + \frac{1}{6}(r+1)(r+3) \sum n^r \\
 & - \frac{1}{6} \binom{r+1}{2} \sum n^{r-1} - \frac{1}{15} \binom{r+1}{4} \sum n^{r-2} + \frac{1}{30} \binom{r+1}{4} \sum n^{r-3} \\
 & + \frac{1}{21} \binom{r+1}{6} \sum n^{r-4} + \dots \\
 & + \frac{1}{2} \left[ (r+5)(r+1)B_r + \sum_{s=1}^{r-1} \binom{r+1}{s} (r-s+1)B_s B_{r-s} \right] \sum n^2 \\
 & + \left( (r+2)B_{r+1} + \sum_{s=1}^r \binom{r+1}{s} B_s B_{r-s+1} \right) \sum n.
 \end{aligned}$$

The last two coefficients can be reduced to the following ones (the classical splitting formulae are applied here [1, 5, 7, 10, 11]):

$$\frac{1}{2} \left[ (r+5)(r+1)B_r - \left( 2 \left\lfloor \frac{r}{2} \right\rfloor + 1 \right) (r+1)B_{2\lfloor \frac{r}{2} \rfloor} \right]$$

since we have the equality

$$\sum_{s=1}^{r-1} \binom{r+1}{s} (r-s+1)B_s B_{r-s} = - \left( 2 \left\lfloor \frac{r}{2} \right\rfloor + 1 \right) (r+1)B_{2\lfloor \frac{r}{2} \rfloor}$$

for every  $r = 3, 4, \dots$  (we have  $(2 \lfloor \frac{r}{2} \rfloor + 1) (r+1)B_{2\lfloor \frac{r}{2} \rfloor}$  for  $r = 2$ ), and

$$(r+2)B_{r+1} + \sum_{s=1}^r \binom{r+1}{s} B_s B_{r-s+1} = -(r+1)B_r.$$

From these relations, we obtain

$$\sum_{s=1}^{r-1} \binom{r+1}{s} (r-s+1)B_s B_{r-s} = (r+1) \sum_{s=1}^{r-1} \binom{r}{s} B_s B_{r-s}$$

if  $r$  is odd and  $r \geq 3$ .

**Proof.** We prove only (13). To this aim let us fix  $r \in \mathbb{N}$  and put

$$(16) \quad T_{k,r} := (r+1)k^r - (k+1)^{r+1} + k^{r+1}.$$

We have

$$\begin{aligned}
 T_{k,r} &= (r+1)k^r + k^{r+1} - \sum_{i=0}^{r+1} \binom{r+1}{i} k^i = - \sum_{i=0}^{r-1} \binom{r+1}{i} k^i, \\
 (17) \quad \sum_{k=1}^n T_{k,r} &= (r+1) \sum n^r + \sum_{k=1}^n (k^{r+1} - (k+1)^{r+1}) \\
 &= (r+1) \sum n^r + 1 - (n+1)^{r+1}.
 \end{aligned}$$

On the other hand

$$(18) \quad \sum_{k=1}^n T_{k,r} = - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i.$$

We note that (13) holds for  $n = 1$  since it is equivalent to the following known relation

$$B_{r+1} = \sum_{s=0}^{r+1} \binom{r+1}{s} B_s.$$

Let us assume that (13) holds for some  $n \in \mathbb{N}$ . Then we get

$$\begin{aligned}
 (19) \quad & (r+1)n \sum n^r - (r+2) \sum n^{r+1} - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i \\
 &= \sum_{s=1}^r \binom{r+1}{s} B_s \sum n^{r-s+1} - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i \\
 &= \sum_{s=1}^r \binom{r+1}{s} B_s \sum (n+1)^{r-s+1} \\
 &\quad - \sum_{s=1}^r \binom{r+1}{s} B_s (n+1)^{r-s+1} - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i,
 \end{aligned}$$

and

$$\begin{aligned}
 & - \sum_{s=1}^r \binom{r+1}{s} B_s (n+1)^{r-s+1} \\
 &= (n+1)^{r+1} - 2(r+1)B_1(n+1)^r - \sum_{s=0}^r (-1)^s \binom{r+1}{s} B_s (n+1)^{r-s+1} \\
 &\stackrel{(12)}{=} (n+1)^{r+1} + (r+1)(n+1)^r - (r+1) \sum (n+1)^r \\
 &= (n+1)^{r+1} - (r+1) \sum n^r,
 \end{aligned}$$

which implies

$$\begin{aligned}
 (20) \quad & - \sum_{s=1}^r \binom{r+1}{s} B_s (n+1)^{r-s+1} - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i \\
 & = (n+1)^{r+1} - \sum_{i=0}^r \binom{r+1}{i} \sum n^i = 1.
 \end{aligned}$$

We note that the last identity can be easily deduced after summing the following equalities

$$\begin{aligned}
 (n+1)^{r+1} &= n^{r+1} + \sum_{i=0}^r \binom{r+1}{i} n^i, \\
 n^{r+1} &= (n-1)^{r+1} + \sum_{i=0}^r \binom{r+1}{i} (n-1)^i, \\
 &\dots \\
 2^{r+1} &= 1^{r+1} + \sum_{i=0}^r \binom{r+1}{i} 1^i.
 \end{aligned}$$

Moreover, by (17) and (18), we obtain

$$\begin{aligned}
 (21) \quad & (r+1)n \sum n^r - (r+2) \sum n^{r+1} - \sum_{i=0}^{r-1} \binom{r+1}{i} \sum n^i \\
 & = (r+1)n \sum n^r - (r+2) \sum n^{r+1} + (r+1) \sum n^r + 1 - (n+1)^{r+1} \\
 & = 1 + (r+1)(n+1) \sum (n+1)^r - (r+2) \sum (n+1)^{r+1}.
 \end{aligned}$$

By comparing (19)–(21) we conclude that relation (16) holds also for  $n$  replaced by  $n+1$  which, by the principle of mathematical induction, ends the proof. ■

**Corollary 3.** *From (15) the following special formulae can be deduced*

$$\begin{aligned}
 2n^2 \sum n &= 4 \sum n^3 - 3 \sum n^2 + \sum n, \\
 3n^2 \sum n^2 &= 5 \sum n^4 - 4 \sum n^2 + \frac{5}{2} \sum n^2 - \frac{1}{2} \sum n, \\
 4n^2 \sum n^3 &= 6 \sum n^5 - 5 \sum n^4 + 4 \sum n^3 - \sum n^2.
 \end{aligned}$$

Some more general formula, than the one in (15), can be also obtained.

**Proposition 4.** *We have*

$$\begin{aligned}
 (22) \quad (m+1)n^k \sum n^m &= (m+k+1) \sum n^{m+k} - \frac{k}{2}(m+k) \sum n^{m+k-1} \\
 &+ \frac{k}{12}(m+2k-1)(m+k-1) \sum n^{m+k-2} \\
 &- \frac{k}{24}(k-1)(m+k-1)(m+k-2) \sum n^{m+k-3} \\
 &+ \frac{1}{720}(m+k-3)k(m+2k-3)(3k^2+3k(m-3)-m(m+4)+6) \sum n^{m+k-4} \\
 &- \frac{1}{360} \binom{m+k-3}{2} \binom{k}{2} (2k^2+2k(m-4)-m(m+5)+8) \sum n^{m+k-5} + \dots \\
 &+ \text{coeff}(m, k) \sum n.
 \end{aligned}$$

We have found the general formula only for the first six coefficients from above. The last absent coefficient  $\text{coeff}(m, k)$  can be found by subtracting from the left side of (22) the expression staying on the right side of this formula with sums  $\sum n^p$  where  $p \geq 2$ .

For example, we find

$$\begin{aligned}
 2n \sum n &= 3 \sum n^2 - \sum n, \\
 2n^3 \sum n &= 5 \sum n^4 - 6 \sum n^3 + \frac{9}{2} \sum n^2 - \frac{7}{2} \sum n, \\
 4n \sum n^3 &= 5 \sum n^4 - 2 \sum n^3 + \sum n^2, \\
 3n^4 \sum n^2 &= \frac{1}{2} n^5 (n+1)(2n+1) \\
 &= 7 \sum n^6 - 12 \sum n^5 + 15 \sum n^4 - 10 \sum n^3 + \frac{7}{2} \sum n^2 - \frac{1}{2} \sum n,
 \end{aligned}$$

and the following special one

$$\begin{aligned}
 (r+1)n^3 \sum n^r &= (r+4) \sum n^{r+3} - \frac{3}{2}(r+3) \sum n^{r+2} + \frac{1}{4}(r+2)(r+5) \sum n^{r+1} \\
 &- \frac{1}{4}(r+1)(r+2) \sum n^r - \frac{1}{240}(r-6)r(r+1)(r+3) \sum n^{r-1} \\
 &+ \frac{1}{240}(r-2)(r-1)r(r+1) \sum n^{r-2} + \dots
 \end{aligned}$$

#### 4. JORDAN NUMBERS

In this section, we present the generalization of relations (3), (5) and (7).



**Proposition 5.** *We have*

$$(23) \quad S(n+k, n) = \sum_{l=0}^{k-1} a_{l,k} \binom{n+k}{2k-l},$$

where

$$(24) \quad \begin{aligned} a_{0,k+1} &= (2k+1)a_{0,k}, & a_{k,k+1} &= a_{k-1,k}, \\ a_{l,k+1} &= (k-l+1)a_{l-1,k} + (2k-l+1)a_{l,k}, \end{aligned}$$

for  $l = 1, 2, \dots, k-1$ . The numbers  $a_{l,k}$  are called the Jordan numbers [2, 3, 4, 8, 15]. Authors of the present paper obtained the above relations independently. For the sake of selfcontainedness, the proof of (23) will be given now. We proceed by induction.

**Proof.** (23) We shall employ the following basic formula

$$(25) \quad \begin{aligned} \sum_{k=m-l}^n k \binom{k+l}{m} &= \sum_{k=m-l}^n ((k+l-m) + (m-l)) \binom{k+l}{m} \\ &= (m+1) \sum_{k=m-l+1}^n \binom{k+l}{m+1} + (m-l) \sum_{k=m-l}^n \binom{k+l}{m} \\ &= (m+1) \binom{n+l+1}{m+2} + (m-l) \binom{n+l+1}{m+1}. \end{aligned}$$

Thus, assuming that formula (23) holds for certain  $k, n \in \mathbb{N}$ , from (25) and (1) the following relation can be derived

$$\begin{aligned} S(n+k+1, n) &= S(n+k+1, n) - S(k+2, 0) \\ &= \sum_{l=0}^{n-1} (S(l+k+2, l+1) - S(l+k+1, l)) = \sum_{l=0}^{n-1} (l+1) S(l+k+1, l+1) \\ &= \sum_{l=0}^{n-1} (l+1) \sum_{\tau=0}^{k-1} a_{\tau,k} \binom{l+k+1}{2k-\tau} = \sum_{\tau=0}^{k-1} a_{\tau,k} \sum_{l=k-1-\tau}^{n-1} (l+1) \binom{l+k+1}{2k-\tau} \\ &= \sum_{\tau=0}^{k-1} a_{\tau,k} \left( (2k-\tau+1) \binom{n+k+1}{2k-\tau+2} + (k-\tau) \binom{n+k+1}{2k-\tau+1} \right) \\ &= a_{0,k} (2k+1) \binom{n+k+1}{2k+2} + \sum_{\tau=0}^{k-2} (a_{\tau,k} (k-\tau) + a_{\tau+1,k} (2k-\tau)) \binom{n+k+1}{2k-\tau+1} \\ &+ a_{k-1,k} \binom{n+k+1}{k+2} = \sum_{l=0}^k a_{l,k+1} \binom{n+k+1}{2k-l+2}. \end{aligned}$$

■

**Remark 6.** It is possible to prove the following formulae:

$$\begin{aligned} a_{0,k} &= (2k-1)!!, & a_{1,k} &= \frac{k-1}{3} a_{0,k}, \\ a_{2,k} &= \frac{1}{12} \binom{2k-2}{3} (2k-3)!!, & a_{3,k} &= S2(k, 2k+3), \\ a_{k-2,k} &= 2^{k+1} - k - 3 & a_{k-1,k} &= 1, \end{aligned}$$

where  $S2(n, k)$  denotes the 2-associated Stirling number of the second kind (see [12] and sequences A000478 and A000247 in [14]).

In Table 1 the triangle of coefficients  $a_{l,k}$ ,  $l = 0, 1, \dots, k-1$  is presented.

Table 1. Triangle of coefficients  $a_{l,k}$

$l, k$	1	2	3	4	5	6	7	8
0	1	3	15	105	945	10395	135135	2027025
1		1	10	105	1260	17325	270270	4729725
2			1	25	490	9450	190575	4099095
3				1	56	1918	56980	1636635
4					1	119	6825	302995
5						1	246	22935
6							1	501
7								1

**Remark 7.** Another formula connecting Stirling numbers of the second kind with binomial coefficients is also known

$$S(n, n-k) = \sum_r \left\langle\left\langle \begin{matrix} k \\ r \end{matrix} \right\rangle\right\rangle \binom{n+k-1-r}{2k},$$

where  $\left\langle\left\langle \begin{matrix} k \\ r \end{matrix} \right\rangle\right\rangle$  are the Eulerian numbers of the second kind (see [6]).

## 5. COMBINATORIC INTERPRETATION OF THE JORDAN AND ŽIVKOVIĆ NUMBERS

As it can be inferred from [12, pp. 76–77], the numbers  $a_{n-2k-1, n-k}$ , for  $k = 0, 1, \dots, \lfloor (n-1)/2 \rfloor$ , enumerate the permutations of  $n$  elements with  $k$  cycles,

none of which is a unit cycle. We note that in [12] the numbers

$$b(n, k) := a_{n-2k-1, n-k}$$

are called the associated Stirling number of the second kind. Moreover, we have the following recurrence relation

$$b(n + 1, k) = k b(n, k) + n b(n - 1, k - 1).$$

Conversely, in [15] the Živković numbers  $G(k, i)$  are defined by the following equality

$$s(n, n - k) = \sum_{i=0}^k (-1)^i G(k, i) \binom{n + i - 1}{k + i},$$

where  $G(k, i) = 0$  if  $k \geq 1$  and  $i > k$  or  $i < 1$  ( $G(0, i) = \delta_{0,i}$ ), and  $s(n, k)$  denotes the Stirling number of the first kind.

Numbers  $G(k, i)$  are determined by the initial condition  $G(1, i) = \delta_{1,i}$  and by the recurrence relation

$$(26) \quad G(k + 1, i) = i G(k, i) + (k + i) G(k, i - 1).$$

It is easy to check that

$$(27) \quad G(r - l, r) = a_{l,r} \iff G(k, k + l) = a_{l,k+l} \quad (r - l := k).$$

Hence, as it is verifiable on the grounds of relation (26),  $a_{l,k+l}$  denotes the number of ways of placing  $2k + l$  labeled balls into  $k$  indistinguishable boxes with at least two balls in each box.

The values of numbers  $G(k, i)$  are compiled in Table 2 (we note that  $G(k, i)$ ,  $k = i, i + 1, \dots$  make up  $i$ -th diagonal – Table 1 – for each  $i \in \mathbb{N}$ ; the diagonal is from the main line upwards).

Table 2. Values of numbers  $G(k, i)$

$G(1, k)$	$G(2, k)$	$G(3, k)$	$G(4, k)$	$G(5, k)$	$G(6, k)$
1					
1	3				
1	10	15			
1	25	105	105		
1	56	490	1260	945	
1	119	1918	9450	17325	10395

## REFERENCES

- [1] Z.I. Borevich and I.R. Szafarevich, *Number Theory* (Nauka, Moscow, 1964, in Russian).
- [2] L. Carlitz, *Note on the numbers of Jordan and Ward*, *Duke Math. J.* **38** (1971) 783–790. doi:10.1215/S0012-7094-71-03894-4
- [3] L. Carlitz, *Some numbers related to the Stirling numbers of the first and second kind*, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **544–576** (1976) 49–55.
- [4] L. Carlitz, *Some remarks on the Stirling numbers*, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **678–715** (1980) 10–14.
- [5] K. Dilcher, *Bernoulli and Euler Polynomials*, 587–600 (in F.W.I. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark, *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press, 2010).
- [6] R.L. Graham, D.E. Knuth and O. Patashnik, *Concrete Mathematics* (Addison-Wesley, Reading, 1994).
- [7] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory* (Springer, 1990). doi:10.1007/978-1-4757-1779-2
- [8] C. Jordan, *Calculus of Finite Differences* (Chelsea, New York, 1960). doi:10.2307/2333783
- [9] D.E. Knuth, *Johann Faulhaber and sums of powers*, *Math. Comp.* **203** (1993) 277–294. doi:10.2307/2152953
- [10] N. Nielsen, *Traité élémentaire des nombres de Bernoulli* (Gauthier – Villars, Paris, 1923).
- [11] S. Rabsztyn, D. Słota and R. Wituła, *Gamma and Beta Functions, Part I* (Silesian Technical University Press, Gliwice, 2011, in Polish).
- [12] J. Riordan, *An Introduction to Combinatorial Analysis* (John Wiley, 1958). doi:10.1063/1.3060724
- [13] J. Riordan, *Combinatorial Identities* (Wiley, New York, 1968).
- [14] N.J.A. Sloane, *The On-Line Encyclopedia of Integer Sequences* (<http://oeis.org/>).
- [15] M. Živković, *On a representation of Stirling's numbers of first kind*, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **498–541** (1975) 217–221.

Received 6 December 2013

Revised 6 January 2015