

A SHORT NOTE ON L_{CBA} —FUZZY LOGIC WITH A NON-ASSOCIATIVE CONJUNCTION

MIROSLAV KOLAŘÍK

Palacký University Olomouc, Faculty of Science
Department of Computer Science
Třída 17. listopadu 12, 77146 Olomouc, Czech Republic
e-mail: miroslav.kolarik@upol.cz

Abstract

We significantly simplify the axiomatic system L_{CBA} for fuzzy logic with a non-associative conjunction.

Keywords: axiomatic system, non-associativity, fuzzy logic.

2010 Mathematics Subject Classification: 06D35, 06F35, 03G10.

Several investigations in probability theory and the theory of expert systems show that it is important to search for some reasonable generalizations of fuzzy logics having a non-associative conjunction, see [5, 6, 8, 9, 10].

In the paper [1] Botur and Halaš introduced and described a non-associative fuzzy logic L_{CBA} having as an equivalent algebraic semantics lattices with section antitone involutions satisfying the contraposition law, so-called commutative basic algebras. The variety of commutative basic algebras was intensively studied in several recent papers and includes the class of MV-algebras. For more details see the book [2].

In [1] Botur and Halaš introduced a non-associative fuzzy logic L_{CBA} by the following nine axioms:

- (i) $x \rightarrow (y \rightarrow x) = 1$
- (ii) $((x \rightarrow y) \rightarrow y) \rightarrow ((y \rightarrow x) \rightarrow x) = 1$
- (iii) $(\neg x \rightarrow \neg y) \rightarrow (y \rightarrow x) = 1$
- (iv) $\neg\neg x \rightarrow x = 1$
- (v) $(x \rightarrow y) \rightarrow (x \rightarrow \neg\neg y) = 1$
- (vi) $x \rightarrow x = 1$
- (vii) $(x = 1 \ \& \ x \rightarrow y = 1) \Rightarrow y = 1$

$$(viii) \quad x \rightarrow y = 1 \Rightarrow (y \rightarrow z) \rightarrow (x \rightarrow z) = 1$$

$$(ix) \quad (x \rightarrow y = 1 \ \& \ y \rightarrow x = 1) \Rightarrow x = y.$$

In the main theorem we significantly simplify the axiomatic system L_{CBA} for fuzzy logic with a non-associative conjunction. We show that three axioms can be omitted.

Theorem 1. *Identities (iv), (v), (vi) follows from identities (i), (ii), (iii) and from quasiidentities (vii), (viii), (ix).*

Proof. First, we show that (vi) is redundant. Using (viii) on (i) we easily obtain

$$(1) \quad ((y \rightarrow x) \rightarrow x) \rightarrow (x \rightarrow x) = 1.$$

Using (viii) on (ii) we obtain

$$(2) \quad (((y \rightarrow x) \rightarrow x) \rightarrow (x \rightarrow x)) \rightarrow (((x \rightarrow y) \rightarrow y) \rightarrow (x \rightarrow x)) = 1.$$

Applying (vii) on (1) and (2) we get

$$(3) \quad ((x \rightarrow y) \rightarrow y) \rightarrow (x \rightarrow x) = 1.$$

Further, using (viii) on (i) in the form $y \rightarrow ((x \rightarrow y) \rightarrow y) = 1$ we derive

$$(4) \quad (((x \rightarrow y) \rightarrow y) \rightarrow (x \rightarrow x)) \rightarrow (y \rightarrow (x \rightarrow x)) = 1.$$

Applying (vii) on (3) and (4) we get $y \rightarrow (x \rightarrow x) = 1$. From the last identity, where $y := 1$, using (vii), we obtain (vi).

Now, we show that (iv) is redundant. From (i), using (vi), we easily derive that

$$(5) \quad x \rightarrow 1 = 1.$$

Putting $y := 1$ in (ii) and applying (5) we get

$$1 \rightarrow ((1 \rightarrow x) \rightarrow x) = 1,$$

which, by (vii), give us

$$(6) \quad (1 \rightarrow x) \rightarrow x = 1.$$

From this and from (i) in the form $x \rightarrow (1 \rightarrow x) = 1$, we obtain

$$(7) \quad 1 \rightarrow x = x.$$

by (ix). From (iii) and (viii) we infer

$$(8) \quad ((y \rightarrow x) \rightarrow z) \rightarrow ((\neg x \rightarrow \neg y) \rightarrow z) = 1,$$

from (i) and (viii) we obtain

$$(9) \quad ((y \rightarrow x) \rightarrow z) \rightarrow (x \rightarrow z) = 1,$$

and from (ii) and (ix) we have

$$(10) \quad (x \rightarrow y) \rightarrow y = (y \rightarrow x) \rightarrow x.$$

Putting $y := 1$ in (iii) and applying (7) we get

$$(11) \quad (\neg x \rightarrow \neg 1) \rightarrow x = 1.$$

From this, for $x := \neg 1$, we get immediately $(\neg \neg 1 \rightarrow \neg 1) \rightarrow \neg 1 = 1$. From (i), where $x := \neg 1$ and $y := \neg \neg 1$, we obtain $\neg 1 \rightarrow (\neg \neg 1 \rightarrow \neg 1) = 1$. Applying (ix) to the last two equations we conclude

$$(12) \quad \neg \neg 1 \rightarrow \neg 1 = \neg 1.$$

Applying (vii) to (11) and to (9), where $y := \neg x$, $x := \neg 1$ and $z := x$, we derive

$$(13) \quad \neg 1 \rightarrow x = 1.$$

Now, put $\neg 1$ instead of x and $x \rightarrow \neg \neg 1$ instead of y in (10) to obtain

$$(\neg 1 \rightarrow (x \rightarrow \neg \neg 1)) \rightarrow (x \rightarrow \neg \neg 1) = ((x \rightarrow \neg \neg 1) \rightarrow \neg 1) \rightarrow \neg 1.$$

The left hand side of the last identity can be reduced to $x \rightarrow \neg \neg 1$, using (13) and (7). The right hand side of the last identity is equal to 1, by (9), where $y := x$, $x := \neg \neg 1$ and $z := \neg 1$, using (12). Therefore, we have

$$(14) \quad x \rightarrow \neg \neg 1 = 1.$$

From (14) we derive easily that $1 \rightarrow \neg \neg 1 = 1$ and from (5) we derive easily that $\neg \neg 1 \rightarrow 1 = 1$, whence, by (ix), we conclude

$$(15) \quad \neg \neg 1 = 1.$$

Putting $y := \neg x$, $x := \neg 1$ and $z := x$ in (8) and applying (11) and (vii) we get

$$(\neg \neg 1 \rightarrow \neg \neg x) \rightarrow x = 1$$

which, by (15) and (5), give us (iv).

Finally, we show that also (v) is redundant. Using (vii) on (iii) in the form $(\neg \neg \neg x \rightarrow \neg x) \rightarrow (x \rightarrow \neg \neg x) = 1$ and on (iv) in the form $\neg \neg \neg x \rightarrow x = 1$ we get $x \rightarrow \neg \neg x = 1$. This with (iv) give us by (ix)

$$x = \neg \neg x.$$

The last identity together with (vi) in the form $(x \rightarrow y) \rightarrow (x \rightarrow y) = 1$ proves (v). ■

Among the recent results on simplification of axiomatic systems related to fuzzy logics belong works of Cintula (based on a proof, see [3]) and Lehmke (applying a solver he programmed, see [7]). Both of them have shown the redundancy of one of the axioms of Hájek's axiomatization of BL-logics [4].

CONCLUSIONS

We briefly recalled what a non-associative fuzzy logic L_{CBA} is and where it can be useful. Then we significantly simplify the axiomatic system L_{CBA} for fuzzy logic with a non-associative conjunction as given in [1]. We show that three axioms can be omitted. Note that generally, the removal of the redundant axioms from a axiomatic system is desirable because it simplifies definitions and shortens proofs.

Note that although the independence of axioms is a highly desirable property, superfluous axioms still could be of use (as tautologies in the considered theory) when proving results in the considered theory, making proofs more transparent.

REFERENCES

- [1] M. Botur and R. Halaš, *Commutative basic algebras and non-associative fuzzy logics*, Arch. Math. Logic **48** (2009) 243–255. doi:10.1007/s00153-009-0125-7
- [2] M. Botur, I. Chajda, R. Halaš, J. Kühr and J. Paseka, *Algebraic Methods in Quantum Logic* (Palacký University, Olomouc, 2014), 200 pages, ISBN 978-80-244-4166-5.
- [3] P. Cintula, *Short note: on the redundancy of axiom (A3) in BL and MTL*, Soft Computing **9** (2005) 942–942. doi:10.1007/s00500-004-0445-9
- [4] P. Hájek, *Metamathematics of Fuzzy Logic*, vol. 4 of Trends in Logic (Dordrecht, Kluwer, 1998), 299 pages, ISBN 978-94-011-5300-3.
- [5] P. Hájek and R. Mesiar, *On copulas, quasicopulas and fuzzy logic*, Soft Computing **12** (2008) 1239–1243. doi:10.1007/s00500-008-0286-z
- [6] V. Kreinovich, *Towards more realistic (e.g., non-associative) and- and or-operations in fuzzy logic*, Soft Computing **8** (2004) 274–280. doi:10.1007/s00500-003-0272-4
- [7] S. Lehmke, *Fun with automated proof search in basic propositional fuzzy logic*, in: Abstracts of the Seventh International Conference FSTA 2004 (P.E. Klement, R. Mesiar, E. Drobná and F. Chovanec, eds.), (Liptovský Mikuláš), 2004, pp. 78–80.
- [8] R.B. Nelsen, *An Introduction to Copulas* (Springer-Verlag, New York, 2006), 286 pages, ISBN 978-0-387-28659-4.
- [9] R.R. Yager, *Modelling holistic fuzzy implication using co-copulas*, Fuzzy Optim. Decis. Making **5** (2006) 207–226. doi:10.1007/s10700-006-0011-2
- [10] H.H. Zimmerman and P. Zysno, *Latent connectives in human decision making*, Fuzzy Sets and Systems **4** (1980) 37–51. doi:10.1016/0165-0114(80)90062-7

Received 25 February 2016

Revised 10 March 2016