Discussiones Mathematicae
General Algebra and Applications 37 (2017) 31–43
doi:10.7151/dmgaa.1265

ZERO-DIVISOR GRAPHS OF REDUCED RICKART $*$-RINGS

A.A. Patil

Department of Mathematics
Garware College of Commerce, Pune-411004, India

e-mail: avipmj@gmail.com

and

B.N. Waphare

Center for Advanced Studies in Mathematics
Department of Mathematics
Savitribai Phule Pune University, Pune-411007, India

e-mail: waphare@yahoo.com
bnwaph@math.unipune.ac.in

Abstract

For a ring A with an involution $*$, the *zero-divisor graph* of A, $\Gamma^*(A)$, is the graph whose vertices are the nonzero left zero-divisors in A such that distinct vertices x and y are adjacent if and only if $xy^* = 0$. In this paper, we study the zero-divisor graph of a Rickart $*$-ring having no nonzero nilpotent element. The distance, diameter, and cycles of $\Gamma^*(A)$ are characterized in terms of the collection of prime strict ideals of A. In fact, we prove that the clique number of $\Gamma^*(A)$ coincides with the cellularity of the hull-kernel topological space $\Sigma(A)$ of the set of prime strict ideals of A, where cellularity of the topological space is the smallest cardinal number m such that every family of pairwise disjoint non-empty open subsets of the space have cardinality at most m.

Keywords: reduced ring, Rickart $*$-ring, zero-divisor graph, prime strict ideals.

2010 Mathematics Subject Classification: Primary: 16W10, Secondary: 05C25, 05C15.

1. Introduction

An *involution* '$*$', on an associative ring A is a mapping $* : A \to A$ such that $(a+b)^* = a^* + b^*$, $(ab)^* = b^*a^*$ and $(a^*)^* = a$, for all $a, b \in A$. A ring with an involution $*$ is called a $*$-*ring*. Clearly, identity mapping is an involution if and only if the ring is commutative. An element e in a $*$-ring A is a *projection* if it is self-adjoint (i.e., $e = e^*$) and idempotent (i.e., $e^2 = e$). By $\tilde{A}$, we denote the set of all projections in A. Let S be a nonempty subset of A. We write $r(S) = \{a \in A \mid sa = 0, \ \forall \ s \in S\}$, the *right annihilator* of S in A. A *Rickart* $*$-*ring* is a $*$-ring in which right annihilator of every element is generated, as a right ideal, by a projection in A. Every Rickart $*$-ring contains a unity. For each element a in a Rickart $*$-ring, there is a unique projection e such that $ae = a$ and $ax = 0$ if and only if $ex = 0$; called the *right projection* of a, $RP(a)$. In fact, $r(\{a\}) = (1 - RP(a))A$. Similarly, the left projection, $LP(a)$, is defined for each element a in a Rickart $*$-ring. The set of projections in a Rickart $*$-ring A forms a lattice, denoted by $L(\tilde{A})$, under the partial order '$e \leq f$ if and only if $e = fe$'. In that case, $e \vee f = f + RP(e(1-f))$ and $e \wedge f = e - LP(e(1-f))$ (see Berberian [5]). An ideal(subring) I is a $*$-*ideal*($*$-*subring*) if $x \in I$ implies $x^* \in I$. An ideal I of a Rickart $*$-ring A is called a *strict ideal* if $x \in I$ implies $RP(x) \in I$. A proper strict ideal P of a Rickart $*$-ring A is called *prime strict*, if for strict ideals I, J of A, $IJ \subseteq P$ implies $I \subseteq P$ or $J \subseteq P$. Let $\Sigma(A)$ denote the set of all prime strict ideals of a Rickart $*$-ring A. Thakare *et al.* [19] studied the hull-kernel topology on $\Sigma(A)$, where A is a reduced Rickart $*$-ring (i.e., a Rickart $*$-ring having no nonzero nilpotent element).

Being motivated from Beck [4], Anderson *et al.* [3] defined the zero-divisor graph for a commutative ring. Further study of this graph, such as connectedness, diameter, girth, etc. is found in [1, 2, 11]. Later on, Redmond [18] generalized the concept of the zero-divisor graph to a non-commutative ring in the following way: Let R be a ring. Then the undirected zero-divisor graph of R, denoted by $\overline{\Gamma(R)}$, is the graph whose vertices are the non-zero zero-divisors of R, and there is an edge between two distinct vertices a and b if and only if either $ab = 0$ or $ba = 0$. This concept is then generalized to semigroups by DeMeyer *et al.* in [7, 8], to semirings by Dolžan in [6]. Further, this concept is well studied in ordered structures such as lattices, meet-semilattices, posets (see [9, 13, 15, 16]). Recently, Sen Gupta *et al.* [10] defined a new graph for a ring with unity by extending the definition of the usual zero-divisor graph. In [17], the authors extended the concept of a zero-divisor graph to $*$-rings as follows: Let A be a $*$-ring. We associate a simple undirected graph $\Gamma^*(A)$ to A whose vertex set is $V(\Gamma^*(A)) = \{a(\neq 0) \in A \mid ab = 0, \text{ for some nonzero } b \in A\}$ (i.e., nonzero left zero-divisors) and two distinct vertices x and y are adjacent if and only if $xy^* = 0$. The zero-divisor graph of a $*$-ring A is denoted by $\Gamma^*(A)$. In the case of reduced Rickart $*$-rings, this graph is the same as the graph introduced by Nimbhorkar in [12].

In this paper, we continue our study of the zero-divisor graphs of Rickart $*$-rings that was started in [14]. Here particularly we deal with the reduced Rickart $*$-rings. The open subsets of the hull-kernel topological space $\Sigma(A)$ of the set of prime strict ideals of A are used to give algebraic and topological characterizations of distances, diameters, and cycles. We show that the clique number of $\Gamma^*(A)$ and the cellularity of $\Sigma(A)$ coincide. In fact, we have obtained the main results of Samei [11] to reduced Rickart $*$-rings.

For undefined concepts in Rickart $*$-rings and graphs, see [5, 20] respectively.

2. Main results

We recall some definitions that are used in the sequel.

Let A be a reduced Rickart $*$-ring and $\Sigma(A)$ be the set of all prime strict ideals in A. Then for $x \in A$ and any subset S of A, we define $H(S) = \{P \in \Sigma(A) \mid S \subseteq P\}$, $H(x) = H(\{x\})$, $B(S) = \{P \in \Sigma(A) \mid S \nsubseteq P\}$, $B(x) = B(\{x\})$ and $< x >=$ the ideal generated by x. For any subset T of $\Sigma(A)$, the kernel of T is the set $K(T) = \bigcap\{P \mid P \in T\}$, and the hull of $C \subseteq A$ is $H(C)$.

At the outset, we list the following observations made in [19, page 67].

Lemma 2.1. *Let A be a reduced Rickart $*$-ring and $\Sigma(A)$ is the set of all prime strict ideals in A. Then*

(1) $B(x) = B(RP(x)) = B(< RP(x) >)$.

(2) $\bigcup_{i\in\Lambda} B(x_i) = B(\{x_i | i \in \Lambda\})$, $B(I \cap J) = B(I) \cap B(J)$, $B(0) = \emptyset$ *and* $B(A) = B(1) = \Sigma(A)$.

(3) $H(x) = H(< x >) = H(< RP(x) >)$, $H(0) = \Sigma(A), H(A) = H(1) = \emptyset$, $H(\cup E_i) = \cap H(E_i)$ *and* $H(I \cap J) = H(I) \cup H(J)$, *where each* E_i, I *and* J *are ideals of* A.

(4) *Every prime strict ideal contains* e *or* $1-e$ *but not both, for any projection* e.

(5) $\bigcap\{P \mid P \in \Sigma(A)\} = \{0\}$.

We note from the above Lemma 2.1 that the sets $\{B(x) \mid x \in A\}$ form a basis for open sets and define a topology on $\Sigma(A)$ called the *hull-kernel topology.* The complement of $B(x)$ is $H(x)$ for any $x \in A$.

Remark 2.2. Following are the simple properties of a reduced Rickart $*$-ring A.

(1) For $a, b \in A$, $ab = 0$ if and only if $ba = 0$. For, let $ab = 0$. Then $(ba)^2 = baba = 0$. Since A is reduced, we get $ba = 0$.

(2) Every projection in A is central. For, if e is a projection, then $e(1-e) = (1-e)e = 0$ and $(ex(1-e))^2 = ex(1-e)ex(1-e) = 0$ for any $x \in A$.

As is reduced, $ex(1-e)=0$. Therefore $ex=exe$ and $ex^*=ex^*e$, i.e., $xe=exe=ex$, for all $x\in A$. Thus e is a central projection.

(3) For any $a\in A$, $RP(a)=RP(a^*)$, since every projection is central, we have $RP(a)=LP(a)$. This together with $RP(a^*)=LP(a)$ gives $RP(a)=RP(a^*)$.

(4) $RP(xy)=RP(x)RP(y)$, for any $x,y\in A$. For, since $RP(x)$ and $RP(y)$ both are central, we get $xyRP(x)RP(y)=xRP(x)yRP(y)=xy$. Let $xyz=0$. By definition of $RP(x)$, we get $RP(x)yz=0$. This together with centrality of $RP(x)$ imply that $yRP(x)z=0$. Again by definition of right projection, we get $RP(y)RP(x)z=0$, i.e., $RP(x)RP(y)z=0$. Note that product of two commuting projections is a projection. Therefore by Remark 2.2(2), the projection $RP(x)RP(y)$ satisfies the conditions of $RP(xy)$, hence $RP(xy)=RP(x)RP(y)$.

(5) For any projection $e\in A$, the ideal $<e>$ is a strict ideal. For, if $x\in<e>$, then $xe=x$. Hence $RP(x)=RP(xe)=RP(x)e\in<e>$.

Lemma 2.3. *Let A be a reduced Rickart $*$-ring. Then the following statements hold.*

(1) *Every strict ideal is a $*$-ideal.*

(2) *Let P be a strict ideal. Then P is a prime strict ideal if and only if $ab\in P$ implies $a\in P$ or $b\in P$.*

Proof. (1) Let I be a strict ideal of A and $a\in I$. Hence $RP(a)\in I$. Then $a^*=a^*RP(a^*)=a^*RP(a)\in I$, by Remark 2.2(3). Thus I is a $*$-ideal.

(2) Let P be a strict ideal. Suppose that P is a prime strict ideal and $ab\in P$ with $a\notin P$. We claim that $b\in P$. Since $ab\in P$ and P is a strict ideal, we get $RP(ab)\in P$. By Remark 2.2(4), $RP(ab)=RP(a)RP(b)$. Observe that $<RP(a)>\cdot<RP(b)>=<RP(a)RP(b)>$. Thus $I=<RP(a)>$ and $J=<RP(b)>$ are two strict ideals (by Remark 2.2(5)) such that $IJ\subseteq P$ with $I\nsubseteq P$. Since P is a prime strict ideal, we get $J\subseteq P$. Hence $RP(b)\in P$. Thus $b=bRP(b)\in P$. Conversely, suppose that $ab\in P$ implies $a\in P$ or $b\in P$. Let I and J be strict ideals of A such that $IJ\subseteq P$ with $I\nsubseteq P$. We claim that $J\subseteq P$. On the contrary assume that $J\nsubseteq P$. Let $a,b\in A$ be such that $a\in I\setminus P$ and $b\in J\setminus P$. Then $ab\in IJ\subseteq P$ with $a,b\notin P$, a contradiction. Hence $J\subseteq P$. Thus P is a prime strict ideal. ■

For a $*$-ring A, two graphs $\Gamma^*(A)$ and $\overline{\Gamma(A)}$ need not be isomorphic, see Example 2.6.

Theorem 2.4. *For a reduced Rickart $*$-ring A, $\Gamma^*(A)\cong\overline{\Gamma(A)}$.*

Recall that, a graph G is said to be *connected* if there is a path between any two distinct vertices of G. For two vertices x and y of G, we define $d(x, y)$ to be the length of the shortest path from x to y ($(d(x, x) = 0$ and $d(x, y) = \infty$ if there is no such path). The *diameter* of G is defined as $diam(G) = sup\{d(x, y) \mid x$ and y are vertices of $G\}$. The *girth* of G, denoted by $gr(G)$, is the length of the shortest cycle in G ($gr(G) = \infty$ if G contains no cycle). A subset S of $V(G)$ is called a *clique* if any two distinct vertices of S are adjacent; the *clique number*, $\omega(G)$, is the least upper bound of the size of the cliques in G.

Theorem 2.5. *Let A be a reduced Rickart $*$-ring such that $V(\Gamma^*(A)) \neq \emptyset$. Then $\Gamma^*(A)$ is connected with $diam(\Gamma^*(A)) \leq 3$.*

Proof. Since A is a reduced ring, by Remark 2.2(1), $ab = 0$ if and only if $ba = 0$ if and only if $ab^* = 0$. Hence the proof follows by Redmond [18, Theorem 3.2]. ■

Following example shows that Rickartness is necessary in Theorem 2.4 and Theorem 2.5.

Example 2.6. Let $A = \mathbb{Z}_3 \times \mathbb{Z}_3$ with $(a, b)^* = (b, a)$ as an involution. Then A is a reduced but not a Rickart $*$-ring (since $(0, 1)^*(0, 1) = (0, 0)$ with $(0, 1) \neq (0, 0)$). Here $V(\Gamma^*(A)) = V(\overline{\Gamma(A)}) = \{a = (0, 1), b = (1, 0), c = (0, 2), d = (2, 0)\}$. Figure 1 shows that $\Gamma^*(A)$ and $\overline{\Gamma(A)}$ are non-isomorphic.



Figure 1.

Next result gives an equivalent condition for adjacency in $\Gamma^*(A)$.

Lemma 2.7. *Let A be a reduced Rickart $*$-ring such that $V(\Gamma^*(A)) \neq \emptyset$. Then the vertices a and b are adjacent in $\Gamma^*(A)$ if and only if $B(a) \cap B(b) = \emptyset$.*

Proof. Suppose a and b are adjacent in $\Gamma^*(A)$. Let P be any prime strict ideal of A. Then $ab^* = 0 \in P$. Since P is a prime strict ideal, we get either $a \in P$ or $b^* \in P$. Since P is a strict ideal, $b^* \in P$ whenever $b \in P$. Hence every prime strict ideal either contains a or b. Therefore $B(a) \cap B(b) = \emptyset$. Conversely, suppose that $B(a) \cap B(b) = \emptyset$. Hence for any prime strict ideal P, we have either $a \in P$ or $b \in P$, i.e., $a \in P$ or $b^* \in P$. Consequently $ab^* \in P$, for all $P \in \Sigma(A)$, hence $ab^* \in \bigcap\{P \mid P \in \Sigma(A)\} = \{0\}$. Thus $ab^* = 0$. ■

Observe that, for each projection e in a reduced Rickart $*$-ring, by Lemma 2.1(4), we have $B(e) = H(1-e)$. This leads to the following:

Corollary 2.8. *Let A be a reduced Rickart $*$-ring such that $V(\Gamma^*(A)) \neq \emptyset$. Then the vertices a and b are adjacent in $\Gamma^*(A)$ if and only if $B(a) \subseteq H(b)$.*

Remark 2.9. In a reduced Rickart $*$-ring, $H(1-RP(a))$ = set of prime strict ideals containing $1-RP(a) = B(RP(a)) = B(a)$.

The following result characterizes the distance between two vertices. We use the notation $a \leftrightarrow b$ to indicate that a and b are adjacent.

Proposition 2.10. *Let A be a reduced Rickart $*$-ring and $a, b, c \in V(\Gamma^*(A))$ be distinct vertices. Then*

(1) *c is adjacent to both a and b if and only if $B(a) \cup B(b) \subseteq H(c)$.*

(2) *$d(a,b) = 1$ if and only if $B(a) \cap B(b) = \emptyset$.*

(3) *$d(a,b) = 2$ if and only if $B(a) \cap B(b) \neq \emptyset$ and $B(a) \cup B(b) \neq \Sigma(A)$.*

(4) *$d(a,b) = 3$ if and only if $B(a) \cap B(b) \neq \emptyset$ and $B(a) \cup B(b) = \Sigma(A)$.*

Proof. (1) Suppose that c is adjacent to both a and b. By Corollary 2.8, $B(a) \subseteq H(c)$ and $B(b) \subseteq H(c)$. Hence $B(a) \cup B(b) \subseteq H(c)$. Conversely, suppose that $B(a) \cup B(b) \subseteq H(c)$. Hence $B(a) \subseteq H(c)$ and $B(b) \subseteq H(c)$. Consequently, $B(a) \cap B(c) = \emptyset$ and $B(b) \cap B(c) = \emptyset$. By Lemma 2.7, c is adjacent to a and c is adjacent to b.

(2) Follows from Lemma 2.7.

(3) Suppose that $d(a,b) = 2$. Let $a \leftrightarrow d \leftrightarrow b$ be a path of length 2. Since a and b are non-adjacent, $B(a) \cap B(b) \neq \emptyset$. Here d is adjacent to both a and b, hence by (1) above, $B(a) \cup B(b) \subseteq H(d)$. Therefore $B(a) \cup B(b) \neq \Sigma(A)$, otherwise $H(d) = \Sigma(A)$ which yields $d \in \bigcap_{P \in \Sigma(A)} P = \{0\}$ giving $d = 0$, a contradiction. Thus $B(a) \cap B(b) \neq \emptyset$ and $B(a) \cup B(b) \neq \Sigma(A)$. Conversely, suppose that $B(a) \cap B(b) \neq \emptyset$ and $B(a) \cup B(b) \neq \Sigma(A)$. By (2) above, a and b are non-adjacent. Let $P \in \Sigma(A)$ be such that $P \notin B(a) \cup B(b)$. Since $a \in P$, we get $RP(a) \in P$. By Lemma 2.1(4), $1 - RP(a) \notin P$. Similarly, $RP(b) \in P$ and $1 - RP(b) \notin P$. Therefore $(1-RP(a))(1-RP(b)) \notin P$. Consequently, $c = (1-RP(a))(1-RP(b)) \neq 0$. Then $a \leftrightarrow c \leftrightarrow b$ is a path. Therefore $d(a,b) = 2$.

(4) Follows from (2), (3) above and Theorem 2.5. ∎

Now, in the following result, we characterize the diameter of $\Gamma^*(A)$.

Theorem 2.11. *For a finite reduced Rickart $*$-ring A following statements hold:*

(1) *$diam(\Gamma^*(A)) = 1$ if and only if $A = \mathbb{Z}_2 \oplus \mathbb{Z}_2$ with identity mapping as an involution.*

(2) *$diam(\Gamma^*(A)) = 2$ if and only if $|V(\Gamma^*(A))| \geq 3$ and $L(\tilde{A})$ contains exactly two atoms.*

(3) *$diam(\Gamma^*(A)) = 3$ if and only if $L(\tilde{A})$ contains at least three atoms.*

Proof. (1) One way is clear. Conversely, suppose that $diam(\Gamma^*(A)) = 1$. Therefore $\Gamma^*(A)$ is a complete graph. Let e be a non-trivial projection in A (since $V(\Gamma^*(A)) \neq \emptyset$, such e exists). Then $A = eA \oplus (1-e)A$. By completeness of $\Gamma^*(A)$, we get $eA = \mathbb{Z}_2$ and $(1-e)A = \mathbb{Z}_2$. Thus $A = \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Further, by Remark 2.2(5), eA is strict ideal. Hence for $x \in eA \setminus \{e\}$, we have $RP(x) = RP(x^*) = e$. This gives $x^* = x^*e \in eA$, which gives $x^* = x$. Therefore the involution on eA and $(1-e)A$, and hence on A is identity.

(2) Let $diam(\Gamma^*(A)) = 2$. Clearly $|V(\Gamma^*(A))| \geq 3$. Also, for a finite Rickart $*$-ring A with $V(\Gamma^*(A)) \neq \emptyset$, the lattice $L(\tilde{A})$ contains at least two atoms. Suppose e, f and g are three atoms in $L(\tilde{A})$, hence $ef = fg = eg = 0$. If $e + f = 1$, then $g = eg + fg = 0$, a contradiction. Hence $e + f \neq 1$. Similarly, $f + g \neq 1$ and $e + g \neq 1$. This gives $(1-e)(1-f) \neq 0$, otherwise $e + f = 1$. If there is a projection h such that $(1-e) \leftrightarrow h \leftrightarrow (1-f)$ is a path, then $h = he = hf$ implies $h = hf = hef = 0$. Thus $1-e$ and $1-f$ are nonadjacent and there is no path of length 2 joining them, hence $d(1-e, 1-f) \geq 3$, a contradiction to the fact that $diam(\Gamma^*(A)) = 2$. Therefore $L(\tilde{A})$ contains exactly two atoms. Conversely, suppose that $|V(\Gamma^*(A))| \geq 3$ and $L(\tilde{A})$ contains exactly two atoms, say e_1 and e_2. Let a and b be any two vertices of $\Gamma^*(A)$. We consider the following two cases.

Case (i) Suppose $RP(a)$ and $RP(b)$ have common atom, say e_1, i.e., $e_1 \leq RP(a)$ and $e_1 \leq RP(b)$. Then $e_1 \leq RP(a)RP(b)$, hence $RP(a)RP(b) \neq 0$. On the other hand, $e_2 \leq 1 - RP(a)$ and $e_2 \leq 1 - RP(b)$ which gives $(1 - RP(a))(1 - RP(b)) \neq 0$. Hence $RP(a) \leftrightarrow (1 - RP(a))(1 - RP(b)) \leftrightarrow RP(b)$ is a path joining $RP(a)$ and $RP(b)$. Thus $d(a, b) = 2$.

Case (ii) Suppose $RP(a)$ and $RP(b)$ don't have common atom. Without loss of generality, assume that $e_1 \leq RP(a)$ and $e_2 \leq RP(b)$. Then $e_2RP(a) = e_1RP(b) = 0$. This gives $e_1RP(a)RP(b) = e_2RP(a)RP(b) = 0$, hence $e_1 \nleq RP(a)RP(b)$ and $e_2 \nleq RP(a)RP(b)$. Since $L(\tilde{A})$ contains exactly two atoms, we must have $RP(a)RP(b) = 0$. Hence $d(a, b) = 1$.

Thus $diam(\Gamma^*(A)) \leq 2$. Since $|V(\Gamma^*(A))| \geq 3$ and $L(\tilde{A})$ contains exactly two atoms, there exists two vertices x and y such that $RP(x)$ and $RP(y)$ contains a common atom in $L(\tilde{A})$. Hence by Case(i) above, we get $d(x, y) = 2$. Therefore $diam(\Gamma^*(A)) = 2$.

(3) If $diam(\Gamma^*(A)) = 3$, then by (2), $L(\tilde{A})$ contains at least three atoms. Conversely, if $L(\tilde{A})$ contains three atoms, then as in the proof of (2), $d(1-e, 1-f) = 3$. By Theorem 2.5, $diam(\Gamma^*(A)) = 3$. ■

It is known that the girth of the zero-divisor graph is either 3 or 4 (if it contains a cycle); see [17]. The following theorem characterizes the diameter and the girth of $\Gamma^*(A)$ in terms of the number of prime strict ideals of A.

Theorem 2.12. *Let A be a reduced Rickart $*$-ring such that $V(\Gamma^*(A)) \neq \emptyset$.*

(1) *If $A \ncong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ with identity involution, then $diam(\Gamma^*(A)) = \min\{|\Sigma(A)|, 3\}$.*

(2) *If $|\Sigma(A)| = 2$, then $gr(\Gamma^*(A)) = 4$ or ∞; otherwise $gr(\Gamma^*(A)) = 3$.*

Proof. (1) Since $V(\Gamma^*(A)) \neq \emptyset$, we get $|\Sigma(A)| \geq 2$. Also, $A \ncong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ which gives $diam(\Gamma^*(A)) \neq 1$. Further, by Theorem 2.5, $diam(\Gamma^*(A)) \leq 3$. It is enough to show that $diam(\Gamma^*(A)) = 3$ if and only if $|\Sigma(A)| \geq 3$. Suppose that $|\Sigma(A)| \geq 3$ and P_1, P_2, P_3 be three distinct prime strict ideals of A. Since $\Sigma(A)$ is Hausdorff [19, Lemma 3.4], there are $a_i \in A$ such that $P_i \in B(a_i)$ and $B(a_i) \cap B(a_j) = \emptyset$, for $i \neq j$ and $i, j \in \{1, 2, 3\}$. By Lemma 2.7, we have a_i and a_j are adjacent in $\Gamma^*(A)$ for $i \neq j$. This implies that $gr(\Gamma^*(A)) = 3$. By Remark 2.9, $P_1 \in H(1 - RP(a_1))$ and $P_2 \in H(1 - RP(a_2))$. Since $a_1a_3 = a_2a_3 = 0 \in P_3$ and $a_3 \notin P_3$, we have $a_1, a_2 \in P_3$. Hence $1 - RP(a_1), 1 - RP(a_2) \notin P_3$. Therefore $P_3 \in B(1 - RP(a_1)) \cap B(1 - RP(a_2))$. Since $B(a_1) \cap B(a_2) = \emptyset$, we get $H(RP(a_1)) \cup H(RP(a_2)) = \Sigma(A)$. This together with $H(RP(a_i)) = B(1 - RP(a_i))$ for $i = 1, 2$, gives $B(1 - RP(a_1)) \cup B(1 - RP(a_2)) = \Sigma(A)$. By Proposition 2.10, $d(1 - RP(a_1), 1 - RP(a_2)) = 3$. Therefore $diam(\Gamma^*(A)) = 3$. Conversely, suppose that $diam(\Gamma^*(A)) = 3$. Let $a \leftrightarrow a_1 \leftrightarrow a_2 \leftrightarrow b$ is a path of length 3 in $\Gamma^*(A)$. Since $V(\Gamma^*(A)) \neq \emptyset$ and $A \ncong \mathbb{Z}_2 \oplus \mathbb{Z}_2$, A has at least two prime strict ideals. On the contrary assume that P_1 and P_2 are the only prime strict ideals of A. Observe that, for no vertex x we have $B(x) = \{P_1, P_2\}$. Without loss of generality, assume that $B(a) = \{P_1\}$. Since a is adjacent to a_1 and nonadjacent to b, we must have $B(a_1) = \{P_2\}$ and $B(b) = \{P_1\}$. Which gives $B(a_1) \cap B(b) = \emptyset$, i.e., a_1 is adjacent to b, a contradiction to the fact that $d(a, b) = 3$. Therefore $|\Sigma(A)| \geq 3$.

(2) By the proof of part (1), $|\Sigma(A)| \geq 3$ implies that $gr(\Gamma^*(A)) = 3$. Now suppose $|\Sigma(A)| = 2$, say $\Sigma(A) = \{P_1, P_2\}$. Since $V(\Gamma^*(A)) \neq \emptyset$, A contains at least one non-trivial projection, say, e. Further, A is a reduced ring, hence e is central. Therefore $A = eA \oplus (1 - e)A$. Next, we claim that eA does not contain nonzero zero-divisor. On the contrary assume that, $xy = 0$ for some nonzero $x, y \in eA$. Then x and y are adjacent in $\Gamma^*(A)$, giving $B(x) \cap B(y) = \emptyset$. Since $\Sigma(A) = \{P_1, P_2\}$ and $H(e) \notin \{\emptyset, \Sigma(A)\}$, without loss of generality assume that, $H(e) = \{P_1\}$. Then $x, y \in eA \subseteq P_1$. Since $H(x), H(y) \notin \{\emptyset, \Sigma(A)\}$, we get $H(x) = H(y) = \{P_1\}$, hence $B(x) = B(y) = \{P_2\}$, a contradiction to the fact that $B(x) \cap B(y) = \emptyset$. Therefore eA does not contain nonzero zero-divisor. Similarly, $(1 - e)A$ does not contain nonzero zero-divisor. If either $|eA| = 2$ or $|(1 - e)A| = 2$, then $gr(\Gamma^*(A)) = \infty$; else it is easy to see that $gr(\Gamma^*(A)) = 4$. ■

The associated number $e(a)$- the *eccentricity of* a denoted by $e(a)$, of a vertex of a graph G is defined to be $e(a) = max\{d(a,b) \mid a \neq b\}$.

Remark 2.13. Note that, for any vertex a of $\Gamma^*(A)$, $d(a,b) \leq 3$ for any $b \in V(\Gamma^*(A)) \setminus \{a\}$. Therefore $e(a) \leq 3$.

Theorem 2.14. *In a reduced Rickart $*$-ring A,*

(1) $e(a) = 1$ *if and only if* a *is a projection and* $|\Sigma(A)| = | < a > | = 2$.
If $e(a) \neq 1$. *Then*

(2) $e(a) = 2$ *if and only if* $|B(a)| = 1$.

(3) $e(a) = 3$ *if and only if* $|B(a)| > 1$.

Proof. (1) Let $e(a) = 1$, i.e., a is adjacent to all the other vertices of $\Gamma^*(A)$. If a is not a projection, then we get a contradiction since a and $RP(a)$ are always nonadjacent in $\Gamma^*(A)$. Therefore a is a projection. Let $b \in < a >$ with $b \neq a$, hence $b = ab$. Since a and b are adjacent, we get $ab = 0$. Consequently, $b = ab = 0$. Therefore $| < a > | = 2$. Also, by Remark 2.2(5), $< a >$ is a strict ideal of A. Next, let P be a prime strict ideal containing a and $y \in P \setminus \{a\}$. Since P is prime strict ideal, we have $RP(y) \neq 1$. If $R(y) \neq 0$, then a is adjacent to both $RP(y)$ and $1 - RP(y)$, which gives $a = 0$, a contradiction. Hence $RP(y) = 0$, consequently $y = 0$. Therefore $P = \{0, a\}$. Thus $\{0, a\}$ is a prime strict ideal. Now, let $P_1 \in B(a)$. Then for any $x \in V(\Gamma^*(A)) \setminus \{a\}$, $aRP(x) = 0$. Hence $aRP(x) \in P_1$ with $a \notin P_1$ and since P_1 is a prime strict ideal, we have $RP(x) \in P_1$ which yields $x \in P_1$, for all $x \in V(\Gamma^*(A)) \setminus \{a\}$. Therefore $V(\Gamma^*(A)) \setminus \{a\} \subseteq P_1$. On the other hand, for any nonzero $y \in P_1$, $RP(y) \neq 1$ and $y(1 - RP(y)) = 0$. Hence $y \in V(\Gamma^*(A))$. Therefore $P_1 = \{0\} \cup [V(\Gamma^*(A)) \setminus \{a\}]$. Thus $|\Sigma(A)| = 2$. Conversely, suppose that a is a projection and $|\Sigma(A)| = | < a > | = 2$. Let P_1 and P_2 be two prime strict ideals of A with $a \in P_1 \setminus P_2$. Hence $1 - a \in P_2$. Then for any $x \in P_1, RP(x)(1 - a) \in P_1 \cap P_2 = \{0\}$, which gives $RP(x) = RP(x)a$. This yields $x = xa \in < a >$. Therefore $P_1 = \{0, a\}$. Since $| < a > | = 2$, there is no b different from a such that $RP(b) = a$. Hence for any $b \in V(\Gamma^*(A)) \setminus \{a\}$, we get $RP(b) \in P_2$. This gives $RP(b)a \in P_1 \cap P_2 = \{0\}$, giving $RP(b)a = 0$, i.e., b and a are adjacent. Thus $e(a) = 1$.

For (2) and (3), suppose that $e(a) \neq 1$. Then by Theorem 2.11, $e(a) = 2$ or 3.

Let $|B(a)| = 1$ and $P \in \Sigma(A)$ be such that $a \notin P$. As $e(a) \neq 1$ there exists $b \in V(\Gamma^*(A)) \setminus \{a\}$ such that a and b are non-adjacent. Hence $B(a) \cap B(b) \neq \emptyset$. Now $B(a) \cap B(b) \subseteq B(a)$ which is a singleton set, hence we get $B(a) \subseteq B(b)$. Therefore $B(a) \cup B(b) = B(b) \neq \Sigma(A)$. By Theorem 2.11, $e(a) = 2$.

Now suppose $|B(a)| > 1$ and $P_1, P_2 \in \Sigma(A)$ such that $a \notin P_1 \cup P_2$. Let $b \in P_1 \setminus P_2$ then $b \in P_1$ and $b \notin P_2$. Therefore $P_2 \in B(a) \cap B(b)$, which gives

$d(a,b) \neq 1$. Let $P \in B(ab)$, i.e., $ab \notin P$. Since P is a prime strict ideal, we have $a \notin P$ and $b \notin P$. Hence $P \in B(a)$. Thus $B(ab) \subseteq B(a)$ which gives $B(a) \cup B(1 - RP(a)RP(b)) = \Sigma(A)$. If $RP(a)(1 - RP(a)RP(b)) = 0$, then $RP(a) = RP(a)RP(b) \in P_1$ giving $a \in P_1$, a contradiction. Hence a and $1 - RP(a)RP(b)$ are non-adjacent in $\Gamma^*(A)$. By Theorem 2.11, $d(a,(1 - aRP(b))) = 3$. Thus $e(a) = 3$. ■

A graph G is *triangulated* if each vertex of G is a vertex of a triangle. Let $Z(A)$ be the set of zero-divisors in A. Now, we characterize the triangulated zero-divisor graphs.

Theorem 2.15. *For a reduced Rickart $*$-ring A,*

(1) *$\Gamma^*(A)$ is triangulated if and only if $|H(a)| > 1$, for all $a \in V(\Gamma^*(A))$.*

(2) *If $2 \notin Z(A)$, then every vertex of $\Gamma^*(A)$ is a 4-cycle-vertex.*

Proof. (1) Suppose that $\Gamma^*(A)$ is triangulated. On the contrary, assume that there is a vertex a such that a belongs to unique prime strict ideal, say, Q of A, i.e., $|H(a)| = 1$. Since $\Gamma^*(A)$ is triangulated, there exists $b, c \in V(\Gamma^*(A))$ such that $ab^* = bc^* = ca^* = 0$. Therefore $B(a) \cap B(b) = B(a) \cap B(c) = B(b) \cap B(c) = \emptyset$. This gives $B(a) \subseteq H(b)$ and $B(c) \subseteq H(a) \cap H(b)$, hence $H(a) \cap H(b) \neq \emptyset$. Also, $H(a) \cap H(b) \subseteq H(a)$ and $H(a)$ is a singleton set. Therefore $H(a) \subseteq H(b)$. Now $\Sigma(A) = B(a) \cup H(a) \subseteq H(b)$ gives $H(b) = \Sigma(A)$. Hence $b \in \bigcap_{P \in \Sigma(A)} P = \{0\}$ giving $b = 0$, a contradiction. Therefore $|H(a)| > 1$, for all $a \in V(\Gamma^*(A))$.

Conversely, suppose that $|H(a)| > 1$, for all $a \in V(\Gamma^*(A))$. Let $a \in V(\Gamma^*(A))$ and P_1, P_2 be two distinct prime strict ideals containing a. Let $e = RP(a)$ and $I = (1-e)A$, i.e., $I = r(\{a\})$. Then $e \in P_1, P_2$. Since $A = eA \oplus (1-e)A$, if $P_1 \cap I = \{0\}$ and $P_2 \cap I = \{0\}$, we get $P_1 \subseteq eA$ and $P_2 \subseteq eA$. Then for any $x \in P_1$, we get $x = xe \in P_2$, i.e; $P_1 \subseteq P_2$. Similarly $P_2 \subseteq P_1$. This gives $P_1 = P_2$, a contradiction. Hence $P_1 \cap I \neq \{0\}$ or $P_2 \cap I \neq \{0\}$. Without loss of generality, suppose $P_1 \cap I \neq \{0\}$. Let $b \in P_1 \cap I$ with $b \neq 0$ and $f = RP(b)$. Then $(1-e)f = f$ giving $ef = 0$, hence a and b are adjacent. If $(1-e)(1-f) = 0$, then $1 - e = (1-e)f \in P_1$. Which gives $1 = 1 - e + e \in P_1$, a contradiction. Hence $(1-e)(1-f) \neq 0$ and $a \leftrightarrow b \leftrightarrow (1-e)(1-f) \leftrightarrow a$ is a triangle. Thus every vertex is a vertex of a triangle.

(2) Suppose that $2 \notin Z(A)$. Let $a \in V(\Gamma^*(A))$. Then there exists nonzero $b \in A$ such that $ab^* = 0$. Since A is reduced and $2 \notin Z(A)$, we get $2a \neq b$, $a \neq 2b$. Therefore $a, b, 2a, 2b$ all are distinct with $ab^* = 0 = (2a)b^* = (2a)(2b)^* = 2a(2b)^*$. Thus $a \leftrightarrow b \leftrightarrow 2a \leftrightarrow 2b \leftrightarrow a$ is a 4-cycle containing a. ■

If a and b are two vertices in $\Gamma^*(A)$, by $c(a,b)$ we mean the length of the smallest cycle containing a and b. For every two vertices a and b, all possible cases for $c(a,b)$ are given in the following theorem.

Theorem 2.16. *Let A be a reduced Rickart $*$-ring, $a, b \in V(\Gamma^*(A))$ and $2 \notin Z(A)$.*

(1) *$c(a,b) = 3$ if and only if $B(a) \cap B(b) = \emptyset$ and $RP(a) + RP(b) \neq 1$.*

(2) *$c(a,b) = 4$ if and only if either $B(a) \cap B(b) = \emptyset$ and $RP(a) + RP(b) = 1$ or $B(a) \cap B(b) \neq \emptyset$ and $d(a,b) = 2$.*

(3) *$c(a,b) = 6$ if and only if $d(a,b) = 3$.*

Proof. (1) Suppose that $B(a) \cap B(b) = \emptyset$ and $RP(a) + RP(b) \neq 1$. Then a and b are adjacent, hence $RP(a)RP(b) = 0$. If $(1 - RP(a))(1 - RP(b)) = 0$, then $RP(a) + RP(b) = 1$, a contradiction. Therefore $(1 - RP(a))(1 - RP(b)) \neq 0$ which yields $a \leftrightarrow (1 - RP(a))(1 - RP(b)) \leftrightarrow b \leftrightarrow a$ a $3-$cycle containing a and b. Thus $c(a,b) = 3$. Conversely, suppose that $c(a,b) = 3$. Then a and b are adjacent, hence $B(a) \cap B(b) = \emptyset$. Let $a \leftrightarrow b \leftrightarrow c$ be a 3-cycle. Then $RP(a)RP(b) = RP(a)RP(c) = RP(b)RP(c) = 0$. If $RP(a) + RP(b) = 1$, then $RP(c) = RP(a)RP(c) + RP(b)RP(c) = 0$, giving $c = 0$, a contradiction. Therefore $RP(a) + RP(b) \neq 1$.

(2) Let $c(a,b) = 4$. Suppose $B(a) \cap B(b) = \emptyset$. If $RP(a) + RP(b) \neq 1$, then by (1) above, $c(a,b) = 3$, a contradiction. Hence $RP(a) + RP(b) = 1$. Now suppose that $B(a) \cap B(b) \neq \emptyset$. Then a and b are non-adjacent. This together with $c(a,b) = 4$ gives $d(a,b) = 2$. Conversely, suppose that $B(a) \cap B(b) = \emptyset$ and $RP(a) + RP(b) = 1$. Since $2 \notin Z(A)$, we get $a \leftrightarrow b \leftrightarrow 2a \leftrightarrow 2b \leftrightarrow a$ a 4-cycle containing a and b. Next, we will show that there is no cycle of length 3 which contains a and b. If $c(a,b) = 3$, then by (1) above, we have $RP(a) + RP(b) \neq 1$, a contradiction. Therefore $c(a,b) = 4$. Suppose $B(a) \cap B(b) \neq \emptyset$ and $d(a,b) = 2$. Let $a \leftrightarrow c \leftrightarrow b$ be a path. Then $a \leftrightarrow c \leftrightarrow b \leftrightarrow 2c \leftrightarrow a$ is a 4-cycle containing a and b. Hence $c(a,b) = 4$.

(3) Suppose $d(a,b) = 3$. Let $a \leftrightarrow c \leftrightarrow d \leftrightarrow b$ be a path. Then $RP(c)RP(b) \neq 0$ (as $d(a,b) = 3$) which gives $a \leftrightarrow c \leftrightarrow d \leftrightarrow b \leftrightarrow (1 - RP(b)) \leftrightarrow RP(c)RP(b) \leftrightarrow a$ a 6-cycle containing a and b. Since $d(a,b) = 3$, we get $c(a,b) = 6$. Conversely, suppose that $c(a,b) = 6$. Then by (1) and (2) above, we get $d(a,b) \neq 1, 2$. Therefore $d(a,b) = 3$. ■

The *cellularity* of a topological space X is denoted by $c(X)$ and it is the smallest cardinal number m such that every family of pairwise disjoint non-empty open subsets of X has cardinality at most m. Next we show that the clique number of $\Gamma^*(A)$ and the cellularity of $\Sigma(A)$ coincide.

Theorem 2.17. *Let A be a reduced Rickart $*$-ring. Then $\omega(\Gamma^*(A)) = c(\Sigma(A))$.*

Proof. Let C be a clique in $\Gamma^*(A)$. Then for every $a, b \in C, ab^* = 0$, i.e., $B(a) \cap B(b) = \emptyset$. Then the collection $\mathcal{C} = \{B(a) \mid a \in C\}$ is a family of pairwise

disjoint non-empty open subsets of $\Sigma(A)$. Hence $\omega(\Gamma^*(A)) \leq c(\Sigma(A))$. Next, suppose that $\mathcal{C} = \{A_\lambda \mid \lambda \in \Lambda\}$ is a collection of pairwise disjoint non-empty open subsets of $\Sigma(A)$. For every $A_\lambda \in \mathcal{C}$ there exists $a_\lambda \in A$ such that $\emptyset \neq B(RP(a_\lambda)) \subseteq A_\lambda$. Clearly for every $A_\lambda, A'_\lambda \in \mathcal{C}$, we have $a_\lambda {a'_\lambda}^* = 0$. Hence $B = \{a_\lambda \mid \lambda \in \Lambda\}$ is a clique in $\Gamma^*(A)$. This implies that $c(\Sigma(A)) \leq \omega(\Gamma^*(A))$. Therefore $\omega(\Gamma^*(A)) = c(\Sigma(A))$. ■

Acknowledgements

The authors are deeply grateful to the referee for many fruitful suggestions which improved the presentation of the paper. The first author was financially supported by the University Grant Commission, New Delhi, India via a minor research project File No. 47-884/14(WRO).

References

[1] S. Akbari and A. Mohammadian, *On the zero-divisor graph of a commutative ring*, J. Algebra **274** (2004) 847–855. doi:10.1016/S0021-8693(03)00435-6

[2] D.F. Anderson, R. Levy and J. Shapirob, *Zero-divisor graphs, von Neumann regular rings and Boolean algebras*, J. Pure Appl. Alg. **180** (2003) 221–241. doi:10.1016/S0022-4049(02)00250-5

[3] D.F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999) 434–447. doi:10.1006/jabr.1998.7840

[4] I. Beck, *Coloring of commutative rings*, J. Algebra **116** (1988) 208–226. doi:10.1016/0021-8693(88)90202-5

[5] S.K. Berberian, Baer $*$-Rings (Springer-Verlag, Berlin and New York, 1972).

[6] D. Dolžan and P. Oblak, *The zero divisor graphs of rings and semirings*, Int. J. Algebra Comput. **22** 1250033 (2012) 20pp. doi:10.1142/S0218196712500336

[7] F. DeMeyer, T. McKenzie and K. Schneider, *The zero-divisor graph of a commutative semigroup*, Semigroup Forum **65** (2002) 206–214. doi:10.1007/s002330010128

[8] F. DeMeyer and L. DeMeyer, *Zero divisor graph of Semigroups*, J. Algebra **283** (2005) 190–198. doi:10.1016/j.jalgebra.2004.08.028

[9] R. Halaš and M. Jukl, *On Beck's colouring of posets*, Discrete Math. **309** (2009) 4584–4589. doi:10.1016/j.disc.2009.02.024

[10] R. Sen Gupta, M.K. Sen and S. Ghosh, *A variation of zero-divisor graphs*, Discuss. Math. Gen. Algebra Appl. **35** (2015) 159–176. doi:10.7151/dmgaa.1238

[11] K. Samei, *The zero divisor graph of a reduced ring*, J. Pure App. Alg. **209** (2007) 813–821.
doi:10.1016/j.jpaa.2006.08.008.

[12] S.K. Nimbhorkar, *Coloring of Rickart *-rings*, Trends in Theory of Rings and Modules, S. Tariq Rizvi and S.M.A. Zaidi (Eds.) Anamya Publishers (New Delhi 2005), 91–95.

[13] S.K. Nimbhorkar, M.P. Wasadikar and L. DeMeyer, *Coloring of semilattices*, Ars Comb. **12** (2007) 97–104.

[14] Avinash Patil and B.N. Waphare, *On the Zero-divisor graph of Rickart ∗-rings*, Asian-Eur. J. Math. **10** (2017) 1750015 (17 pages).
doi:10.1142/S1793557117500152

[15] Avinash Patil, B.N. Waphare, V.V. Joshi and H.Y. Paurali, *Zero-divisor graphs of lower dismantlable lattices*-I, Math. Slovaca **67** (2017) 1–12.

[16] Avinash Patil, B.N. Waphare and V.V. Joshi, *Zero-divisor graphs of lower dismantlable lattices*-II, to appear in Math. Slovaca.

[17] Avinash Patil and B.N. Waphare, *The zero-divisor graph of a ring with involution*, J. Algebra Appl. (to appear).

[18] S.P. Redmond, *The zero-divisor graph of a non-commutative ring*, Int. J. Commut. Rings **1** (2002) 203–211.

[19] N.K. Thakare and S.K. Nimbhorkar, *Prime strict ideals in Rickart *-rings*, Ind. J. Pure appl. Math. **22** (1991) 63–72.

[20] D.B. West, *Introduction to Graph Theory*, Second Edition (Prentice-Hall of India, New Delhi, 2002).

Received 28 June 2016
1st Revised 17 October 2016
2st Revised 23 January 2017