

## FOLDING THEORY OF IMPLICATIVE AND OBSTINATE IDEALS IN BL-ALGEBRAS

AKBAR PAAD

*Department of Mathematics*  
*University of Bojnord, Bojnord, Iran*

**e-mail:** akbar.paad@gmail.com

### Abstract

In this paper, the concepts of  $n$ -fold implicative ideals and  $n$ -fold obstinate ideals in  $BL$ -algebras are introduced. With respect to this concepts, some related results are given. In particular, it is proved that an ideal is an  $n$ -fold implicative ideal if and only if is an  $n$ -fold Boolean ideal. Also, it is shown that a  $BL$ -algebra is an  $n$ -fold integral  $BL$ -algebra if and only if trivial ideal  $\{0\}$  is an  $n$ -fold obstinate ideal. Moreover, the relation between  $n$ -fold obstinate ideals and  $n$ -fold (integral) obstinate filters in  $BL$ -algebras are studied by using the set of complement elements. Finally, it is proved that ideal  $I$  of  $BL$ -algebra  $L$  is an  $n$ -fold obstinate ideal if and only if  $\frac{L}{I}$  is an  $n$ -fold obstinate  $BL$ -algebra.

**Keywords:**  $BL$ -algebra, ideal,  $n$ -fold implicative ideal,  $n$ -fold obstinate ideal.

**2010 Mathematics Subject Classification:** 03G25, 03G05, 06D35, 06E99.

### 1. INTRODUCTION

$BL$ -algebras are the algebraic structure for Hájek basic logic [7] in order to investigate many valued logic by algebraic means. His motivations for introducing  $BL$ -algebras were of two kinds. The first one was providing an algebraic counterpart of a propositional logic, called Basic Logic, which embodies a fragment common to some of the most important many-valued logics, namely Lukasiewicz Logic, Gödel Logic and Product Logic. This Basic Logic ( $BL$  for short) is proposed as "the most general" many-valued logic with truth values in  $[0, 1]$  and  $BL$ -algebras are the corresponding Lindenbaum-Tarski algebras. The second one was to provide an algebraic mean for the study of continuous t-norms (or triangular norms)

on  $[0, 1]$ . In 1958, Chang [1] introduced the concept of an *MV*-algebra which is one of the most classes of *BL*-algebras. Turunen [12] introduced the notion of an implicative filter and a Boolean filter in *BL*-algebras. Boolean filters are an important class of filters, because the quotient *BL*-algebra induced by these filters are Boolean algebras. The notion of (fuzzy) ideal has been introduced in many algebraic structures such as lattices, rings, *MV*-algebras. Ideal theory is very effective tool for studying various algebraic and logical systems. In the theory of *MV*-algebras, as various algebraic structures, the notion of ideal is at the center, while in *BL*-algebras, the focus has been on deductive systems also filters. The study of *BL*-algebras has experienced a tremendous growth over recent years and the main focus has been on filters. In 2013, Lele [6], introduced the notions of (Boolean, prime) ideals and analyzed the relationship between ideals and filters by using the set of complement elements. In 2017, Yang and Xin [11], introduced implicative ideals in *BL*-algebras and studied some characterizations of them by the pseudo implication operation and proved the implicative ideals coincide with Boolean ideals in *BL*-algebras.

This motivates us to introduce the notions of  $n$ -fold implicative and  $n$ -fold obstinate ideals in *BL*-algebras and investigate the relations among  $n$ -fold implicative ideals,  $n$ -fold obstinate ideals and the other ideals in *BL*-algebras. In particular, we prove that an ideal is an  $n$ -fold implicative ideal if and only if is an  $n$ -fold Boolean. Also, we prove that a *BL*-algebra is an  $n$ -fold integral *BL*-algebra if and only if trivial ideal  $\{0\}$  is an  $n$ -fold obstinate ideal. Moreover, we study relation between  $n$ -fold obstinate ideals and  $n$ -fold (integral) obstinate filters in *BL*-algebras by using the set of complement elements. Finally, we prove that ideal  $I$  of *BL*-algebra  $L$  is an  $n$ -fold obstinate ideal if and only if  $\frac{L}{I}$  is an  $n$ -fold obstinate *BL*-algebra.

## 2. PRELIMINARIES

In this section, we give some fundamental definitions and results. For more details, refer to the references.

**Definition** [7]. A *BL*-algebra is an algebra  $(L, \vee, \wedge, \odot, \rightarrow, 0, 1)$  of type  $(2, 2, 2, 2, 0, 0)$  such that

- (BL1)  $(L, \vee, \wedge, 0, 1)$  is a bounded lattice,
- (BL2)  $(L, \odot, 1)$  is a commutative monoid,
- (BL3)  $z \leq x \rightarrow y$  if and only if  $x \odot z \leq y$ , for all  $x, y, z \in L$ ,
- (BL4)  $x \wedge y = x \odot (x \rightarrow y)$ ,
- (BL5)  $(x \rightarrow y) \vee (y \rightarrow x) = 1$ .

We denote  $x^n = \overbrace{x \odot \cdots \odot x}^{n\text{-times}}$ , if  $n > 0$  and  $x^0 = 1$ , for all  $x, y \in L$ .

A  $BL$ -algebra  $L$  is called a Gödel algebra (1-fold implicative  $BL$ -algebra) if  $x^2 = x \odot x = x$ , for all  $x \in L$  and  $L$  is called an  $MV$ -algebra if  $(x^-)^- = x$ , for all  $x \in L$ , where  $x^- = x \rightarrow 0$ . A  $BL$ -algebra  $L$  is called a Boolean algebra if  $x \vee x^- = 1$ , for all  $x \in L$ .

**Proposition 1** [2, 3]. *In any  $BL$ -algebra the following hold:*

- (BL6)  $x \leq y$  if and only if  $x \rightarrow y = 1$ ,
- (BL7)  $y \leq x \rightarrow y$ , and  $x \odot y \leq x, y$ ,
- (BL8)  $x \leq y$  implies  $y \rightarrow z \leq x \rightarrow z$  and  $z \rightarrow x \leq z \rightarrow y$ ,
- (BL9)  $(x \rightarrow y)^{-} = x^{-} \rightarrow y^{-}$ ,
- (BL10)  $(x \odot y)^{-} = x^{-} \odot y^{-}$ ,
- (BL11)  $(x \odot y)^- = x \rightarrow y^-$ ,
- (BL12)  $x^{-} = x^{-}$ ,  $x \leq x^{-}$  and  $x \odot x^- = 0$ ,
- (BL13)  $x \rightarrow (y \rightarrow z) = (x \odot y) \rightarrow z$ ,
- (BL14)  $x \leq y$  implies  $y^- \leq x^-$ ,
- (BL15)  $x \leq y$  implies  $z \odot x \leq z \odot y$ ,
- (BL16)  $(x \wedge y)^{-} = x^{-} \wedge y^{-}$ , for all  $x, y, z \in L$ .

Note that by (BL13)  $(x \rightarrow (\cdots (x \rightarrow (x \rightarrow y))) \cdots) = x^n \rightarrow y$ , for all  $x, y \in L$ . The following theorems and definitions are from [4, 5, 8, 10] and we refer the reader to them, for more details.

**Definition.** Let  $L$  be a  $BL$ -algebra,  $n$  be a natural number and  $F$  be a nonempty subset of  $L$ . Then

- (i)  $F$  is called a *filter* of  $L$  if  $x \odot y \in F$ , for any  $x, y \in F$  and if  $x \in F$  and  $x \leq y$  then  $y \in F$ , for all  $x, y \in L$ . A proper filter  $F$  is called a *maximal filter* of  $L$  if it is not properly contained in any other proper filter of  $L$ .
- (ii)  $F$  is called an  *$n$ -fold implicative filter* of  $L$  if  $1 \in F$  and for all  $x, y, z \in L$ ,

$$x^n \rightarrow (y \rightarrow z) \in F \text{ and } x^n \rightarrow y \in F \text{ imply } x^n \rightarrow z \in F.$$

- (iii) A proper filter  $F$  is called an  *$n$ -fold obstinate filter* if for all  $x, y \in L$ ,

$$x, y \notin F \text{ imply } x^n \rightarrow y \in F \text{ and } y^n \rightarrow x \in F.$$

- (iv) A proper filter  $F$  is called an  *$n$ -fold integral filter* if for all  $x, y \in L$ ,

$$(x^n \odot y^n)^- \in F \text{ implies } (x^n)^- \in F \text{ or } (y^n)^- \in F.$$

**Definition** [10]. Let  $L$  be a  $BL$ -algebra and  $n$  be a natural number. Then

- (i)  $L$  is called an  $n$ -fold integral  $BL$ -algebra if for all  $x, y \in L$

$$x^n \odot y^n = 0 \text{ then } x^n = 0 \text{ or } y^n = 0.$$

- (ii)  $L$  is called an  $n$ -fold obstinate  $BL$ -algebra if  $L$  is an  $MV$ -algebra and  $x^n = 0$ , for all  $x \in L \setminus \{1\}$ .

**Definition** [6, 8, 9]. Let  $L$  be a  $BL$ -algebra and  $I$  be a nonempty subset of  $L$ . Then

- (i)  $I$  is called an *ideal* of  $L$ , if  $x \odot y := x^- \rightarrow y \in I$ , for any  $x, y \in I$  and if  $y \in I$  and  $x \leq y$  then  $x \in I$ , for all  $x, y \in L$ . The operation  $\odot$  is associative. Moreover, a set  $I$  containing 0 of  $L$  is an ideal if and only if for all  $x, y \in L$ ,  $x^- \odot y \in I$  and  $x \in I$  imply  $y \in I$ .
- (ii) A proper ideal  $I$  of  $L$  is called a *prime ideal* of  $L$  if  $x \wedge y \in I$  implies  $x \in I$  or  $y \in I$ , for all  $x, y \in L$ .
- (iii) A proper ideal  $I$  is called a *maximal ideal* of  $L$  if it is not properly contained in any other proper ideal of  $L$ .
- (iv) An ideal  $I$  of  $L$  is called a  *$n$ -fold Boolean ideal* if  $x^n \wedge (x^n)^- \in I$ , for all  $x \in L$  and an ideal  $I$  of  $L$  is called a *Boolean ideal* if  $x \wedge x^- \in I$ , for all  $x \in L$ .
- (v) An ideal  $I$  of  $L$  is called an  *$n$ -fold integral ideal*, if for all  $x, y \in L$ ,

$$(x \odot y)^n \in I \text{ implies } x^n \in I \text{ or } y^n \in I.$$

Let  $L$  be a  $BL$ -algebra, we define the pseudo implication operation  $\rightarrow$  by  $x \rightarrow y := x \odot y^-$ , for any  $x, y \in L$ . It is easy to see that  $z \leq x \odot y$  if and only if  $z \rightarrow x \leq y$ .

Moreover, we denote  $x^n_{\odot} = \overbrace{x \odot \cdots \odot x}^{n\text{-times}}$ , when  $n$  is a natural number.

**Lemma 2** [11]. Let  $L$  be a  $BL$ -algebra, for any  $x, y, z \in L$ , we have:

- (i)  $x \leq y$  implies  $z \rightarrow y \leq z \rightarrow x$  and  $x \rightarrow z \leq y \rightarrow z$ ,
- (ii)  $(x \rightarrow y) \rightarrow z = (x \rightarrow z) \rightarrow y = x \rightarrow (y \odot z)$ ,
- (iii)  $x \rightarrow 0 = x$ ,  $0 \rightarrow x = 0$ ,  $x \rightarrow x = 0$ ,
- (iv)  $(x \rightarrow z) \rightarrow (y \rightarrow z) \leq x \rightarrow y$ ,
- (v)  $(x \rightarrow z) \leq (y \rightarrow z) \odot (x \rightarrow y)$ ,
- (vi)  $x \leq x \odot x$ .

**Lemma 3** [11]. Let  $I$  be a nonempty subset of a  $BL$ -algebra  $L$ . Then  $I$  is an ideal of  $L$  if and only if it satisfies:

- (i)  $0 \in I$ ,
- (ii) for any  $x, y \in L$ , if  $x \rightarrow y \in I$  and  $y \in I$ , then  $x \in I$ .

**Lemma 4** [11]. Let  $I$  be an ideal of BL-algebra  $L$ . Then the following hold: for any  $x, y, z \in L$

- (i)  $x \rightarrow y \in I$  if and only if  $y^- \rightarrow x^- \in I$ .
- (ii)  $x \rightarrow y \in I$  if and only if  $x^{--} \rightarrow y \in I$ .
- (iii)  $(y \rightarrow x^-) \rightarrow z \in I$  if and only if  $(z^- \rightarrow y^-) \rightarrow x^- \in I$ .
- (iv)  $x \in I$  if and only if  $x^{--} \in I$ .

**Theorem 5** [11]. Let  $P$  be a proper ideal of BL-algebra  $L$ . Then  $P$  is a prime ideal if and only if  $x \rightarrow y \in P$  or  $y \rightarrow x \in P$ , for all  $x, y \in L$ .

**Definition** [6]. Let  $L$  be a BL-algebra and  $X$  any subset of  $L$ . Then the set of complement elements (with respect to  $X$ ) is denoted by  $N(X)$  and is defined by

$$N(X) = \{x \in L \mid x^- \in X\}.$$

**Theorem 6** [6]. Let  $I$  be an ideal of BL-algebra  $L$ . Then the binary relation  $\equiv_I$  on  $L$  which is defined by

$$x \equiv_I y \text{ if and only if } x^- \odot y \in I \text{ and } y^- \odot x \in I$$

is a congruence relation on  $L$ . Define  $\cdot, \rightarrow, \sqcup, \sqcap$  on  $\frac{L}{I}$ , the set of all congruence classes of  $L$ , as follows:

$$[x] \cdot [y] = [x \odot y], [x] \rightarrow [y] = [x \rightarrow y]$$

$$[x] \sqcup [y] = [x \vee y], [x] \sqcap [y] = [x \wedge y].$$

Then  $(\frac{L}{I}, \cdot, \rightarrow, \sqcup, \sqcap, [0], [1])$  is a BL-algebra which is called quotient BL-algebra with respect to  $I$ . In addition, it is clear  $[x]^{--} = [x]$ , for all  $x \in L$ . Consequently, the quotient BL-algebra via any ideal is always an MV-algebra.

**Theorem 7** [9]. Let  $I$  be an ideal of  $L$ . Then the following conditions are equivalent:

- (i)  $I$  is an  $n$ -fold integral ideal of  $L$ ,
- (ii)  $I$  is a maximal and  $n$ -fold Boolean ideal of  $L$ ,
- (iii)  $I$  is a prime and  $n$ -fold Boolean ideal of  $L$ ,
- (iv)  $I$  is a proper ideal and for all  $x \in L$ ,  $x^n \in I$  or  $(x^n)^- \in I$ .

**Theorem 8** [9]. Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold integral ideal if and only if  $N(I)$  is an  $n$ -fold obstinate filter of  $L$ .

**Theorem 9** [9]. *Let  $F$  be a proper filter of  $L$ . Then  $F$  is an  $n$ -fold integral filter if and only if  $N(F)$  is an  $n$ -fold integral ideal of  $L$ .*

**Theorem 10** [9]. *In any  $BL$ -algebra  $L$ , the following conditions are equivalent:*

- (i)  $\{0\}$  is an  $n$ -fold integral ideal of  $L$ ,
- (ii) any ideal of  $L$  is an  $n$ -fold integral ideal,
- (iii)  $L$  is an  $n$ -fold integral  $BL$ -algebra.

**Theorem 11** [9]. *Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold integral ideal of  $L$  if and only if  $\frac{L}{I}$  is an  $n$ -fold obstinate  $BL$ -algebra.*

**Theorem 12** [9]. *Let  $L$  be a Boolean algebra or a Gödel algebra. Then every ideal of  $L$  is implicative.*

From now on, in this paper  $(L, \wedge, \vee, \odot, \rightarrow, 0, 1)$  (or simply)  $L$  is a  $BL$ -algebra, unless otherwise stated.

### 3. $N$ -FOLD IMPLICATIVE IDEALS IN $BL$ -ALGEBRAS

In this section we introduce two new class of ideals in  $BL$ -algebras that called  $n$ -fold implicative ideals and we give some related results.

**Definition.** A nonempty subset  $I$  of  $L$  is called an  $n$ -fold implicative ideal if it satisfies:

- (i)  $0 \in I$ ,
- (ii)  $(x \rightarrow y) \rightarrow z_{\odot}^n \in I$  and  $y \rightarrow z_{\odot}^n \in I$  imply  $x \rightarrow z_{\odot}^n \in I$ , for all  $x, y, z \in L$ .

An 1-fold implicative ideal is called an implicative ideal of  $L$ .

**Example 13** [6]. Let  $L = \{0, a, b, c, d, e, f, 1\}$  be such that  $0 < a < b < c < 1$ ,  $0 < d < e < f < 1$ ,  $a < e$  and  $b < f$ . Define  $\odot$  and  $\rightarrow$  as follows:

Table 1

$\odot$	0	a	b	c	d	e	f	1
0	0	0	0	0	0	0	0	0
a	0	a	a	a	0	a	a	a
b	0	a	a	b	0	a	a	b
c	0	a	b	c	0	a	b	c
d	0	0	0	0	d	d	d	d
e	0	a	a	a	d	e	e	e
f	0	a	a	b	d	e	e	f
1	0	a	b	c	d	e	f	1

Table 2

$\rightarrow$	0	a	b	c	d	e	f	1
0	1	1	1	1	1	1	1	1
a	d	1	1	1	d	1	1	1
b	d	f	1	1	d	f	1	1
c	d	e	f	1	d	e	f	1
d	c	c	c	c	1	1	1	1
e	0	c	c	c	d	1	1	1
f	0	b	c	c	d	f	1	1
1	0	a	b	c	d	e	f	1

Then  $(L, \wedge, \vee, \odot, \rightarrow, 0, 1)$  is a  $BL$ -algebra. Let  $I = \{0, d\}$ . Then  $I$  is a 2-fold implicative ideal of  $L$ .

**Proposition 14.** *Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then  $I$  is an ideal of  $L$ .*

**Proof.** Suppose that  $I$  is an  $n$ -fold implicative ideal of  $L$  and  $x, y \in L$ . If  $x \rightarrow y \in I$  and  $y \in I$ , then  $(x \rightarrow y) \rightarrow 0_{\odot}^n = x \rightarrow y \in I$  and  $y \rightarrow 0_{\odot}^n = y \in I$ . By hypothesis  $x = x \rightarrow 0_{\odot}^n \in I$ , hence  $I$  is an ideal of  $L$ . ■

The following example shows that the converse of Proposition 14, does not hold in general.

**Example 15** [6]. Let  $L = \{0, a, b, 1\}$ , where  $0 < a < b < 1$ . Let  $x \wedge y = \min\{x, y\}$ ,  $x \vee y = \max\{x, y\}$  and operations  $\odot$  and  $\rightarrow$  are defined as the following tables:

Table 3

$\odot$	0	a	b	1
0	0	0	0	0
a	0	0	0	a
b	0	0	a	b
1	0	a	b	1

Table 4

$\rightarrow$	0	a	b	1
0	1	1	1	1
a	b	1	1	1
b	a	b	1	1
1	0	a	b	1

Then  $(L, \vee, \wedge, \odot, \rightarrow, 0, 1)$  is a  $BL$ -algebra. Now, let  $I = \{0\}$ . Then  $I$  is an ideal of  $L$  and since  $(1 \rightarrow b) \rightarrow b = b^- \odot b^- = a \odot a = 0 \in I$ ,  $b \rightarrow b = b \odot b^- = b \odot a = 0 \in I$  and  $1 \rightarrow b = 1 \odot b^- = a \notin I$ , then  $I$  is not a 1-fold implicative ideal of  $L$ .

**Theorem 16.** *Let  $I$  be an ideal of  $L$ . Then the following conditions are equivalent:*

- (i)  $I$  is an  $n$ -fold implicative ideal of  $L$ ,
- (ii) for any  $a \in L$ , the set  $I_{a_{\odot}^n} := \{x \in L \mid x \rightarrow a_{\odot}^n \in I\}$  is an ideal of  $L$ .

**Proof.** (i)  $\Rightarrow$  (ii) Suppose that  $I$  is an  $n$ -fold implicative ideal of  $L$  and  $a \in L$ . For any  $x, y \in L$ , if  $x \rightarrow y \in I_{a_{\odot}^n}$  and  $y \in I_{a_{\odot}^n}$ , then  $(x \rightarrow y) \rightarrow a_{\odot}^n \in I$  and  $y \rightarrow a_{\odot}^n \in I$ , hence  $x \rightarrow a_{\odot}^n \in I$ , and so  $x \in I_{a_{\odot}^n}$ . Moreover, since  $0 \rightarrow a_{\odot}^n = 0 \odot (a_{\odot}^n)^- = 0 \in I$ , we obtain  $0 \in I_{a_{\odot}^n}$ . Therefore,  $I_{a_{\odot}^n}$  is an ideal of  $L$ .

(ii)  $\Rightarrow$  (i) Suppose that  $I_{a_{\odot}^n}$  is an ideal of  $L$ , for any  $a \in L$ . For any  $x, y, z \in L$ , if  $(x \rightarrow y) \rightarrow z_{\odot}^n \in I$  and  $y \rightarrow z_{\odot}^n \in I$ , then  $x \rightarrow y \in I_{z_{\odot}^n}$  and  $y \in I_{z_{\odot}^n}$ . Now, since  $I_{z_{\odot}^n}$  is an ideal of  $L$ , we have  $x \in I_{z_{\odot}^n}$ , and so  $x \rightarrow z_{\odot}^n \in I$ . Therefore,  $I$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Theorem 17.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then for any  $a \in L$ ,  $I_{a_{\odot}^n}$  is the least ideal of  $L$  containing  $I$  and  $a$ .

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$  and  $a \in L$ . Then by Theorem 16,  $I_{a_{\odot}^n}$  is an ideal of  $L$  and by (BL7), for any  $x \in I$ ,  $x \rightarrow a_{\odot}^n = x \odot (a_{\odot}^n)^- \leq x$ , we get  $x \rightarrow a_{\odot}^n \in I$ , and so  $x \in I_{a_{\odot}^n}$ . Hence  $I \subseteq I_{a_{\odot}^n}$ . Moreover, by (BL7), (BL12), (BL14) and (BL15),

$$\begin{aligned} a \rightarrow a_{\odot}^n &= a \rightarrow (a_{\odot}^{n-1} \odot a) = a \odot (a_{\odot}^{n-1} \odot a)^- \\ &= a \odot \left( (a_{\odot}^{n-1})^- \rightarrow a \right)^- \leq a \odot a^- = 0. \end{aligned}$$

Hence,  $a \rightarrow a_{\odot}^n = 0 \in I$ , and so  $a \in I_{a_{\odot}^n}$ . Now, if  $J$  is an ideal of  $L$  containing  $I$  and  $a$ , then for any  $x \in I_{a_{\odot}^n}$ , we get that  $x \rightarrow a_{\odot}^n \in I \subseteq J$ . Since  $J$  is an ideal of

$L$  and  $a \in J$ , we have  $a_{\odot}^n = \overbrace{a \odot \cdots \odot a}^{n\text{-times}} \in J$  and so  $x \in J$ . Therefore,  $I_{a_{\odot}^n} \subseteq J$  and so  $I_{a_{\odot}^n}$  is the least ideal of  $L$  containing  $I$  and  $a$ . ■

**Theorem 18.** Let  $I$  be a nonempty subset of  $L$ . Then the following conditions are equivalent:

- (i)  $I$  is an  $n$ -fold implicative ideal of  $L$ ,
- (ii)  $I$  is an ideal of  $L$  and for any  $x, y \in L$ ,  $x \rightarrow y_{\odot}^{n+1} \in I$  implies  $x \rightarrow y_{\odot}^n \in I$ ,
- (iii)  $I$  is an ideal of  $L$  and for any  $x, y, z \in L$ ,  $(x \rightarrow y) \rightarrow z_{\odot}^n \in I$  implies  $(x \rightarrow z_{\odot}^n) \rightarrow (y \rightarrow z_{\odot}^n) \in I$ ,
- (iv)  $0 \in I$ , and if  $(x \rightarrow y_{\odot}^{n+n}) \rightarrow z \in I$  and  $z \in I$ , then  $x \rightarrow y_{\odot}^n \in I$ , for any  $x, y, z \in L$ .
- (v)  $0 \in I$ , and if  $(x \rightarrow y_{\odot}^{n+1}) \rightarrow z \in I$  and  $z \in I$ , then  $x \rightarrow y_{\odot}^n \in I$ , for any  $x, y, z \in L$ .

**Proof.** (i)  $\Rightarrow$  (ii) Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then by Proposition 14,  $I$  is an ideal of  $L$ . Now, if  $x \rightarrow y_{\odot}^{n+1} \in I$ , for  $x, y \in L$ , then by Lemma



2(ii),  $(x \multimap y) \multimap y_{\mathcal{O}}^n = x \multimap y_{\mathcal{O}}^{n+1} \in I$  and since by Lemma 2(ii) and (iii),  $y \multimap y_{\mathcal{O}}^n = y \multimap y \odot y_{\mathcal{O}}^{n-1} = (y \multimap y) \multimap y_{\mathcal{O}}^{n-1} = 0 \multimap y_{\mathcal{O}}^{n-1} = 0 \in I$ , we get  $x \multimap y_{\mathcal{O}}^n \in I$ .

(ii)  $\Rightarrow$  (iii) Assume that (ii) holds. Let  $x, y, z \in L$  and  $(x \multimap y) \multimap z_{\mathcal{O}}^n \in I$ . By Lemma 2(i), (ii) and (iv),

$$((x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^n) \multimap z_{\mathcal{O}}^n = ((x \multimap z_{\mathcal{O}}^n) \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^n \leq (x \multimap y) \multimap z_{\mathcal{O}}^n.$$

Then  $((x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^n) \multimap z_{\mathcal{O}}^n \in I$ , and so  $(x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^{(n+n-1)+1} \in I$  and by hypothesis  $(x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^{(n+n-1)} \in I$ . By continuing this process we get that  $(x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^{(n+1)} \in I$ . Hence,  $(x \multimap (y \multimap z_{\mathcal{O}}^n)) \multimap z_{\mathcal{O}}^n \in I$ . Therefore,  $(x \multimap z_{\mathcal{O}}^n) \multimap (y \multimap z_{\mathcal{O}}^n) \in I$ .

(iii)  $\Rightarrow$  (iv) Assume that (iii) holds. Obviously,  $0 \in I$ . Let  $(x \multimap y_{\mathcal{O}}^{n+n}) \multimap z \in I$  and  $z \in I$ , for  $x, y, z \in L$ . Since  $I$  is an ideal of  $L$ , we have  $x \multimap y_{\mathcal{O}}^{n+n} \in I$ . Now, since by Lemma 2(ii),  $(x \multimap y_{\mathcal{O}}^n) \multimap y_{\mathcal{O}}^n = x \multimap y_{\mathcal{O}}^{n+n} \in I$ , then by (iii),  $(x \multimap y_{\mathcal{O}}^n) \multimap (y_{\mathcal{O}}^n \multimap y_{\mathcal{O}}^n) \in I$  and since  $y_{\mathcal{O}}^n \multimap y_{\mathcal{O}}^n = 0$ , then  $x \multimap y_{\mathcal{O}}^n \in I$ .

(iv)  $\Rightarrow$  (i) Suppose that (iv) is valid. Firstly, we show that  $I$  is an ideal of  $L$ . For any  $x, y \in L$ , if  $x \multimap y \in L$  and  $y \in I$ , then

$$\begin{aligned} (x \multimap 0_{\mathcal{O}}^{n+n}) \multimap y &= (\cdots (x \multimap \overbrace{0 \multimap 0}^{(n+n)\text{-times}}) \cdots \multimap 0) \cdots \multimap y \\ &= (\cdots (x \multimap \overbrace{0 \multimap 0}^{(2n-1)\text{-times}}) \cdots \multimap 0) \cdots \multimap y \\ &\vdots \\ &= x \multimap y \in I. \end{aligned}$$

And since  $y \in I$ , it follows that by (iv),  $x = x \multimap 0_{\mathcal{O}}^n \in I$ . Hence,  $I$  is an ideal of  $L$ . Now, let  $(x \multimap y) \multimap z_{\mathcal{O}}^n \in I$  and  $y \multimap z_{\mathcal{O}}^n \in I$ , for  $x, y, z \in L$ . Then by Lemma 2(ii) and (iv),

$$((x \multimap z_{\mathcal{O}}^n) \multimap z_{\mathcal{O}}^n) \multimap (y \multimap z_{\mathcal{O}}^n) \leq ((x \multimap z_{\mathcal{O}}^n) \multimap y = (x \multimap y) \multimap z_{\mathcal{O}}^n).$$

And since  $(x \multimap y) \multimap z_{\mathcal{O}}^n \in I$ , we obtain  $((x \multimap z_{\mathcal{O}}^n) \multimap z_{\mathcal{O}}^n) \multimap (y \multimap z_{\mathcal{O}}^n) \in I$ , hence  $(x \multimap z_{\mathcal{O}}^{n+n}) \multimap (y \multimap z_{\mathcal{O}}^n) \in I$ . Now, since  $y \multimap z_{\mathcal{O}}^n \in I$ , so by (iv),  $x \multimap z_{\mathcal{O}}^n \in I$ . Therefore,  $I$  is an  $n$ -fold implicative ideal of  $L$ .

(iv)  $\Rightarrow$  (v) Let  $(x \multimap y_{\mathcal{O}}^{n+1}) \multimap z \in I$  and  $z \in I$ , for  $x, y, z \in L$ . Then by the similarly proof ((iv)  $\Rightarrow$  (i)),  $I$  is an ideal of  $L$ . Moreover, since  $y_{\mathcal{O}}^{n+1} \leq y_{\mathcal{O}}^{n+n}$ , we conclude that by Lemma 2(i),  $x \multimap y_{\mathcal{O}}^{n+n} \leq x \multimap y_{\mathcal{O}}^{n+1}$ . Hence,  $(x \multimap y_{\mathcal{O}}^{n+n}) \multimap z \leq x \multimap (y_{\mathcal{O}}^{n+1}) \multimap z$  and since  $(x \multimap y_{\mathcal{O}}^{n+1}) \multimap z \in I$ , we get  $(x \multimap y_{\mathcal{O}}^{n+n}) \multimap z \in I$ . Now, since  $z \in I$ , we have by (iv),  $x \multimap y_{\mathcal{O}}^n \in I$ .

(v)  $\Rightarrow$  (ii) By the similarly proof ((iv)  $\Rightarrow$  (i)),  $I$  is an ideal of  $L$ . Now, if  $x \multimap y_{\mathcal{O}}^{n+1} \in I$ , then  $(x \multimap y_{\mathcal{O}}^{n+1}) \multimap 0 \in I$  and so by (v),  $x \multimap y_{\mathcal{O}}^n \in I$ . ■

**Theorem 19.** *Let  $I \subseteq J$ , where  $I$  and  $J$  be two ideals of  $L$  and  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then  $J$  is an  $n$ -fold implicative ideal, too.*

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$ ,  $I \subseteq J$  and  $(x \rightarrow y) \rightarrow z_{\odot}^n \in J$ , for  $x, y, z \in L$ . Denote  $u = (x \rightarrow y) \rightarrow z_{\odot}^n$ . Then by Lemma 2(i) and (iii),  $((x \rightarrow u) \rightarrow y) \rightarrow z_{\odot}^n = ((x \rightarrow y) \rightarrow z_{\odot}^n) \rightarrow u = u \rightarrow u = 0 \in I$ . Since  $I$  is an  $n$ -fold implicative ideal of  $L$ , it follows by Theorem 18,

$$((x \rightarrow u) \rightarrow z_{\odot}^n) \rightarrow (y \rightarrow z_{\odot}^n) \in I \subseteq J.$$

Hence, by Lemma 2(ii),  $((x \rightarrow z_{\odot}^n) \rightarrow (y \rightarrow z_{\odot}^n)) \rightarrow u \in J$  and since  $J$  is an ideal of  $L$  and  $u \in J$ , we have  $(x \rightarrow z_{\odot}^n) \rightarrow (y \rightarrow z_{\odot}^n) \in J$ . Therefore, by Theorem 18,  $J$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Lemma 20.** *For any BL-algebra  $L$  and  $x, y \in L$ ,*

- (i)  $(x_{\odot}^n)^- = (x^-)^n$ .
- (ii)  $(x^n)^- = (x^-)_{\odot}^n$ .
- (iii)  $(x \odot y)^{- -} = x^{- -} \odot y^{- -}$ .
- (iv)  $(x \odot y)^- = x^- \rightarrow y^{- -}$ .

**Proof.** (i) For any  $x \in L$ , by (BL9), (BL11) and (BL12),

$$(x^- \rightarrow x)^{- -} = x^{- - -} \rightarrow x^{- -} = x^- \rightarrow x^{- -} = (x^- \odot x^-)^-.$$

Then  $(x^- \rightarrow x)^- = (x^- \rightarrow x)^{- - -} = (x^- \odot x^-)^{- -} = x^{- - -} \odot x^{- - -} = x^- \odot x^-$ . Hence,

$$(x \odot x)^- = (x^- \rightarrow x)^- = x^- \odot x^-.$$

Now, since the operation  $\odot$  is associative, we get

$$\begin{aligned} (x_{\odot}^n)^- &= \overbrace{(x \odot \cdots \odot x)}^{n\text{-times}}^- \\ &= \overbrace{((x \odot \cdots \odot x) \odot x)}^{(n-1)\text{-times}}^- \\ &= \overbrace{(x \odot \cdots \odot x)}^{(n-1)\text{-times}}^- \odot x^- \\ &\vdots \\ &= (x \odot x)^- \odot \overbrace{(x^- \odot \cdots \odot x^-)}^{(n-2)\text{-times}} \\ &= \overbrace{x^- \odot \cdots \odot x^-}^{n\text{-times}} \\ &= (x^-)^n. \end{aligned}$$

(ii) For any  $x \in L$ , by (BL9), (BL11) and (BL12),

$$\begin{aligned}
 (x \odot x)^- &= ((x \odot x)^-)^{-} \\
 &= (x \rightarrow x^-)^{-} \\
 &= x^{-} \rightarrow x^{-} \\
 &= x^{-} \rightarrow x^- \\
 &= x^- \odot x^-.
 \end{aligned}$$

Now,

$$\begin{aligned}
 (x^n)^- &= \overbrace{(x \odot \cdots \odot x)^-}^{n\text{-times}} \\
 &= \overbrace{((x \odot \cdots \odot x) \odot x)^-}^{(n-1)\text{-times}} \\
 &= \overbrace{(x \odot \cdots \odot x)^-}^{(n-1)\text{-times}} \odot x^- \\
 &\vdots \\
 &= (x \odot x)^- \odot \overbrace{(x^- \odot \cdots \odot x^-)}^{(n-2)\text{-times}} \\
 &= \overbrace{x^- \odot \cdots \odot x^-}^{n\text{-times}} \\
 &= (x^-)_{\odot}^n.
 \end{aligned}$$

(iii) Let  $x, y \in L$ . Then by the definition  $\odot$  and (BL9),  $(x \odot y)^{-} = (x^- \rightarrow y)^{-} = x^{-} \rightarrow y^- = x^- \odot y^-$ .

(iv) Let  $x, y \in L$ . Then by the definition  $\odot$ ,  $(x \odot y)^- = (x^- \rightarrow y)^-$ . Now, by (BL9), (BL11) and (BL12),

$$\begin{aligned}
 ((x^- \rightarrow y)^-)^- &= (x^- \rightarrow y)^{-} \\
 &= x^{-} \rightarrow y^- \\
 &= x^- \rightarrow y^- \\
 &= (x^- \odot y^-)^-.
 \end{aligned}$$

And by (BL10) and (BL12),

$$\begin{aligned}
 (x^- \rightarrow y)^- &= ((x^- \rightarrow y)^-)^{-} \\
 &= ((x^- \odot y^-)^-)^- \\
 &= x^- \odot y^- \\
 &= x^- \odot y^- \\
 &= x^- \rightarrow y^-.
 \end{aligned}$$

Therefore,  $(x \odot y)^- = x^- \rightarrow y^-$ . ■

**Theorem 21.** *Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold implicative ideal of  $L$  if and only if it satisfies the condition*

**(n-PI):**  $(y \rightarrow (x^n)^-) \rightarrow z \in I$  and  $x^n \rightarrow y \in I$  imply  $x^n \rightarrow z \in I$ , for any  $x, y, z \in L$ .

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . For any  $x, y, z \in L$ , let  $(y \rightarrow (x^n)^-) \rightarrow z \in I$  and  $x^n \rightarrow y \in I$ . Then by Lemma 4(i) and (iii),  $(z^- \rightarrow y^-) \rightarrow (x^n)^- \in I$  and  $y^- \rightarrow (x^n)^- \in I$ . Now, by Lemma 20(ii),  $(z^- \rightarrow y^-) \rightarrow (x^-)^n_{\odot} \in I$  and  $y^- \rightarrow (x^-)^n_{\odot} \in I$  and since  $I$  is an  $n$ -fold implicative ideal of  $L$ , we have  $z^- \rightarrow (x^-)^n_{\odot} \in I$ . Now, by Lemma 4(i),  $((x^-)^n_{\odot})^- \rightarrow z^{--} \in I$  and so by Lemma 20(i),  $(x^{--})^n \rightarrow z^{--} \in I$ . Moreover, since by (BL12) and (BL15),  $x^n \leq (x^{--})^n$ , it follows that by Lemma 2(i),  $x^n \rightarrow z^{--} \leq (x^{--})^n \rightarrow z^{--}$  and so  $x^n \rightarrow z^{--} \in I$ . Hence,  $x^n \odot (z^{--})^- \in I$  and so  $x^n \odot z^- \in I$ . Therefore,  $x^n \rightarrow z \in I$ .

Conversely, let  $I$  satisfy the condition **(n-PI)** and  $(x \rightarrow y) \rightarrow z^n_{\odot} \in I$ ,  $y \rightarrow z^n_{\odot} \in I$ , for  $x, y, z \in L$ . Then by Lemma 2(ii),  $x \rightarrow y \odot z^n_{\odot} \in I$  and so by Lemma 4(i),  $(y \odot z^n_{\odot})^- \rightarrow x^- \in I$  and  $(z^n_{\odot})^- \rightarrow y^- \in I$ . Now, by Lemma 20(iv),  $(y^- \rightarrow ((z^n_{\odot})^-)^-) \rightarrow x^- \in I$  and so Lemma 20(i),  $(y^- \rightarrow ((z^n_{\odot})^-)^-) \rightarrow x^- \in I$  and since  $(z^n_{\odot})^- \rightarrow y^- \in I$ , we get by condition **(n-PI)**,  $(z^-)^n \rightarrow x^- \in I$ . Hence, by Lemma 20(ii),  $(z^n_{\odot})^- \rightarrow x^-$ , and so by Lemma 4(i),  $x \rightarrow z^n_{\odot} \in I$ . Therefore,  $I$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Theorem 22.** *Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then  $I$  is an  $(n+1)$ -fold implicative ideal of  $L$ .*

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$  and  $x \rightarrow y^{n+2}_{\odot} \in I$ , for  $x, y \in L$ . Then by Lemma 2(ii),

$$(x \rightarrow y) \rightarrow y^{n+1}_{\odot} = x \rightarrow y \odot y^{n+1}_{\odot} = x \rightarrow y^{n+2}_{\odot} \in I.$$

Now, by Theorem 18,  $(x \rightarrow y) \rightarrow y^n_{\odot} \in I$  and so  $x \rightarrow y^{n+1}_{\odot} = (x \rightarrow y) \rightarrow y^n_{\odot} \in I$ . Hence, by Theorem 18,  $I$  is an  $(n+1)$ -fold implicative ideal of  $L$ . ■

**Theorem 23.** *Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold implicative ideal of  $L$  if and only if  $x^{2n} \rightarrow x^n_{\odot} \in I$ , for any  $x \in L$ .*

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$  and  $x \in L$ . Since by Lemma 2(ii),  $(x^{2n} \rightarrow x^n_{\odot}) \rightarrow x^n_{\odot} = x^{2n} \rightarrow x^n_{\odot} \odot x^n_{\odot} = x^{2n} \rightarrow x^{2n}_{\odot} = 0 \in I$ , and  $x^n_{\odot} \rightarrow x^n_{\odot} = 0 \in I$ , we get  $x^{2n} \rightarrow x^n_{\odot} \in I$ .

Conversely, suppose that for any  $x \in L$ ,  $x^{2n} \rightarrow x^n_{\odot} \in I$ , and  $(x \rightarrow y) \rightarrow z^n_{\odot} \in I$ ,  $y \rightarrow z^n_{\odot} \in I$ , for  $x, y, z \in L$ . Then by Lemma 2(ii) and (iv),  $((x \rightarrow z^n_{\odot}) \rightarrow z^n_{\odot}) \rightarrow (y \rightarrow z^n_{\odot}) \leq (x \rightarrow z^n_{\odot}) \rightarrow y = (x \rightarrow y) \rightarrow z^n_{\odot}$ . Since  $(x \rightarrow y) \rightarrow z^n_{\odot} \in I$  and  $I$  is an ideal of  $L$ , we have

$$((x \rightarrow z^n_{\odot}) \rightarrow z^n_{\odot}) \rightarrow (y \rightarrow z^n_{\odot}) \in I.$$

And since  $y \rightarrow z_{\odot}^n \in I$ , by Lemma 3,

$$(x \rightarrow z_{\odot}^n) \rightarrow z_{\odot}^n \in I.$$

Moreover, by Lemma 2(v),

$$x \rightarrow z_{\odot}^n \leq (z_{\odot}^n \odot z_{\odot}^n \rightarrow z_{\odot}^n) \odot (x \rightarrow z_{\odot}^n \odot z_{\odot}^n).$$

And since  $x \rightarrow z_{\odot}^n \odot z_{\odot}^n = (x \rightarrow z_{\odot}^n) \rightarrow z_{\odot}^n \in I$  and by hypothesis  $z_{\odot}^n \odot z_{\odot}^n \rightarrow z_{\odot}^n = z_{\odot}^{2n} \rightarrow z_{\odot}^n \in I$ , we have  $(z_{\odot}^n \odot z_{\odot}^n \rightarrow z_{\odot}^n) \odot (x \rightarrow z_{\odot}^n \odot z_{\odot}^n) \in I$ . Hence  $x \rightarrow z_{\odot}^n \in I$ . Therefore,  $I$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Theorem 24.** *Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold implicative ideal of  $L$  if and only if  $I$  is an  $n$ -fold Boolean ideal of  $L$ .*

**Proof.** Let  $I$  be an  $n$ -fold implicative ideal of  $L$ . Then by Theorem 22,  $x_{\odot}^{2n} \rightarrow x_{\odot}^n \in I$ , for any  $x \in L$ . By Lemma 2(ii),

$$\begin{aligned} x_{\odot}^{2n} \rightarrow x_{\odot}^n &= x_{\odot}^n \odot x_{\odot}^n \rightarrow x_{\odot}^n \\ &= ((x_{\odot}^n)^- \rightarrow x_{\odot}^n) \rightarrow x_{\odot}^n \\ &= ((x_{\odot}^n)^- \rightarrow x_{\odot}^n) \odot (x_{\odot}^n)^- \\ &= (x_{\odot}^n)^- \odot ((x_{\odot}^n)^- \rightarrow x_{\odot}^n) \\ &= (x_{\odot}^n)^- \wedge x_{\odot}^n \\ &= x_{\odot}^n \wedge (x_{\odot}^n)^-. \end{aligned}$$

Hence, for any  $x \in L$ ,  $x_{\odot}^n \wedge (x_{\odot}^n)^- \in I$  and since  $(x^-)_{\odot}^{2n} \rightarrow (x^-)_{\odot}^n \in I$ , by similar way  $(x^-)_{\odot}^n \wedge ((x^-)_{\odot}^n)^- \in I$ . Now, since by Lemma 20(i),  $((x^-)_{\odot}^n)^- = (x^{--})^n$ , then  $(x^-)_{\odot}^n \wedge (x^{--})^n \in I$  and since by (BL12),  $(x^-)_{\odot}^n \wedge x^n \leq (x^-)_{\odot}^n \wedge (x^{--})^n$ , we get  $(x^-)_{\odot}^n \wedge x^n \in I$ . Moreover, by Lemma 4(iv),  $((x^-)_{\odot}^n \wedge x^n)^{-} \in I$ . Hence, applying (BL16), we have  $((x^-)_{\odot}^n)^{-} \wedge (x^n)^{-} \in I$ . Now, by Lemma 20(i),  $((x^-)_{\odot}^n)^{-} = (((x^-)_{\odot}^n)^-)^- = ((x^{--})^n)^- = ((x^n)^{-})^- = (x^n)^-$ . Hence,

$$(x^n)^- \wedge (x^n)^{-} = ((x^-)_{\odot}^n)^{-} \wedge (x^n)^{-} \in I.$$

By (BL12),  $x^n \leq (x^n)^{-}$  and so  $(x^n)^- \wedge x^n \leq (x^n)^- \wedge (x^n)^{-}$  and since  $I$  is an ideal of  $L$ , we have  $(x^n)^- \wedge x^n \in I$ , for any  $x \in L$ . Therefore,  $I$  is an  $n$ -fold Boolean ideal of  $L$ .

Conversely, Let  $I$  be an  $n$ -fold Boolean ideal of  $L$ . Then for any  $x \in L$ ,  $((x^-)_{\odot}^n)^- \wedge (x^-)_{\odot}^n \in I$ . By Lemma 20(i),

$$((x_{\odot}^n)^-)^- \wedge (x_{\odot}^n)^- = ((x^-)_{\odot}^n)^- \wedge (x^-)_{\odot}^n \in I.$$

Since  $I$  is an ideal of  $L$  and by (BL12),

$$x_{\mathcal{O}}^n \wedge (x_{\mathcal{O}}^n)^- \leq (x_{\mathcal{O}}^n)^{- -} \wedge (x_{\mathcal{O}}^n)^- = ((x_{\mathcal{O}}^n)^-)^- \wedge (x_{\mathcal{O}}^n)^-,$$

we obtain  $x_{\mathcal{O}}^n \wedge (x_{\mathcal{O}}^n)^- \in I$ , and so  $x_{\mathcal{O}}^{2n} \rightarrow x_{\mathcal{O}}^n \in I$ . Therefore, by Theorem 23,  $I$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Theorem 25.** *In a BL-algebra  $L$ , the following conditions are equivalent:*

- (i) *any ideal  $I$  of  $L$  is an  $n$ -fold implicative,*
- (ii)  *$\{0\}$  is an  $n$ -fold implicative ideal of  $L$ ,*
- (iii) *for any  $a \in L$ , the set  $L(a) = \{x \in L \mid x \rightarrow a^n = 0\}$  is an ideal of  $L$ .*

**Proof.** (i)  $\Leftrightarrow$  (ii) It follows from Theorem 19.

(ii)  $\Leftrightarrow$  (iii) For any  $a, x, y \in L$ , if  $x \rightarrow y \in L(a)$  and  $y \in L(a)$ , then  $(x \rightarrow y) \rightarrow a^n = 0 \in \{0\}$ ,  $y \rightarrow a^n = 0 \in \{0\}$  and since  $\{0\}$  is an  $n$ -fold implicative ideal of  $L$ , we have  $x \rightarrow a^n \in \{0\}$ . Hence,  $x \rightarrow a^n = 0$  and so  $x \in L(a)$ . Therefore,  $L(a)$  is an ideal of  $L$ .

(iii)  $\Leftrightarrow$  (ii) Let  $(x \rightarrow y) \rightarrow z_{\mathcal{O}}^n \in \{0\}$  and  $y \rightarrow z_{\mathcal{O}}^n \in \{0\}$ , for  $x, y, z \in L$ . Then  $(x \rightarrow y) \in L(z_{\mathcal{O}}^n)$  and  $y \in L(z_{\mathcal{O}}^n)$  and since  $L(z_{\mathcal{O}}^n)$  is an ideal of  $L$ , we get  $x \in L(z_{\mathcal{O}}^n)$ , and so  $x \rightarrow z_{\mathcal{O}}^n = 0$ . Hence,  $\{0\}$  is an  $n$ -fold implicative ideal of  $L$ . ■

**Proposition 26.** *Let  $L$  be Boolean algebra or Gödel algebra. Then any ideal of  $L$  is an  $n$ -fold implicative ideal of  $L$  for any natural number  $n$ .*

**Proof.** It follows from Theorems 12 and 22. ■

**Theorem 27.** *Let  $I$  be a proper ideal of a  $L$ . Then the following conditions are equivalent:*

- (i)  *$I$  is a maximal and  $n$ -fold implicative ideal of  $L$ ,*
- (ii)  *$x, y \notin I$  imply  $x \rightarrow y_{\mathcal{O}}^n \in I$  and  $y \rightarrow x_{\mathcal{O}}^n \in I$ , for all  $x, y \in L$ ,*
- (iii) *if  $x \notin I$ , then there exists natural number  $m$  such that  $((x_{\mathcal{O}}^n)^-)^m \in I$ ,*
- (iv)  *$(x^-)_{\mathcal{O}}^n \in I$  or  $((x^-)_{\mathcal{O}}^n)^- \in I$ , for all  $x \in L$ ,*
- (v)  *$I$  is a prime and  $n$ -fold implicative ideal of  $L$ ,*
- (vi)  *$I$  is a prime and  $n$ -fold Boolean ideal of  $L$ .*

**Proof.** (i)  $\Leftrightarrow$  (ii) Let  $I$  be a maximal and  $n$ -fold implicative ideal of  $L$  and  $x, y \notin I$ . Then by Theorem 17,  $I_{y_{\mathcal{O}}^n} = \{z \in L \mid z \rightarrow y_{\mathcal{O}}^n \in I\}$  is the least ideal of  $L$  containing  $I$  and  $y$  and since  $I$  is maximal ideal of  $L$  and  $y \notin I$ , we have  $I_{y_{\mathcal{O}}^n} = L$ , and so  $x \in I_{y_{\mathcal{O}}^n}$ . Therefore,  $x \rightarrow y_{\mathcal{O}}^n \in I$ . By similar way  $y \rightarrow x_{\mathcal{O}}^n \in I$ .

(ii)  $\Leftrightarrow$  (iii) Suppose that  $x \notin I$ . Since  $I$  is a proper ideal, we have  $1 \notin I$  and so by hypothesis  $1 \rightarrow x_{\mathcal{O}}^n = (x_{\mathcal{O}}^n)^- \in I$ . Hence, for some natural number  $m$ ,  $((x_{\mathcal{O}}^n)^-)^m \in I$ .

(iii)  $\Leftrightarrow$  (iv) For any  $x \in L$ , if  $x^- \in I$ , then  $(x^-)_{\odot}^n \in I$ . Assume that  $x^- \notin I$ , then there exists natural number  $m$  such that  $((x^-)_{\odot}^n)^m \in I$  and since by Lemma 2(vi),  $((x^-)_{\odot}^n)^- \leq (((x^-)_{\odot}^n)^-)^m$  and  $I$  is an ideal of  $L$ , we get that  $((x^-)_{\odot}^n)^- \in I$ . Thus, (iv) is valid.

(iv)  $\Leftrightarrow$  (v) Let  $(x^-)_{\odot}^n \in I$  or  $((x^-)_{\odot}^n)^- \in I$ , for all  $x \in L$ . Then by Lemma 20(ii),  $(x^n)^- \in I$  or  $(x^n)^{- -} \in I$ , for all  $x \in L$ , and since  $I$  is an ideal of  $L$ , we obtain  $(x^n)^- \in I$  or  $x^n \text{implicationalgebran} I$ , for all  $x \in L$ . Now, by Theorem 7,  $I$  is a prime and  $n$ -fold Boolean ideal of  $L$  and so by Theorem 24,  $I$  is a prime and  $n$ -fold implicative ideal of  $L$ .

(v)  $\Leftrightarrow$  (vi) It follows from Theorem 24.

(vi)  $\Leftrightarrow$  (i) Let  $I$  be a prime and  $n$ -fold Boolean ideal of  $L$ . Then by Theorem 7,  $I$  is a maximal and  $n$ -fold Boolean ideal of  $L$ . Hence, by Theorem 24,  $I$  is a maximal and  $n$ -fold implicative ideal of  $L$ . ■

#### 4. N-FOLD OBSTINATE IDEALS IN BL-ALGEBRAS

In this section we introduce a new class of ideals in  $BL$ -algebras that called  $n$ -fold obstinate ideals and we give some results.

**Definition.** Let  $I$  be an ideal of  $L$ .  $I$  is called an  $n$ -fold obstinate ideal if it satisfies:

$$x, y \notin I \text{ imply } x \rightarrow y_{\odot}^n \in I \text{ and } y \rightarrow x_{\odot}^n \in I, \text{ for all } x, y \in L$$

**Example 28.** [6] Let  $L = \{0, a, b, 1\}$ , where  $0 < a < b < 1$ . Let  $x \wedge y = \min\{x, y\}$ ,  $x \vee y = \max\{x, y\}$  and operations  $\odot$  and  $\rightarrow$  are defined as the following tables:

Table 3

$\odot$	0	a	b	1
0	0	0	0	0
a	0	0	a	a
b	0	a	b	b
1	0	a	b	1

Table 4

$\rightarrow$	0	a	b	1
0	1	1	1	1
a	a	1	1	1
b	0	a	1	1
1	0	a	b	1

Then  $(L, \vee, \wedge, \odot, \rightarrow, 0, 1)$  is a  $BL$ -algebra. Now, let  $I = \{0\}$ . Then  $I$  is a 2-fold obstinate ideal of  $L$ , but it is not a 1-fold obstinate ideal. Indeed,  $a, b \notin \{0\}$  and  $b \rightarrow a = b \odot a^- = b \odot a = a \notin \{0\}$ .

**Theorem 29.** Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold obstinate ideal of  $L$  if and only if  $I$  is an  $n$ -fold integral ideal of  $L$ .

**Proof.** It follows from Theorems 7 and 27. ■

**Theorem 30.** *Let  $I$  be a proper ideal and  $F$  be a proper filter of  $L$ . Then*

- (i)  *$I$  is an  $n$ -fold obstinate ideal if and only if  $N(I)$  is an  $n$ -fold obstinate filter of  $L$ .*
- (ii)  *$F$  is an  $n$ -fold integral filter if and only if  $N(F)$  is an  $n$ -fold obstinate ideal of  $L$ .*

**Proof.** It follows from Theorems 8, 9 and 29. ■

The following theorem describes the relationship between  $n$ -fold obstinate ideals and  $n$ -fold integral  $BL$ -algebras.

**Theorem 31.** *In any  $BL$ -algebra  $L$ , the following conditions are equivalent:*

- (i)  *$\{0\}$  is an  $n$ -fold obstinate ideal of  $L$ ,*
- (ii) *any ideal of  $L$  is an  $n$ -fold obstinate ideal,*
- (iii)  *$L$  is an  $n$ -fold integral  $BL$ -algebra.*

**Proof.** It follows from Theorems 10 and 29. ■

**Theorem 32.** *Let  $I$  be an ideal of  $L$ . Then  $I$  is an  $n$ -fold obstinate ideal of  $L$  if and only if  $\frac{L}{I}$  is an  $n$ -fold obstinate  $BL$ -algebra.*

**Proof.** It follows from Theorems 11 and 29. ■

**Example 33.** Let  $L$  be  $BL$ -algebra given in Example 28 and  $I = \{0\}$ , which is a 2-fold obstinate ideal of  $L$ . We have  $\frac{L}{I} = \{[0], [a], [1]\}$ , where  $[0] = \{0\}$ ,  $[a] = \{a\}$  and  $[1] = \{b, 1\}$ . Note that  $\frac{L}{I}$  is an  $MV$ -algebra and  $[a]^2 = [a^2] = [0]$ . Hence,  $\frac{L}{I}$  is a 2-fold obstinate  $BL$ -algebra.

## 5. CONCLUSION

The results of this paper are devoted to study two new classes of ideals that is called  $n$ -fold implicative ideals and  $n$ -fold obstinate ideals. We presented a characterization and several important properties of  $n$ -fold implicative ideals and  $n$ -fold obstinate ideals. In particular, we proved that an ideal is  $n$ -fold implicative ideal if and only if is an  $n$ -fold Boolean ideal. Also, we proved that a  $BL$ -algebra is an  $n$ -fold integral  $BL$ -algebra if and only if trivial ideal  $\{0\}$  is an  $n$ -fold obstinate ideal. Moreover, we studied the relation between  $n$ -fold obstinate ideals and  $n$ -fold (integral) obstinate filters in  $BL$ -algebras by using the set of complement elements.



## REFERENCES

- [1] C.C. Chang, *Algebraic analysis of many valued logics*, Trans. Amer. Math. Soc. **88** (1958) 467–490.  
doi:10.1090/S0002-9947-1958-0094302-9
- [2] A. Di Nola, G. Georgescu and A. Iorgulescu, *Pseudo BL-algebras Part I*, Mult. Val. Logic, **8** (2002) 673–714.
- [3] A. Di Nola and L. Leustean, *Compact representations of BL-algebras*, Department of Computer Science, University Aarhus. BRICS Report Series, (2002).
- [4] M. Haveski and E. Eslami, *n-Fold filters in BL-algebras*, Math. Log. Quart. **54** (2008) 178–186.
- [5] S. Motamed and A.B. Saeid, *n-Fold obstinate filters in BL-algebras*, Neural. Comput. Applic. **20** (2011) 461–472.
- [6] C. Lele and J.B. Nganou, *MV-algebras derived from ideals in BL-algebras*, Fuzzy Sets and Systems **218** (2013) 103–113.
- [7] P. Hájek, *Metamathematics of Fuzzy Logic*, Trends in Logic **4** (Kluwer Academic Publishers, 1998), ISBN:9781402003707.
- [8] A. Paad, *Integral ideals and maximal ideals in BL-algebras*, An.Univ. Craiova Ser. Mat. Inform. **43** (2016) 231–242.
- [9] A. Paad, *n-Fold integral ideals and n-fold Boolean ideals in BL-algebras*, Afr. Mat. **28** (2017) 971–984.
- [10] A. Paad and R.A. Borzooei, *Generalization of integral filters in BL-algebras and n-fold integral BL-algebras*, Afr. Mat. **26** (2015) 1299–1311.
- [11] Y. Yang and X. Xin, *On characterization of BL-algebras via implicative ideals*, Italian J. Pure and Appl. Math. **37** (2017) 493–506.
- [12] E. Turunen, *Boolean deductive systems of BL-algebras*, Arch. Math. Logic. **40** (2001) 467–473.

Received 5 July 2018

Revised 3 September 2018

Accepted 8 September 2018

