

f -FIXED POINTS OF ISOTONE f -DERIVATIONS ON A LATTICE

LEMNAOUAR ZEDAM, MOURAD YETTOU

AND

ABDELAZIZ AMROUNE

Laboratory of Pure and Applied Mathematics
Department of Mathematics
Mohamed Boudiaf University
Msila 28000, Algeria

e-mail: l.zedam@gmail.com
yettou28mourad28@gmail.com
aamrounedz@yahoo.fr

Abstract

In a recent paper, Çeven and Öztürk have generalized the notion of derivation on a lattice to f -derivation, where f is a given function of that lattice into itself. Under some conditions, they have characterized the distributive and modular lattices in terms of their isotone f -derivations. In this paper, we investigate the most important properties of isotone f -derivations on a lattice, paying particular attention to the lattice (resp. ideal) structures of isotone f -derivations and the sets of their f -fixed points. As applications, we provide characterizations of distributive lattices and principal ideals of a lattice in terms of principal f -derivations.

Keywords: lattice, isotone f -derivation, principal f -derivation, f -fixed points set.

2010 Mathematics Subject Classification: 03G10, 06B05, 06B10, 06B99.

1. INTRODUCTION

The notion of derivation appeared on the ring structures and it has many applications (see, e.g. [1]). Szász [15, 16] has extended the notion of derivation on a lattice structure L as a function d of L into itself satisfying the following two conditions:

$$d(x \wedge y) = (d(x) \wedge y) \vee (x \wedge d(y)) \text{ and } d(x \vee y) = d(x) \vee d(y),$$

for any $x, y \in L$. Ferrari [5] has investigated some properties of this notion and provided some interesting examples in particular classes of lattices. Xin *et al.* [19] have ameliorated the notion of derivation on a lattice by considering only the first condition, and they have showed that the second condition is obviously holds for the isotone derivations on a distributive lattice. In the same paper, they characterized also the distributive and modular lattices in terms of their isotone derivations. Later on, Xin [20] has focused his attention to the structure of the fixed sets of derivations on a lattice and showed some relationships between lattice ideals and these fixed sets.

In the same direction, Çeven and Öztürk [3] have generalized the notion of derivation on a lattice L to f -derivation on L by using a function f of L into itself. For a given function f of L into itself, an f -derivation on a lattice L is a function d of L into itself satisfying:

$$d(x \wedge y) = (d(x) \wedge f(y)) \vee (f(x) \wedge d(y)), \text{ for any } x, y \in L.$$

In this context, they also characterized the distributive and modular lattices by isotone f -derivations.

This notion of f -derivation on a lattice is witnessing increased attention. It studies, among others, in semi-lattices [21], in bounded hyperlattices [17], in quantales and residuated lattices [6, 18], in distributive lattices [12], and in several kinds of algebras [7, 9, 10]. Furthermore, it is used in the definition of congruences and ideals in a distributive lattice [11].

The aim of the present paper is to investigate the most important properties of isotone and principal f -derivations on a lattice. We pay particular attention to the lattice structure of isotone f -derivations on a lattice, and to the ideal structure of their f -fixed points sets. More specifically, we show some cases that the set of principal f -derivations on a lattice has a lattice structure, and we provide a representation of any lattice in terms of its principal f -derivations. We give a relationship between a distributive lattice and its lattice of isotone f -derivations, and we show a characterization theorem of a distributive lattice in terms of its principal f -derivations. Furthermore, we investigate the structure of the set of f -fixed points of an isotone f -derivation on a lattice, and we show some cases that this set is an ideal (resp. a principal ideal). As applications, we provide a representation of any lattice in terms of its principal f -derivations, and we show characterization theorems of distributive lattices (resp. principal ideals of a lattice) in terms of principal f -derivations.

The remainder of the paper is structured as follows. In Section 2, we recall the necessary basic concepts and properties of lattices and f -derivations on lattices. In Section 3, we provide a representation (resp. a characterization theorem) of any lattice (resp. distributive lattice) in terms of its principal f -derivations.

In Section 4, we study the structure of the f -fixed points set of an isotone f -derivation on a lattice, and we provide a characterization theorem of principal ideals of a lattice in terms of its principal f -derivations. In Section 5, we show that the set of f -fixed points sets of isotone f -derivations on a distributive lattice has also a structure of a distributive lattice, and we provide a representation of any distributive lattice based on the f -fixed points sets of its principal f -derivations. Finally, we present some concluding remarks in Section 6.

2. BASIC CONCEPTS

In this section, we recall the necessary basic concepts and properties of lattices and f -derivations on lattices.

2.1. Lattices

In this subsection, we recall some definitions and properties of lattices that will be needed throughout this paper. Further information can be found in [2, 4, 8, 13, 14].

An *order relation* \leq on a set X is a binary relation on X that is *reflexive*, *antisymmetric* and *transitive*. A set X equipped with an order relation \leq is called a *partially ordered set* (*poset*, for short), denoted (X, \leq) . Let (X, \leq) be a poset and A be a subset of X . An element $x_0 \in X$ is called a *lower bound* of A if $x_0 \leq x$, for any $x \in A$. x_0 is called the *greatest lower bound* (or the *infimum*) of A if x_0 is a lower bound and $m \leq x_0$, for any lower bound m of A . *Upper bound* and *least upper bound* (or *supremum*) are defined dually. Let (X, \leq_X) and (Y, \leq_Y) be two posets. A mapping $\varphi : X \rightarrow Y$ is called an *order isomorphism* if it is surjective and satisfies the following condition:

$$x \leq_X y \text{ if and only if } \varphi(x) \leq_Y \varphi(y), \text{ for any } x, y \in X.$$

If $X = Y$, an order isomorphism $\varphi : X \rightarrow X$ is called an order automorphism.

A poset (L, \leq) is called a \wedge -semi-lattice if any two elements x and y have a greatest lower bound, denoted by $x \wedge y$ and called the *meet* (infimum) of x and y . Analogously, it is called a \vee -semi-lattice if any two elements x and y have a smallest upper bound, denoted by $x \vee y$ and called the *join* (supremum) of x and y . A poset (L, \leq) is called a *lattice* if it is both a \wedge -semi-lattice and a \vee -semi-lattice. A lattice can also be defined as an algebraic structure: a set L equipped with two binary operations \wedge and \vee that are idempotent, commutative and associative, and satisfy the absorption laws (i.e., $x \wedge (x \vee y) = x$ and $x \vee (x \wedge y) = x$, for any $x, y \in L$). The order relation and the meet and join operations are then related as follows: $x \leq y$ if and only if $x \wedge y = x$; $x \leq y$ if and only if $x \vee y = y$. Usually, the notation (L, \leq, \wedge, \vee) is used for a lattice. A poset (L, \leq) is called *bounded*

if it has a least and a greatest element, respectively denoted by 0 and 1. Often, the notation $(L, \leq, \wedge, \vee, 0, 1)$ is used to describe a bounded lattice. A non-empty subset M of a lattice (L, \leq, \wedge, \vee) is called a sublattice of L if, for any $x, y \in M$, it holds that $x \wedge y \in M$ and $x \vee y \in M$. A poset (L, \leq) is called a complete lattice if every subset A of L has both a greatest lower bound, denoted by $\bigwedge A$ and called the infimum of A , and a least upper bound, denoted by $\bigvee A$ and called the supremum of A , in (L, \leq) . A lattice (L, \leq, \wedge, \vee) is called distributive if one of the following two equivalent conditions holds:

- (a) $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$, for any $x, y, z \in L$;
- (a^δ) $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$, for any $x, y, z \in L$.

Let (L, \leq, \wedge, \vee) and $(M, \preceq, \wedge, \vee)$ be two lattices. A mapping $\varphi : L \rightarrow M$ is called a \wedge -homomorphism (resp. \vee -homomorphism), if it satisfies $\varphi(x \wedge y) = \varphi(x) \wedge \varphi(y)$ (resp. $\varphi(x \vee y) = \varphi(x) \vee \varphi(y)$), for any $x, y \in L$. A \wedge -monomorphism is an injective \wedge -homomorphism. Also, a \wedge -epimorphism is a surjective \wedge -homomorphism. \vee -monomorphism and \vee -epimorphism are defined dually. A lattice homomorphism is both a \wedge -homomorphism and a \vee -homomorphism, a lattice isomorphism is a bijective lattice homomorphism. If $L = M$, a lattice isomorphism $\varphi : L \rightarrow L$ is called a lattice automorphism.

Proposition 2.1 [4]. *Let L and M be two lattices, and $\varphi : L \rightarrow M$ be a mapping. The following statements are equivalent:*

- (i) φ is an order isomorphism;
- (ii) φ is a lattice isomorphism.

2.2. f -derivations on a lattice

In this subsection, we recall the definition and some properties of f -derivation on a lattice. Further information can be found in [3, 19, 21].

Definition 2.1 [19]. Let (L, \leq, \wedge, \vee) be a lattice. A function $d : L \rightarrow L$ is called a *derivation* on L if it satisfies the following condition:

$$d(x \wedge y) = (d(x) \wedge y) \vee (x \wedge d(y)), \text{ for any } x, y \in L.$$

Definition 2.2 [3]. Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a function. A function $d : L \rightarrow L$ is called an f -derivation on L if it satisfies the following condition:

$$d(x \wedge y) = (d(x) \wedge f(y)) \vee (f(x) \wedge d(y)), \text{ for any } x, y \in L.$$

Throughout this paper, we shortly write dx instead of $d(x)$ and fx instead of $f(x)$.

Definition 2.3 [3]. Let (L, \leq, \wedge, \vee) be a lattice and d be an f -derivation on L . d is called *isotone* if it satisfies the following condition:

$$x \leq y \text{ implies } dx \leq dy, \text{ for any } x, y \in L.$$

The following proposition gives some proprieties of f -derivations on a lattice.

Proposition 2.2 [3]. Let (L, \leq, \wedge, \vee) be a lattice and d be an f -derivation on L . Then the following holds.

- (i) $dx \leq fx$, for any $x \in L$;
- (ii) If (L, \leq, \wedge, \vee) is distributive, f is a \vee -homomorphism and d is isotone, then $d(x \vee y) = dx \vee dy$.

Proposition 2.3 [3]. Let (L, \leq, \wedge, \vee) be a lattice, $\alpha \in L$ and $f : L \rightarrow L$ be a function satisfies $f(x \wedge y) = fx \wedge fy$, for any $x, y \in L$. Then the function $d_{(\alpha, f)} : L \rightarrow L$ defined by $d_{(\alpha, f)}(x) = \alpha \wedge fx$, for any $x \in L$, is an f -derivation on L . In addition, if f is an increasing function, then $d_{(\alpha, f)}$ is an isotone f -derivation.

The following sets are the key notions of this paper.

Notation 2.1. Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a function. We denote by:

- (i) $\mathfrak{I}_f(L)$ the set of isotone f -derivations on L ;
- (ii) $\mathcal{P}_f(L) := \{d_{(\alpha, f)} \mid \alpha \in L\}$.

The following result shows that the set of isotone f -derivations on a distributive lattice has also a structure of a distributive lattice.

Theorem 1 [21]. Let (L, \leq, \wedge, \vee) be a distributive lattice and d_1, d_2 be two isotone f -derivations on L . Define $(d_1 \sqcap d_2)(x) = d_1x \wedge d_2x$ and $(d_1 \sqcup d_2)(x) = d_1x \vee d_2x$, for any $x \in L$. Then the structure $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$ is a distributive lattice, where the order relation \preceq is defined as:

$$d_1 \preceq d_2 \text{ if and only if } d_1 \sqcup d_2 = d_2, \text{ for any } d_1, d_2 \in \mathfrak{I}_f(L).$$

3. PRINCIPAL f -DERIVATIONS ON A LATTICE

In this section, we show a necessary and sufficient condition that the functions $d_{(\alpha, f)}$ on a lattice L are f -derivations (called principal f -derivations). Also, we show that their set has a lattice structure. Furthermore, we provide a representation (resp. a characterization) theorem of any lattice (resp. distributive lattice) in terms of its principal f -derivations.

3.1. Poset structure for the set of principal f -derivations on a lattice

The following result shows a necessary and sufficient condition that the functions $d_{(\alpha,f)}$ on a lattice L being f -derivations on L .

Theorem 2. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a function. Then it holds that f is a \wedge -homomorphism if and only if $d_{(\alpha,f)}$ is an f -derivation on L , for any $\alpha \in L$.*

Proof. The direct implication follows from Proposition 2.3. For the converse implication, we assume that $d_{(\alpha,f)}$ is an f -derivation on L , for any $\alpha \in L$. It follows that

$$\begin{aligned} d_{(\alpha,f)}(x \wedge y) &= \alpha \wedge f(x \wedge y) \\ &= (d_{(\alpha,f)}(x) \wedge fy) \vee (fx \wedge d_{(\alpha,f)}(y)) \\ &= (\alpha \wedge fx \wedge fy) \vee (fx \wedge \alpha \wedge fy) \\ &= \alpha \wedge fx \wedge fy, \text{ for any } \alpha, x, y \in L. \end{aligned}$$

Hence, $\alpha \wedge f(x \wedge y) = \alpha \wedge fx \wedge fy$, for any $\alpha, x, y \in L$. On the one hand, setting $\alpha = f(x \wedge y)$. Then it follows that $f(x \wedge y) = f(x \wedge y) \wedge (fx \wedge fy)$, for any $x, y \in L$. Hence, $f(x \wedge y) \leq fx \wedge fy$, for any $x, y \in L$. On the other hand, setting $\alpha = fx \wedge fy$. Then it follows that $(fx \wedge fy) \wedge f(x \wedge y) = (fx \wedge fy) \wedge (fx \wedge fy) = fx \wedge fy$, for any $x, y \in L$. Hence, $fx \wedge fy \leq f(x \wedge y)$, for any $x, y \in L$. Thus, $f(x \wedge y) = fx \wedge fy$, for any $x, y \in L$. Therefore, f is a \wedge -homomorphism. ■

The following corollary expresses the relationship between $\mathcal{P}_f(L)$ and $\mathfrak{I}_f(L)$.

Corollary 3.1. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a function. Then it holds that f is a \wedge -homomorphism if and only if $\mathcal{P}_f(L)$ is a subset of $\mathfrak{I}_f(L)$.*

Proof. The proof is directly from Theorem 2. ■

In what follows, for a given lattice L , the f -derivations $d_{(\alpha,f)}$ will be called principal f -derivations on L , and $\mathcal{P}_f(L)$ denotes their set. On $\mathcal{P}_f(L)$, we define a binary relation \leq' as follows:

$$d_{(\alpha,f)} \leq' d_{(\beta,f)} \text{ if and only if } d_{(\alpha,f)}(x) \leq d_{(\beta,f)}(x), \text{ for any } x \in L.$$

One easily verifies that \leq' is an order relation on $\mathcal{P}_f(L)$.

Remark 3.1. If $(L, \leq, \wedge, \vee, 0, 1)$ is a bounded lattice, then the poset $(\mathcal{P}_f(L), \leq')$ is also bounded, where $0_{\mathcal{P}_f(L)} = d_{(0,f)}$ and $1_{\mathcal{P}_f(L)} = d_{(1,f)}$ such that $d_{(0,f)}(x) = 0$ and $d_{(1,f)}(x) = fx$, for any $x \in L$.

3.2. Lattice structure for the poset of principal f -derivations on a lattice

In this subsection, we show some cases in which the poset $(\mathcal{P}_f(L), \leq')$ of principal f -derivations on a lattice L has a lattice structure. Also, we provide a representation of a lattice in terms of its principal f -derivations. First, we show that $(\mathcal{P}_f(L), \leq')$ is a \wedge -semi-lattice.

Proposition 3.1. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then the poset $(\mathcal{P}_f(L), \leq')$ is a \wedge -semi-lattice.*

Proof. Let $d_{(\alpha, f)}, d_{(\beta, f)} \in \mathcal{P}_f(L)$. It is easy to verify that $d_{(\alpha \wedge \beta, f)}$ is the greatest lower bound of $d_{(\alpha, f)}$ and $d_{(\beta, f)}$. Thus, $(\mathcal{P}_f(L), \leq')$ is a \wedge -semi-lattice. ■

The following theorem shows that the set of principal f -derivations on a complete lattice is also a complete lattice.

Theorem 3. *Let (L, \leq, \wedge, \vee) be a complete lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then the poset $(\mathcal{P}_f(L), \leq')$ is a complete lattice.*

Proof. Proposition 3.1 guarantees that $(\mathcal{P}_f(L), \leq')$ is a \wedge -semi-lattice. Let A be a non-empty subset of $\mathcal{P}_f(L)$ and P^u be the set of upper bounds of A . The fact that (L, \leq, \wedge, \vee) is a complete lattice implies that $d_{(\beta, f)} = d_{(\bigwedge \alpha_i, f)}$ with $d_{(\alpha_i, f)} \in P^u$ is the least upper bound of A . Thus, $(\mathcal{P}_f(L), \leq')$ is a complete lattice. ■

The following corollary follows from the above theorem.

Corollary 3.2. *Let (L, \leq, \wedge, \vee) be a finite lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then $(\mathcal{P}_f(L), \leq')$ is a finite lattice.*

The following theorem shows that the set of principal f -derivations on a distributive lattice is also a distributive lattice.

Theorem 4. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then the poset $(\mathcal{P}_f(L), \leq')$ is a distributive lattice.*

Proof. Let $d_{(\alpha, f)}, d_{(\beta, f)} \in L$. The fact that (L, \leq, \wedge, \vee) is distributive implies that $d_{(\alpha \vee \beta, f)}$ is the least upper bound of $d_{(\alpha, f)}$ and $d_{(\beta, f)}$. Thus, $(\mathcal{P}_f(L), \leq')$ is a lattice. Moreover, its distributivity follows from that of (L, \leq, \wedge, \vee) . ■

Next, we provide a representation of a lattice L based on its principal f -derivations. This representation gives also another case where the poset $(\mathcal{P}_f(L), \leq')$ is a lattice. First, we need to show the following lemma.

Lemma 5. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a function. If f is surjective, then for any $\alpha, \beta \in L$, the following equivalence holds:*

$$\alpha \leq \beta \text{ if and only if } d_{(\alpha,f)} \leq' d_{(\beta,f)}.$$

Proof. The direct implication is immediate. For the converse implication, assume that f is surjective. Let $\alpha, \beta \in L$ such that $d_{(\alpha,f)} \leq' d_{(\beta,f)}$. Then $\alpha \wedge fx \leq \beta \wedge fx$, for any $x \in L$. Since f is surjective, it holds that there exists $m \in L$ such that $fm = \alpha$. Hence, $\alpha \wedge fm \leq \beta \wedge fm$. Thus, $\alpha \leq \beta$. ■

Now, we are able to provide a representation of a lattice in terms of its principal f -derivations.

Theorem 6. *Let (L, \leq, \wedge, \vee) be a lattice and f be a \wedge -homomorphism. If f is a \wedge -epimorphism, then the poset $(\mathcal{P}_f(L), \leq')$ is a lattice, where $d_{(\alpha,f)} \wedge' d_{(\beta,f)} = d_{(\alpha \wedge \beta, f)}$ and $d_{(\alpha,f)} \vee' d_{(\beta,f)} = d_{(\alpha \vee \beta, f)}$, for any $d_{(\alpha,f)}, d_{(\beta,f)} \in \mathcal{P}_f(L)$. Moreover, (L, \leq, \wedge, \vee) and $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ are isomorphic.*

Proof. Assume that $f : L \rightarrow L$ is a \wedge -epimorphism. Proposition 3.1 guarantees that $(\mathcal{P}_f(L), \leq')$ is a \wedge -semi-lattice, where $d_{(\alpha,f)} \wedge' d_{(\beta,f)} = d_{(\alpha \wedge \beta, f)}$ is the greatest lower bound of $d_{(\alpha,f)}$ and $d_{(\beta,f)}$, for any $d_{(\alpha,f)}, d_{(\beta,f)} \in \mathcal{P}_f(L)$. Now, we show that $(\mathcal{P}_f(L), \leq')$ is a \vee -semi-lattice. Let $d_{(\alpha,f)}, d_{(\beta,f)} \in \mathcal{P}_f(L)$, since f is surjective, it follows from Lemma 5 that $d_{(\alpha,f)} \vee' d_{(\beta,f)} = d_{(\alpha \vee \beta, f)}$ is the least upper bound of $d_{(\alpha,f)}$ and $d_{(\beta,f)}$. Hence, $(\mathcal{P}_f(L), \leq')$ is a \vee -semi-lattice. Thus, the structure $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ is a lattice.

Next, let $\psi : L \rightarrow \mathcal{P}_f(L)$ be a mapping defined as $\psi(\alpha) = d_{(\alpha,f)}$, for any $\alpha \in L$. It is obvious to verify that ψ is surjective. Furthermore, Lemma 5 guarantees that

$$\alpha \leq \beta \text{ if and only if } \psi(\alpha) \leq' \psi(\beta), \text{ for any } \alpha, \beta \in L.$$

Thus, ψ is an order isomorphism between L and $\mathcal{P}_f(L)$. Proposition 2.1 guarantees that ψ is a lattice isomorphism. Therefore, (L, \leq, \wedge, \vee) and $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ are isomorphic. ■

In the following, we present an illustrative example of Theorem 6.

Example 3.1. Let $L = D(30)$ be the lattice of the positive divisors of 30 given by Hasse diagram in Figure 1, and $f : D(30) \rightarrow D(30)$ be a function defined by the following table:

x	1	2	3	5	6	10	15	30
fx	1	2	5	3	10	6	15	30

The following table presents the elements of $\mathcal{P}_f(D(30))$.

x	1	2	3	5	6	10	15	30
$d_{(1,f)}(x)$	1	1	1	1	1	1	1	1
$d_{(2,f)}(x)$	1	2	1	1	2	2	1	2
$d_{(3,f)}(x)$	1	1	1	3	1	3	3	3
$d_{(5,f)}(x)$	1	1	5	1	5	1	5	5
$d_{(6,f)}(x)$	1	2	1	3	2	6	3	6
$d_{(10,f)}(x)$	1	2	5	1	10	2	5	10
$d_{(15,f)}(x)$	1	1	5	3	5	3	15	15
$d_{(30,f)}(x)$	1	2	5	3	10	6	15	30

One easily verifies that f is a lattice automorphism. Hence, Theorem 6 guarantees that $(\mathcal{P}_f(D(30)), \leq', \wedge', \vee')$ is a lattice and isomorphic to $(D(30), |, gcd, lcm)$.




Figure 1. The Hasse diagrams of the lattices $(D(30), |, gcd, lcm)$ and $(\mathcal{P}_f(D(30)), \leq', \wedge', \vee')$.

Note that the converse of the Theorem 6 does not necessarily hold, as can be seen in the following example.

Example 3.2. Let $L = D(6)$ be the lattice of the positive divisors of 6 given by Hasse diagram in Figure 2, and $f : D(6) \rightarrow D(6)$ be a function defined by the following table:

x	1	2	3	6
fx	1	1	6	6

The following table presents the elements of $\mathcal{P}_f(D(6))$.

$d_{(1,f)}(x)$	1	1	1	1
$d_{(2,f)}(x)$	1	1	2	2
$d_{(3,f)}(x)$	1	1	3	3
$d_{(6,f)}(x)$	1	1	6	6

One easily verifies that f is a \wedge -homomorphism. Moreover, since there not exists $x \in D(6)$ such that $fx = 2$, it holds that f is not surjective. But, as can be seen in Figure 2 that the poset $(\mathcal{P}_f(D(6)), \leq')$ is a lattice and isomorphic to $(D(6), |, gcd, lcm)$.




Figure 2. The Hasse diagrams of the lattices $(D(6), |, gcd, lcm)$ and $(\mathcal{P}_f(D(6)), \leq', \wedge', \vee')$.

3.3. A relationship between a distributive lattice and its lattice of isotone f -derivations

In this subsection, we give a relationship between a distributive lattice L and its lattice of isotone f -derivations $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$. Also, we show a characterization theorem of a distributive lattice in terms of its principal f -derivations. First, we need to recall the following result.

Proposition 3.2 [21]. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then the structure $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ is a sublattice of the distributive lattice $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$.*

In the case of (L, \leq, \wedge, \vee) is a distributive lattice, the following proposition shows that the lattice structures $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ and $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ coincide.

Proposition 3.3. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ coincides with $(\mathcal{P}_f(L), \leq', \wedge', \vee')$.*

Proof. On the one hand, $(d_{(\alpha,f)} \sqcap d_{(\beta,f)})(x) = (\alpha \wedge fx) \wedge (\beta \wedge fx) = (\alpha \wedge \beta) \wedge fx = d_{(\alpha \wedge \beta, f)}(x) = (d_{(\alpha,f)} \wedge' d_{(\beta,f)})(x)$, for any $\alpha, \beta, x \in L$. Then \sqcap coincides with \wedge' on $\mathcal{P}_f(L)$, for any lattice (L, \leq, \wedge, \vee) . On the other hand, we assume that (L, \leq, \wedge, \vee) is a distributive lattice. Let $d_{(\alpha,f)}, d_{(\beta,f)} \in \mathcal{P}_f(L)$, then $(d_{(\alpha,f)} \sqcup d_{(\beta,f)})(x) = (\alpha \wedge fx) \vee (\beta \wedge fx) = (\alpha \vee \beta) \wedge fx = d_{(\alpha \vee \beta, f)}(x) = (d_{(\alpha,f)} \vee' d_{(\beta,f)})(x)$, for any $x \in L$. Thus, \sqcup coincides with \vee' on $\mathcal{P}_f(L)$. Therefore, the lattice structures $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ and $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ coincide. ■

Combining Propositions 3.2 and 3.3 leads to the following corollary.

Corollary 3.3. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a \wedge -homomorphism. Then $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ is a sublattice of $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$.*

The following theorem shows a relationship between a distributive lattice and its lattice of isotone f -derivations.

Theorem 7. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a \wedge -epimorphism. Then (L, \leq, \wedge, \vee) is isomorphic to a sublattice of $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$.*

Proof. Assume that (L, \leq, \wedge, \vee) is a distributive lattice and $f : L \rightarrow L$ is a \wedge -epimorphism. On the one hand, Theorem 6 guarantees that (L, \leq, \wedge, \vee) is isomorphic to $(\mathcal{P}_f(L), \leq', \wedge', \vee')$. On the other hand, Corollary 3.3 shows that $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ is a sublattice of $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$. Consequently, (L, \leq, \wedge, \vee) is isomorphic to a sublattice of $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$. ■

We conclude this subsection by a characterization theorem of a distributive lattice in terms of its principal f -derivations.

Theorem 8. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a \wedge -epimorphism. The following statements are equivalent:*

- (i) (L, \leq, \wedge, \vee) is distributive;
- (ii) $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ is a distributive lattice;
- (iii) \sqcup is a binary operation on $\mathcal{P}_f(L)$;
- (iv) \sqcup coincides with \vee' on $\mathcal{P}_f(L)$.

Proof. (i) \Rightarrow (ii): A straightforward application of Proposition 3.2.

(ii) \Rightarrow (iii): The proof is immediate.

(iii) \Rightarrow (iv): Let $d_{(\alpha,f)}, d_{(\beta,f)} \in \mathcal{P}_f(L)$. The fact that \sqcup is a binary operation on $\mathcal{P}_f(L)$ implies that there exists $d_{(\gamma,f)} \in \mathcal{P}_f(L)$ such that $d_{(\alpha,f)} \sqcup d_{(\beta,f)} = d_{(\gamma,f)}$, this equivalent to

$$(\alpha \wedge fx) \vee (\beta \wedge fx) = \gamma \wedge fx, \text{ for any } x \in L.$$

Since f is surjective, it follows that there exist $a, b, c \in L$ such that $fa = \alpha$, $fb = \beta$ and $fc = \gamma$. Setting $x = a$ (resp. $x = b$), it holds that $\alpha = \gamma \wedge \alpha$ (resp. $\beta = \gamma \wedge \beta$). Moreover, setting $x = c$, we obtain that $\gamma = \alpha \vee \beta$. Hence, $d_{(\alpha,f)} \sqcup d_{(\beta,f)} = d_{(\alpha \vee \beta, f)} = d_{(\alpha,f)} \vee' d_{(\beta,f)}$. Thus, \sqcup coincides with the binary operation \vee' on $\mathcal{P}_f(L)$.

(iv) \Rightarrow (i): Let $\alpha, \beta, \gamma \in L$. Since \sqcup coincides with \vee' on $\mathcal{P}_f(L)$, it holds that $(d_{(\alpha,f)} \sqcup d_{(\beta,f)})(x) = (d_{(\alpha,f)} \vee' d_{(\beta,f)})(x) = d_{(\alpha \vee \beta, f)}(x)$ for any $x \in L$, this equivalent to

$$(\alpha \wedge fx) \vee (\beta \wedge fx) = (\alpha \vee \beta) \wedge fx, \text{ for any } x \in L.$$

The fact that f is surjective implies that there exists $c \in L$ satisfying $fc = \gamma$. Setting $x = c$, we obtain that $(\alpha \wedge \gamma) \vee (\beta \wedge \gamma) = (\alpha \vee \beta) \wedge \gamma$. Thus, (L, \leq, \wedge, \vee) is distributive. \blacksquare

Remark 3.2. From Theorem 8, we conclude that if (L, \leq, \wedge, \vee) is not distributive and $f : L \rightarrow L$ is a \wedge -epimorphism, then $(\mathcal{P}_f(L), \preceq, \sqcap, \sqcup)$ has not a lattice structure. It is only a \sqcap -semi-lattice, indeed, in this case \sqcup can not be a binary operation on $\mathcal{P}_f(L)$.

4. IDEAL STRUCTURE OF f -FIXED POINTS OF AN ISOTONE f -DERIVATION ON A LATTICE

This section is devoted to study the structure of the set of f -fixed points of an isotone f -derivation on a lattice L . More specifically, we present some cases that this set is an ideal of L , and we provide a characterization theorem of principal ideals of L in terms of its principal f -derivations. Furthermore, we show a relationship between prime ideals of L and f -derivations on L . First, we recall the following definitions.

4.1. Definitions

A non-empty subset I of a lattice L is called an *ideal*, if the following two conditions hold:

- (i) if $x \in L$ and $y \in I$ such that $x \leq y$, then $x \in I$;
- (ii) if $x, y \in I$, then $x \vee y \in I$.

An ideal I is called *prime* if $x \wedge y \in I$ implies that $x \in I$ or $y \in I$, for any $x, y \in L$. An ideal is called *principal*, if it is generated by an element $x \in L$. It is the smallest ideal contains x and is given by the set $\downarrow x = \{y \in L \mid y \leq x\}$.

Definition 4.1 [3]. Let (L, \leq, \wedge, \vee) be a lattice and d be an f -derivation on L . The set of f -fixed points of d is given by:

$$\text{Fix}_{(d,f)}(L) = \{x \in L \mid dx = fx\}.$$

Notation 4.1. Let (L, \leq, \wedge, \vee) be a lattice and $f : L \longrightarrow L$ be a function. We denote by:

- (i) $\mathfrak{F}_f(L) := \{\text{Fix}_{(d,f)}(L) \mid d \in \mathfrak{I}_f(L)\};$
- (ii) $\mathcal{F}_f(L) := \{\text{Fix}_{(d_{(\alpha,f)},f)}(L) \mid d_{(\alpha,f)} \in \mathcal{P}_f(L)\}.$

4.2. Ideal structure of the set of f -fixed points of an isotone f -derivation on a lattice

In this subsection, we present some cases that the set of f -fixed points of an isotone (resp. a principal) f -derivations on a lattice L is an ideal of L .

Theorem 9. Let (L, \leq, \wedge, \vee) be a distributive lattice and d be an isotone f -derivation on L . If f is a \vee -homomorphism and $\text{Fix}_{(d,f)}(L)$ is a non-empty set, then $\text{Fix}_{(d,f)}(L)$ is an ideal of L .

Proof. Assume that $f : L \longrightarrow L$ is a \vee -homomorphism. Let d be an isotone f -derivation on L such that $\text{Fix}_{(d,f)}(L)$ is a non-empty set. On the one hand, let $x, y \in L$ such that $x \in \text{Fix}_{(d,f)}(L)$ and $y \leq x$. The fact that d is an f -derivation on L implies from Proposition 2.2 that $dy \leq fy$. Since f is increasing, $y \leq x$ and $x \in \text{Fix}_{(d,f)}(L)$, it follows that $fy \leq fx = dx$. Hence, $dy = d(x \wedge y) = (dx \wedge fy) \vee (fx \wedge dy) = fy \vee dy$, and this implies that $fy \leq dy$. Thus, $dy = fy$, i.e., $y \in \text{Fix}_{(d,f)}(L)$. On the other hand, let $x, y \in \text{Fix}_{(d,f)}(L)$. This implies that $dx = fx$ and $dy = fy$. Since (L, \leq, \wedge, \vee) is distributive, f is a \vee -homomorphism and d is an isotone f -derivation on L , it follows from Proposition 2.2 that $d(x \vee y) = dx \vee dy = fx \vee fy = f(x \vee y)$. Hence, $x \vee y \in \text{Fix}_{(d,f)}(L)$. Finally, we conclude that $\text{Fix}_{(d,f)}(L)$ is an ideal of L . ■

Remark 4.1. In general, the set of f -fixed points of an f -derivation on a lattice L is a non-empty set. Indeed, if (L, \leq, \wedge, \vee) is a lattice has a least element $0 \in L$ and $f0 = 0$, then 0 is an f -fixed point of any f -derivation on L .

Theorem 10. Let (L, \leq, \wedge, \vee) be a lattice and $d_{(\alpha,f)}$ be a principal f -derivation on L such that $\text{Fix}_{(d_{(\alpha,f)},f)}(L)$ is a non-empty set. If f is a lattice homomorphism, then $\text{Fix}_{(d_{(\alpha,f)},f)}(L)$ is an ideal of L .

Proof. Assume that $f : L \longrightarrow L$ is a lattice homomorphism. Let $d_{(\alpha,f)} \in \mathcal{P}_f(L)$ such that $\text{Fix}_{(d_{(\alpha,f)},f)}(L)$ is a non-empty set. On the one hand, let $x, y \in L$ such that $x \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$ and $y \leq x$. The fact that $x \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$ implies

that $d_{(\alpha,f)}(x) = \alpha \wedge fx = fx$. Hence, $fx \leq \alpha$. Now, since f is increasing and $y \leq x$, it holds that $fy \leq fx$. Hence, $fy \leq \alpha$. Thus, $d_{(\alpha,f)}(y) = \alpha \wedge fy = fy$. Therefore, $y \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$. On the other hand, let $x, y \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$. Then $d_{(\alpha,f)}(x) = \alpha \wedge fx = fx$ and $d_{(\alpha,f)}(y) = \alpha \wedge fy = fy$. This implies that $fx \vee fy \leq \alpha$. The fact that f is a \vee -homomorphism and $x, y \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$ imply that $f(x \vee y) = fx \vee fy \leq \alpha$. Hence, $d_{(\alpha,f)}(x \vee y) = \alpha \wedge f(x \vee y) = f(x \vee y)$. Thus, $x \vee y \in \text{Fix}_{(d_{(\alpha,f)},f)}(L)$. Therefore, $\text{Fix}_{(d_{(\alpha,f)},f)}(L)$ is an ideal of L . ■

4.3. Characterization of principal ideals in terms of principal f -derivations on a lattice

In this subsection, we show a characterization theorem of principal ideals of a lattice in terms of its principal f -derivations. First, we show the following key results.

Proposition 4.1. *Let (L, \leq, \wedge, \vee) be a lattice, $\downarrow x$ be a principal ideal of L and $f : L \rightarrow L$ be a \wedge -monomorphism. Then there exists a principal f -derivation $d_{(\alpha,f)} \in \mathcal{P}_f(L)$ such that $\downarrow x = \text{Fix}_{(d_{(\alpha,f)},f)}(L)$, where $\alpha = f(x)$.*

Proof. Let $\downarrow x$ be a principal ideal of L . Since f is a \wedge -monomorphism, it follows that

$$\begin{aligned} \downarrow x &= \{y \in L \mid y \leq x\} \\ &= \{y \in L \mid x \wedge y = y\} \\ &= \{y \in L \mid f(x \wedge y) = f(y)\} \\ &= \{y \in L \mid f(x) \wedge f(y) = f(y)\} \\ &= \{y \in L \mid d_{(f(x),f)}(y) = f(y)\} \\ &= \text{Fix}_{(d_{(f(x),f)},f)}(L). \end{aligned}$$

Thus, there exists $d_{(\alpha,f)} \in \mathcal{P}_f(L)$ such that $\downarrow x = \text{Fix}_{(d_{(\alpha,f)},f)}(L)$, where $\alpha = f(x)$. ■

Proposition 4.2. *Let (L, \leq, \wedge, \vee) be a lattice, $f : L \rightarrow L$ be a lattice automorphism and $d_{(\alpha,f)}$ be a principal f -derivation on L . Then $\text{Fix}_{(d_{(\alpha,f)},f)}(L)$ is a principal ideal of L generated by $f^{-1}(\alpha)$.*

Proof. Let $d_{(\alpha,f)} \in \mathcal{P}_f(L)$. Since f is a lattice automorphism, it follows that

$$\begin{aligned} \text{Fix}_{(d_{(\alpha,f)},f)}(L) &= \{y \in L \mid d_{(\alpha,f)}(y) = fy\} \\ &= \{y \in L \mid \alpha \wedge fy = fy\} \\ &= \{y \in L \mid fy \leq \alpha\} \\ &= \{y \in L \mid y \leq f^{-1}(\alpha)\} \\ &= \downarrow f^{-1}(\alpha). \end{aligned}$$

■

Combining Propositions 4.1 and 4.2 leads to the following characterization theorem of principal ideals of a lattice L in terms of its principal f -derivations.

Theorem 11. *Let (L, \leq, \wedge, \vee) be a lattice and $f : L \rightarrow L$ be a lattice automorphism. Then $\mathcal{F}_f(L) = \{Fix_{(d_{(\alpha,f)},f)}(L) \mid d_{(\alpha,f)} \in \mathcal{P}_f(L)\}$ is exactly the set of principal ideals of L .*

In the following, we present an illustrative example of the above Theorem 11.

Example 4.1. Let $L = D(30)$ be the lattice of the positive divisors of 30 given by Hasse diagram in Figure 1, and f be the $D(30)$ -automorphism given in Example 3.1. Then the following holds:

$$\left\{ \begin{array}{l} Fix_{(d_{(1,f)},f)}(D(30)) = \{1\} = \downarrow 1 = \downarrow f^{-1}(1); \\ Fix_{(d_{(2,f)},f)}(D(30)) = \{1, 2\} = \downarrow 2 = \downarrow f^{-1}(2); \\ Fix_{(d_{(3,f)},f)}(D(30)) = \{1, 5\} = \downarrow 5 = \downarrow f^{-1}(3); \\ Fix_{(d_{(5,f)},f)}(D(30)) = \{1, 3\} = \downarrow 3 = \downarrow f^{-1}(5); \\ Fix_{(d_{(6,f)},f)}(D(30)) = \{1, 2, 5, 10\} = \downarrow 10 = \downarrow f^{-1}(6); \\ Fix_{(d_{(10,f)},f)}(D(30)) = \{1, 2, 3, 6\} = \downarrow 6 = \downarrow f^{-1}(10); \\ Fix_{(d_{(15,f)},f)}(D(30)) = \{1, 3, 5, 15\} = \downarrow 15 = \downarrow f^{-1}(15); \\ Fix_{(d_{(30,f)},f)}(D(30)) = D(30) = \downarrow 30 = \downarrow f^{-1}(30). \end{array} \right.$$

Thus, $\mathcal{F}_f(D(30))$ is the set of principal ideals of $D(30)$.

4.4. A relationship between prime ideals and f -derivations on a lattice

In this subsection, we show a relationship between prime ideals of a lattice L and f -derivations on L . This relationship is a generalization of the result of Theorem 4.13 given by Xin in [20].

Theorem 12. *Let (L, \leq, \wedge, \vee) be a lattice, $f : L \rightarrow L$ be a function and I be a prime ideal of L . The following implications hold:*

- (i) *if f is a \wedge -homomorphism, then there exists an f -derivation d on L such that $I \subseteq Fix_{(d,f)}(L)$;*
- (ii) *if f is a \wedge -monomorphism, then there exists an f -derivation d on L such that $I = Fix_{(d,f)}(L)$.*

Proof. Let $\alpha \in I$ and $d : L \rightarrow L$ be a function defined as

$$dx = \begin{cases} fx, & \text{if } x \in I; \\ f(\alpha \wedge x), & \text{otherwise.} \end{cases}$$

(i) The fact that f is a \wedge -homomorphism and I is a prime ideal of L imply that $d(x \wedge y) = (dx \wedge fy) \vee (fx \wedge dy)$, for any $x, y \in L$. Thus, d is an f -derivation on L . The proof of $I \subseteq \text{Fix}_{(d,f)}(L)$ is straightforward.

(ii) Assume that f is a \wedge -monomorphism. On the one hand, (i) guarantees that d is an f -derivation on L and $I \subseteq \text{Fix}_{(d,f)}(L)$. On the other hand, let $x \in \text{Fix}_{(d,f)}(L)$. Here, we distinguish two possible cases, which are $x \in I$ or $x \notin I$. Now, we prove that the case of $x \notin I$ is an impossible case. Suppose that $x \notin I$, then $dx = f(\alpha \wedge x)$. The fact that $x \in \text{Fix}_{(d,f)}(L)$ implies that $dx = fx$. Hence, $f(\alpha \wedge x) = fx$. Since f is injective, it holds that $\alpha \wedge x = x$, i.e., $x \leq \alpha$. Since $\alpha \in I$ and I is an ideal, it holds that $x \in I$, which contradicts the hypothesis that $x \notin I$. Hence, necessarily $x \in I$. Thus, $\text{Fix}_{(d,f)}(L) \subseteq I$. Finally, we conclude that $I = \text{Fix}_{(d,f)}(L)$. ■

5. STRUCTURE OF THE SET OF f -FIXED POINTS SETS OF ISOTONE f -DERIVATIONS ON A DISTRIBUTIVE LATTICE

In this section, for a given distributive lattice L , we show that the set of f -fixed points sets $\mathfrak{F}_f(L)$ of its isotone f -derivations has also a structure of a distributive lattice. Moreover, we prove that the set of f -fixed points sets $\mathcal{F}_f(L)$ of principal f -derivations on L is a sublattice of $\mathfrak{F}_f(L)$. Finally, we provide a representation of any distributive lattice based on the f -fixed points of its principal f -derivations. First, we prove the following key result.

Proposition 5.1. *Let (L, \leq, \wedge, \vee) be a distributive lattice. For any $\text{Fix}_{(d_1,f)}(L), \text{Fix}_{(d_2,f)}(L) \in \mathfrak{F}_f(L)$, we define:*

$$\text{Fix}_{(d_1,f)}(L) \sqcap' \text{Fix}_{(d_2,f)}(L) = \text{Fix}_{(d_1 \sqcap d_2, f)}(L),$$

and

$$\text{Fix}_{(d_1,f)}(L) \sqcup' \text{Fix}_{(d_2,f)}(L) = \text{Fix}_{(d_1 \sqcup d_2, f)}(L).$$

Then \sqcap' and \sqcup' are idempotent, commutative and associative binary operations on $\mathfrak{F}_f(L)$, and they satisfy the absorption laws.

Proof. Let $\text{Fix}_{(d_1,f)}(L), \text{Fix}_{(d_2,f)}(L) \in \mathfrak{F}_f(L)$. Then $d_1, d_2 \in \mathfrak{I}_f(L)$, i.e., d_1 and d_2 are two isotone f -derivations on L . Since (L, \leq, \wedge, \vee) is distributive, it follows from Theorem 1 that $d_1 \sqcap d_2$ and $d_1 \sqcup d_2$ are also isotone f -derivations on L , i.e., $d_1 \sqcap d_2, d_1 \sqcup d_2 \in \mathfrak{I}_f(L)$. Hence, $\text{Fix}_{(d_1,f)}(L) \sqcap' \text{Fix}_{(d_2,f)}(L), \text{Fix}_{(d_1,f)}(L) \sqcup' \text{Fix}_{(d_2,f)}(L) \in \mathfrak{F}_f(L)$. Thus, \sqcap' and \sqcup' are binary operations on $\mathfrak{F}_f(L)$. Furthermore, the fact that $(\mathfrak{I}_f(L), \preceq, \sqcap, \sqcup)$ is a lattice implies that \sqcap and \sqcup are idempotent, commutative and associative binary operations on $\mathfrak{I}_f(L)$, and they satisfy the absorption laws. These imply that \sqcap' and \sqcup' are also idempotent,

commutative and associative binary operations on $\mathfrak{F}_f(L)$, and they satisfy the absorption laws. ■

Theorem 13. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a function. Then the structure $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$ is a distributive lattice, where the order relation \preceq' is defined as $\text{Fix}_{(d_1, f)}(L) \preceq' \text{Fix}_{(d_2, f)}(L)$ if and only if $\text{Fix}_{(d_1, f)}(L) \sqcup' \text{Fix}_{(d_2, f)}(L) = \text{Fix}_{(d_2, f)}(L)$, for any $\text{Fix}_{(d_1, f)}(L), \text{Fix}_{(d_2, f)}(L) \in \mathfrak{F}_f(L)$.*

Proof. Proposition 5.1 guarantees that $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$ is a lattice. Moreover, from the distributivity of $(\mathfrak{F}_f(L), \preceq, \sqcap, \sqcup)$, we easily verify that $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$ is also distributive. ■

The following Proposition lists some proprieties of the sets of f -fixed points of principal f -derivations on a lattice.

Proposition 5.2. *Let (L, \leq, \wedge, \vee) be a lattice, $f : L \rightarrow L$ be a \wedge -homomorphism and $d_{(\alpha, f)}, d_{(\beta, f)}$ be two principal f -derivations on L . Then it holds that*

- (i) $\text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cap \text{Fix}_{(d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha, f)} \sqcap d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L)$;
- (ii) $\text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cup \text{Fix}_{(d_{(\beta, f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha, f)} \sqcup d_{(\beta, f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L)$;
- (iii) *If (L, \leq, \wedge, \vee) is distributive, then $\text{Fix}_{(d_{(\alpha, f)} \sqcup d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L)$.*

Proof. (i) Let $d_{(\alpha, f)}, d_{(\beta, f)} \in \mathcal{P}_f(L)$. We only prove that $\text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cap \text{Fix}_{(d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L)$, as the fact that $d_{(\alpha, f)} \sqcap d_{(\beta, f)} = d_{(\alpha \wedge \beta, f)}$ implies that $\text{Fix}_{(d_{(\alpha, f)} \sqcap d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L)$. Then

$$\begin{aligned} \text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cap \text{Fix}_{(d_{(\beta, f)}, f)}(L) &= \{x \in L \mid d_{(\alpha, f)}(x) = d_{(\beta, f)}(x) = fx\} \\ &= \{x \in L \mid \alpha \wedge fx = \beta \wedge fx = fx\} \\ &= \{x \in L \mid fx \leq \alpha \wedge \beta\} \\ &= \{x \in L \mid (\alpha \wedge \beta) \wedge fx = fx\} \\ &= \{x \in L \mid d_{(\alpha \wedge \beta, f)}(x) = fx\} \\ &= \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L). \end{aligned}$$

Thus, $\text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cap \text{Fix}_{(d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha, f)} \sqcap d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L)$.

(ii) On the one hand, let $x \in \text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cup \text{Fix}_{(d_{(\beta, f)}, f)}(L)$. Then $x \in \text{Fix}_{(d_{(\alpha, f)}, f)}(L)$ or $x \in \text{Fix}_{(d_{(\beta, f)}, f)}(L)$. Assume that $x \in \text{Fix}_{(d_{(\alpha, f)}, f)}(L)$, it holds that $d_{(\alpha, f)}(x) = \alpha \wedge fx = fx$. Then $(d_{(\alpha, f)} \sqcup d_{(\beta, f)})(x) = (\alpha \wedge fx) \vee (\beta \wedge fx) = fx \vee (\beta \wedge fx) = fx$. Thus, $x \in \text{Fix}_{(d_{(\alpha, f)} \sqcup d_{(\beta, f)}, f)}(L)$. The case of $x \in \text{Fix}_{(d_{(\beta, f)}, f)}(L)$ can be proved similarly. Therefore,

$$\text{Fix}_{(d_{(\alpha, f)}, f)}(L) \cup \text{Fix}_{(d_{(\beta, f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha, f)} \sqcup d_{(\beta, f)}, f)}(L).$$

On the other hand, let $y \in \text{Fix}_{(d_{(\alpha,f)} \sqcup d_{(\beta,f)}, f)}(L)$. Then $(d_{(\alpha,f)} \sqcup d_{(\beta,f)})(y) = (\alpha \wedge fy) \vee (\beta \wedge fy) = fy$. This implies that $d_{(\alpha \vee \beta, f)}(y) = (\alpha \vee \beta) \wedge fy = (\alpha \vee \beta) \wedge [(\alpha \wedge fy) \vee (\beta \wedge fy)] = (\alpha \wedge fy) \vee (\beta \wedge fy) = fy$. Hence, $y \in \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L)$. Thus,

$$\text{Fix}_{(d_{(\alpha,f)} \sqcup d_{(\beta,f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L).$$

Therefore,

$$\text{Fix}_{(d_{(\alpha,f)}, f)}(L) \cup \text{Fix}_{(d_{(\beta,f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha,f)} \sqcup d_{(\beta,f)}, f)}(L) \subseteq \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L).$$

(iii) Since (L, \leq, \wedge, \vee) is distributive, it follows from Proposition 3.3 that $d_{(\alpha,f)} \sqcup d_{(\beta,f)} = d_{(\alpha,f)} \vee' d_{(\beta,f)} = d_{(\alpha \vee \beta, f)}$. Thus,

$$\text{Fix}_{(d_{(\alpha,f)} \sqcup d_{(\beta,f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L). \quad \blacksquare$$

The following result shows that the set of f -fixed points sets $\mathcal{F}_f(L)$ of principal f -derivations on L is a sublattice of $\mathfrak{F}_f(L)$.

Theorem 14. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \longrightarrow L$ be a \wedge -homomorphism. Then the structure $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ is a sublattice of $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$.*

Proof. Since (L, \leq, \wedge, \vee) is a distributive lattice, it holds from Theorem 13 that $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$ is a distributive lattice. The fact that f is a \wedge -homomorphism implies that $\mathcal{F}_f(L)$ is a subset of $\mathfrak{F}_f(L)$. Furthermore, Proposition 5.2 guarantees that

$$\text{Fix}_{(d_{(\alpha,f)}, f)}(L) \sqcap' \text{Fix}_{(d_{(\beta,f)}, f)}(L) = \text{Fix}_{(d_{(\alpha,f)} \sqcap d_{(\beta,f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \wedge \beta, f)}, f)}(L)$$

and

$$\text{Fix}_{(d_{(\alpha,f)}, f)}(L) \sqcup' \text{Fix}_{(d_{(\beta,f)}, f)}(L) = \text{Fix}_{(d_{(\alpha,f)} \sqcup d_{(\beta,f)}, f)}(L) = \text{Fix}_{(d_{(\alpha \vee \beta, f)}, f)}(L),$$

for any $\text{Fix}_{(d_{(\alpha,f)}, f)}(L), \text{Fix}_{(d_{(\beta,f)}, f)}(L) \in \mathcal{F}_f(L)$. Thus, $\mathcal{F}_f(L)$ is closed under \sqcap' and \sqcup' . Therefore, $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ is a sublattice of $(\mathfrak{F}_f(L), \preceq', \sqcap', \sqcup')$. \blacksquare

Combining Theorems 13 and 14 leads to the following corollary.

Corollary 5.1. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \longrightarrow L$ be a \wedge -homomorphism. Then the structure $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ is a distributive lattice.*

Next, we provide a representation of any distributive lattice based on the f -fixed points of its principal f -derivations.

Theorem 15. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \longrightarrow L$ be a lattice automorphism. Then (L, \leq, \wedge, \vee) is isomorphic to $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$.*

Proof. Assume that (L, \leq, \wedge, \vee) is a distributive lattice and $f : L \rightarrow L$ is a lattice automorphism. Corollary 5.1 guarantees that $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ is a distributive lattice. Moreover, let $\psi : L \rightarrow \mathcal{F}_f(L)$ be a mapping defined as:

$$\psi(\alpha) = \text{Fix}_{(d_{(f(\alpha), f)}, f)}(L), \text{ for any } \alpha \in L.$$

Now, we show that ψ is surjective. Let $\text{Fix}_{(d_{(f(\beta), f)}, f)}(L) \in \mathcal{F}_f(L)$. Since f is a lattice automorphism, it holds that there exists $\alpha \in L$ such that $f(\alpha) = \beta$. Then $\psi(\alpha) = \text{Fix}_{(d_{(f(\alpha), f)}, f)}(L) = \text{Fix}_{(d_{(f(\beta), f)}, f)}(L)$. Hence, ψ is surjective.

Next, we prove that

$$\alpha \leq \beta \text{ if and only if } \psi(\alpha) \preceq' \psi(\beta), \text{ for any } \alpha, \beta \in L.$$

Since (L, \leq, \wedge, \vee) is a distributive lattice and f is a lattice automorphism, it follows from Propositions 4.1 and 5.2 that

$$\begin{aligned} \alpha \leq \beta &\iff \alpha \vee \beta = \beta \\ &\iff \downarrow(\alpha \vee \beta) = \downarrow \beta \\ &\iff \text{Fix}_{(d_{(f(\alpha \vee \beta), f)}, f)}(L) = \text{Fix}_{(d_{(f(\beta), f)}, f)}(L) \\ &\iff \text{Fix}_{(d_{(f(\alpha) \vee f(\beta), f)}, f)}(L) = \text{Fix}_{(d_{(f(\beta), f)}, f)}(L) \\ &\iff \text{Fix}_{(d_{(f(\alpha), f)} \sqcup d_{(f(\beta), f)}, f)}(L) = \text{Fix}_{(d_{(f(\beta), f)}, f)}(L) \\ &\iff \text{Fix}_{(d_{(\alpha, f)}, f)} \sqcup' \text{Fix}_{(d_{(\beta, f)}, f)}(L) = \text{Fix}_{(d_{(f(\beta), f)}, f)}(L) \\ &\iff \psi(\alpha) \sqcup' \psi(\beta) = \psi(\beta) \\ &\iff \psi(\alpha) \preceq' \psi(\beta), \end{aligned}$$

for any $\alpha, \beta \in L$. Hence, ψ is an order isomorphism between L and $\mathcal{F}_f(L)$. Proposition 2.1 guarantees that ψ is a lattice isomorphism. Thus, the distributive lattices (L, \leq, \wedge, \vee) and $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ are isomorphic. ■

Combining Theorems 6 and 15 leads to the following corollary.

Corollary 5.2. *Let (L, \leq, \wedge, \vee) be a distributive lattice and $f : L \rightarrow L$ be a lattice automorphism. Then the three distributive lattices (L, \leq, \wedge, \vee) , $(\mathcal{P}_f(L), \leq', \wedge', \vee')$ and $(\mathcal{F}_f(L), \preceq', \sqcap', \sqcup')$ are isomorphic.*

6. CONCLUSION

In this work, we have investigated the most important properties of isotone (resp. principal) f -derivations on a lattice. In particular, we have focused on the lattice (resp. ideal) structures of isotone f -derivations and their f -fixed points sets. These properties and structures lead to some interesting results, such as the characterizations of principal ideals and distributive lattices in terms of principal

f -derivations. Also, a representation of a lattice (resp. a distributive lattice) in terms of its principal f -derivations (resp. the f -fixed points sets of its principal f -derivations).

REFERENCES

- [1] M. Ashraf, S. Ali and C. Haetinge, *On derivations in rings and their applications*, Aligarh Bull. Math. **25** (2006) 79–107.
- [2] G. Birkhoff, *Lattice Theory*, 3rd edition, Amer. Math. Soc. (Providence, RI, 1967).
- [3] Y. Çeven and M. Öztürk, *On f -derivations of lattices*, Bull. Korean Math. Soc. **45** (2008) 701–707.
- [4] B.A. Davey and H.A. Priestley, *Introduction to Lattices and Order*, 2nd edition (Cambridge University Press, 2002).
- [5] L. Ferrari, *On derivations of lattices*, Pure Math. and Appl. **12** (2001) 365–382.
- [6] P. He, X. Xin and J. Zhan, *On derivations and their fixed point sets in residuated lattices*, Fuzzy Sets and Systems **303** (2016) 97–113.
doi:10.1016/j.fss.2016.01.006
- [7] K.H. Kim and B. Davvaz, *On f -derivations of BE-algebras*, J. Chungcheong Math. Soc. **28** (2015) 127–138.
doi:10.14403/jcms.2015.28.1.127
- [8] B. Kolman, R.C. Busby and S.C. Ross, *Discrete Mathematical Structures*, 4th edition (Prentice Hall PTR, 2000).
- [9] S.M. Lee and K.H. Kim, *A note on f -derivations of BCC-algebras*, Pure Math. Sci. **1** (2012) 87–93.
- [10] Ş.A. Özbal and A. Firat, *On f -derivations of incline algebras*, Int. Electronic J. Pure and Appl. Math. **3** (2011) 83–90.
- [11] M.S. Rao, *Congruences and ideals in a distributive lattice with respect to a derivation*, Bulletin of the Section of Logic **42** (2013) 1–10.
- [12] G.C. Rao and K.R. Babu, *The theory of derivations in almost distributive lattice*, Bulletin of the International Mathematical Virtual Institute **7** (2017) 203–216.
- [13] S. Roman, *Lattices and Ordered Sets* (Springer Science+Business Media, New York, 2008).
- [14] B.S. Schröder, *Ordered Sets* (Birkhauser, Boston, 2003).
- [15] G. Szász, *Translationen der verbände*, Acta Fac. Rer. Nat. Univ. Comenianae **5** (1961) 53–57.
- [16] G. Szász, *Derivations of lattices*, Acta Sci. Math. **37** (1975) 149–154.
- [17] J. Wang, Y. Jun, X.L. Xin, T.Y. Li and Y. Zou, *On derivations of bounded hyperlattices*, J. Math. Res. Appl. **36** (2016) 151–161.
doi:10.3770/j.issn:2095-2651.2016.02.003

- [18] Q. Xiao and W. Liu, *On derivations of quantales*, Open Mathematics **14** (2016) 338–346.
doi:10.1515/math-2016-0030
- [19] X.L. Xin, T.Y. Li and J.H. Lu, *On derivations of lattices*, Information Sciences **178** (2008) 307–316.
doi:10.1016/j.ins.2007.08.018
- [20] X.L. Xin, *The fixed set of a derivation in lattices*, Fixed Point Theory and Applications **218** (2012) 1–12.
doi:10.1186/1687-1812-2012-218
- [21] Y.H. Yon and K.H. Kim, *On f -derivations from semilattices to lattices*, Commun. Korean Math. Soc. **29** (2014) 27–36.
doi:10.4134/CKMS.2014.29.1.027

Received 11 November 2018

Revised 10 January 2019

Accepted 13 January 2019