

**SOME UPPER BOUNDS FOR THE DIMENSION
OF THE c -NILPOTENT MULTIPLIER
OF A PAIR OF LIE ALGEBRAS**

HOMAYOON ARABYANI

MOHAMMAD JAVAD SADEGHIFARD

Department of Mathematics
Neyshabur Branch, Islamic Azad University
Neyshabur, Iran

e-mail: arabyani.h@gmail.com
h.arabyani@iau-neyshabur.ac.ir
math.sadeghifard85@gmail.com

AND

SEDIGHEH SHEIKH-MOHSENI

Higher Education Center of Eghlid, Eghlid, Iran

e-mail: sh.mohseni.s@gmail.com

Abstract

The notion of the Schur multiplier of a Lie algebra L was introduced by Batten in 1996. Recently, the first author introduced the concept of the c -nilpotent multiplier of a pair of Lie algebras and gave some exact sequences for the c -nilpotent multiplier of a pair of Lie algebras. The purpose of this paper is to derive some inequalities for dimension of the c -nilpotent multiplier of a pair of Lie algebras.

Keywords: pair of Lie algebras, Schur multiplier, nilpotent Lie algebra.

2010 Mathematics Subject Classification: 17B30, 17B60, 17B99.

1. INTRODUCTION AND PRELIMINARY

The Schur multiplier $\mathcal{M}(G)$ of a group G was introduced by Schur [16] in 1904. Let $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ be a free presentation of a group G . Then the c -nilpotent multiplier of G is defined to be $\mathcal{M}^{(c)}(G) = (R \cap \gamma_{c+1}(F))/[R, {}_c F]$,

where $[R, {}_c F]$ denotes the commutator subgroup $[R, \underbrace{F, \dots, F}_{c\text{-times}}]$ and $c \geq 1$. The case $c = 1$ which has been much studied is the Schur multiplier of G , denoted by $\mathcal{M}(G)$. (See [8], for more information). Recently, authors investigated to develop some results on the group theory case to Lie algebra. In [15], analogues to the c -nilpotent multiplier of groups, for a given Lie algebra L , the c -nilpotent multiplier of L is defined as $\mathcal{M}^{(c)}(L) = (R \cap \gamma_{c+1}(F))/\gamma_{c+1}(R, F)$, where $\gamma_{c+1}(F)$ is the $(c + 1)$ -st term of the lower central series of F , $\gamma_1(R, F) = R$ and $\gamma_{c+1}(R, F) = [\gamma_c(R, F), F]$. The Lie algebra $\mathcal{M}^{(1)}(L) = \mathcal{M}(L)$ is the Schur multiplier of L (see [5, 10, 11] for more information). One may check that $\mathcal{M}^{(c)}(L)$ is independent of the choice of the free presentation of L .

Let (N, L) be a pair of Lie algebras, in which N is an ideal in L . The Schur multiplier of (N, L) to be the abelian Lie algebra $\mathcal{M}(N, L)$ appearing in the following natural exact sequence of Lie algebras

$$\begin{aligned} H_3(L) &\rightarrow H_3(L/N) \rightarrow \mathcal{M}(N, L) \rightarrow \mathcal{M}(L) \\ &\rightarrow \mathcal{M}(L/N) \rightarrow \frac{L}{[N, L]} \rightarrow \frac{L}{L^2} \rightarrow \frac{L}{(L^2 + N)} \rightarrow 0, \end{aligned}$$

where $\mathcal{M}(-)$ and $H_3(-)$ denote the Schur multiplier and the third homology of a Lie algebra, respectively. This is analogous to the definition of the Schur multiplier of a pair of groups given by Ellis [7]. For every free presentation $0 \rightarrow R \rightarrow F \rightarrow L \rightarrow 0$ of L , $\mathcal{M}(N, L)$ is isomorphic to the factor Lie algebra $(R \cap [S, F])/[R, F]$, where S is an ideal in F such that $S/R \cong N$, see [2, 12, 14] for more information. In particular, if $N = L$, then the Schur multiplier of (N, L) is $\mathcal{M}(L)$. Using the above notion, we can define the c -nilpotent multiplier of a pair (N, L) as $\mathcal{M}^{(c)}(N, L) = (R \cap [S, {}_c F])/[R, {}_c F]$. In [1], we introduced some exact sequences and upper bounds for the c -nilpotent multiplier of a pair of Lie algebras (Also see [3, 4, 13] for more information). In this paper, we give some inequalities for the dimension of the c -nilpotent multiplier of a pair of Lie algebras.

All Lie algebras are considered over a fixed field Λ and $[\cdot, \cdot]$ denotes the Lie bracket. We define the subalgebras $Z_c(N, L)$ and $[N, {}_c L]$, for all $c \geq 1$, as follows:

$$Z_c(N, L) = \{n \in N \mid [n, l_1, \dots, l_c] = 0, \forall l_1, \dots, l_c \in L\},$$

$$[N, {}_c L] = \langle [n, l_1, \dots, l_c] \mid n \in N, l_1, \dots, l_c \in L \rangle,$$

where $[n, l_1, \dots, l_c] = [\dots [n, l_1], l_2], \dots, l_c]$. (See [12, 13]).

Let X and Y be Lie algebras. Then $X \wedge Y$ is the non-abelian exterior product of X and Y (see [6]). Moreover, $\wedge^c X$ is the c -th exterior product of X , which is the free Λ -module generated by $x_1 \wedge \dots \wedge x_c$ with $x_i \in X$.

2. SOME INEQUALITIES ON $\dim \mathcal{M}^{(c)}(N, L)$

In this section, we give some inequalities for the dimension of the c -nilpotent multiplier of a pair of Lie algebras. Note that $X \otimes^c Y = X \otimes \underbrace{Y \otimes \cdots \otimes Y}_{c\text{-times}}$ is the abelian tensor product.

The following lemmas are useful for the next results.

Lemma 2.1. *Let L and K be two Lie algebras with central subalgebras N and M , respectively. If $\theta : L \rightarrow K$ is an epimorphism with $\theta(N) = M$, then*

$$\dim \mathcal{M}^{(c)}(M, K) \leq \dim \mathcal{M}^{(c)}(N, L).$$

Proof. We can see, that θ induces an epimorphism from $N \otimes^c L^{ab}$ on to $M \otimes^c K^{ab}$ such that

$$\theta(n \otimes (l_1 + L^2) \otimes \cdots \otimes (l_c + L^2)) = \theta(n) \otimes (\theta(l_1) + K^2) \otimes \cdots \otimes (\theta(l_c) + K^2)$$

for all $n \in N$ and $l_1, l_2, \dots, l_c \in L$.

So, one can easily check that there exists an epimorphism from $\mathcal{M}^{(c)}(N, L)$ on to $\mathcal{M}^{(c)}(M, K)$. Therefore,

$$\dim \mathcal{M}^{(c)}(M, K) \leq \dim \mathcal{M}^{(c)}(N, L). \quad \blacksquare$$

Lemma 2.2. *Let (N, L) be a pair of finite dimensional Lie algebras and M be an ideal in L contained in $Z(N, L)$. Then*

$$\dim(M \cap [N, {}_c L]) \leq \dim \mathcal{M}^{(c)}(N/M, L/M).$$

Proof. If $\sigma : N \wedge^c L \rightarrow L$ is a Lie homomorphism defined by

$$n \wedge (l_1 \wedge l_2 \wedge \cdots \wedge l_c) \mapsto [n, l_1, l_2, \dots, l_c],$$

then $Im(\sigma) = [N, {}_c L]$ and $Ker(\sigma) \cong \mathcal{M}^{(c)}(N, L)$. So, there exists an epimorphism

$$\varphi : N \wedge^c L \rightarrow N/M \wedge^c L/M$$

$$\varphi(n \wedge (l_1 \wedge \cdots \wedge l_c)) = (n + M) \wedge ((l_1 + M) \wedge \cdots \wedge (l_c + M)),$$

for $l_1, l_2, \dots, l_c \in L$ and $n \in N$. So, we obtain an epimorphism $\delta : N/M \wedge^c L/M \rightarrow [N, {}_c L]$ such that $\delta\varphi = \sigma$. Thus,

$$\dim([N, {}_c L]) \leq \dim(N/M \wedge^c L/M).$$

Hence, we have

$$\begin{aligned} & \dim(N/M \wedge^c L/M) + \dim(M \cap [N, {}_c L]) \\ &= \dim \mathcal{M}^{(c)}(N/M, L/M) + \dim([N, {}_c L]), \end{aligned}$$

and so,

$$\dim(M \cap [N, {}_c L]) \leq \dim \mathcal{M}^{(c)}(N/M, L/M). \quad \blacksquare$$

In the following theorem, we generalize a result of Salemkar and Niri (2012) [14].

Theorem 2.3. *Let (M, K) be a pair of nilpotent Lie algebras. If (N, L) is a pair of finite dimensional Lie algebras such that $L/Z_c(N, L) \cong K$ and $N/Z_c(N, L) \cong M$, then*

$$\begin{aligned} \dim([N, {}_c L]) &\leq \dim \mathcal{M}^{(c)}(M/[M, {}_c K], K/[M, {}_c K]) \\ &\quad + \dim([M, {}_c K]) \cdot d(K/Z_c(M, K)), \end{aligned}$$

where $d(X)$ is the minimal number of generators of a Lie algebra X .

Proof. We proceed by induction on the dimension of $[M, {}_c K]$. If $\dim([M, {}_c K]) = 0$, then the result follows from Lemma 2.2. Suppose that $\dim([M, {}_c K]) = n > 0$, and the result holds for any pair (M', K') of finite dimensional nilpotent Lie algebras with $\dim([M', {}_c K']) < n$. Assume that $Z_{c+1}(N, L)$ is the pre-image in the ideal N of $Z(N/Z_c(N, L), L/Z_c(N, L))$, we can see that

$$Z_c(N, L) \not\cong Z_{c+1}(N, L) \cap ([N, {}_c L] + Z_c(N, L))$$

and so, there exists $x \in (Z_{c+1}(N, L) \cap ([N, {}_c L] + Z_c(N, L)) - Z_c(N, L)$. Hence, the following map is a well-defined epimorphism.

$$\begin{aligned} \delta : L/Z_{c+1}(N, L) &\rightarrow [x, {}_c L] \\ \delta(l + Z_{c+1}(N, L)) &= [x, \underbrace{l, \dots, l}_c] \end{aligned}$$

Put $T = [x, {}_c L]$, by [9] we have

$$\dim T \leq d\left(\frac{K}{Z_c(M, K)}\right).$$

Now, put

$$(N^*, L^*) = (N/T, L/T), (M^*, K^*) = (N^*/Z_c(N^*, L^*), L^*/Z_c(N^*, L^*)).$$

As $x + T \in Z_c(N^*, L^*) - (Z_c(N, L)/T)$, it follows that

$$Z_c(N, L)/T \not\cong Z_c(N^*, L^*).$$

Also, the following map is an epimorphism with $\theta(M) = M^*$ and $\text{Ker}\theta \neq 0$.

$$\begin{aligned} \theta : K \cong L/Z_c(N, L) &\rightarrow K^* \\ \theta(l + Z_c(N, L)) &= (l + T) + Z_c(N^*, L^*) \end{aligned}$$

By Lemma 2.1, we obtain

$$\begin{aligned} & \dim \left(\mathcal{M}^{(c)} \left(\frac{M^*}{Z_c(M^*, K^*)}, \frac{K^*}{Z_c(M^*, K^*)} \right) \right) \\ & \leq \dim \left(\mathcal{M}^{(c)} \left(\frac{M}{Z_c(M, K)}, \frac{K}{Z_c(M, K)} \right) \right). \end{aligned}$$

Moreover,

$$d(K^*/Z_c(M^*, K^*)) \leq d(K/Z_c(M, K))$$

and

$$\dim([M^*, {}_c L^*]) < \dim([M, {}_c K]).$$

Hence, by the induction hypothesis

$$\begin{aligned} \dim([N^*, {}_c L^*]) & \leq \dim \mathcal{M}^{(c)}(M^*/Z_c(M^*, K^*), K^*/Z_c(M^*, K^*)) \\ & \quad + \dim([M^*, {}_c K^*]) d(K^*/Z_c(M^*, K^*)) \\ & \leq \dim(\mathcal{M}^{(c)}(M/Z_c(M, K), K/Z_c(M, K))) \\ & \quad + (\dim([M, {}_c K]) - 1) d(K/Z_c(M, K)). \end{aligned}$$

But $\dim([N, {}_c L]) = \dim([N^*, {}_c L]) + \dim T$. Thus,

$$\begin{aligned} \dim([N, {}_c L]) & \leq \dim \mathcal{M}^{(c)}(M/Z_c(M, K), K/Z_c(M, K)) \\ & \quad + (\dim([M, {}_c K]) - 1) d(K/Z_c(M, K)) + \dim T \\ & \leq \dim(\mathcal{M}^{(c)}(M/Z_c(M, K), K/Z_c(M, K))) \\ & \quad + \dim([M, {}_c K]) d(K/Z_c(M, K)), \end{aligned}$$

as required. ■

Using Theorem 2.3, we obtain the following corollary.

Corollary 2.4. *Let (M, K) be a pair of nilpotent Lie algebras. Then for each pair (N, L) of finite dimensional Lie algebras with $L/Z_c(N, L) \cong K$ and $N/Z_c(N, L) \cong M$,*

$$\begin{aligned} \dim([N, {}_c L] \cap Z_c(N, L)) & \leq \dim \mathcal{M}^{(c)}(M/[M, {}_c K], K/[M, {}_c K]) \\ & \quad + \dim([M, {}_c K])(d(K/Z_c(M, K)) - 1). \end{aligned}$$

REFERENCES

- [1] H. Arabyani, *Bounds for the dimension of the c -nilpotent multiplier of a pair of Lie algebras*, Bull. Iranian Math. Soc. **43** (2017) 2411–2418.
doi:10.1142/S1793557119500074

- [2] H. Arabyani, F. Saeedi, M.R.R. Moghaddam and E. Khamseh, *Characterization of nilpotent Lie algebras pair by their Schur multipliers*, Comm. Algebra **42** (2014) 5474–5483. doi:10.1080/00927872.2012.677081
- [3] H. Arabyani and H. Safa, *Some properties of c -covers of a pair of Lie algebras*, Quaest. Math. **42** (2019) 37–45. doi:10.2989/16073606.2018.1437482
- [4] H. Arabyani, *Some results on the c -nilpotent multiplier of a pair of Lie algebras*, Bull. Iranian Math. Soc. **45** (2019) 205–212. doi:10.1007/s41980-018-0126-6
- [5] P. Batten, K. Moneyhun and E. Stitzinger, *On characterizing nilpotent Lie algebras by their multipliers*, Comm. Algebra **24** (1996) 4319–4330. doi:10.1080/00927879608825817
- [6] G. Ellis, *Nonabelian exterior products of Lie algebras and an exact sequence in the homology of Lie algebras*, J. Pure Appl. Algebra **46** (1987) 111–115. doi:10.1016/0022-4049(87)90089-2
- [7] G. Ellis, *The Schur multiplier of a pair of groups*, Appl. Categ. Structures **6** (1998) 355–371. doi:10.1023/A:1008652316165
- [8] G. Karpilovsky, *The Schur Multiplier* (Clarendon Press, Oxford, 1987).
- [9] E.I. Marshal, *The Frattini subalgebra of a Lie algebra*, J. London Math. Soc. **42** (1967) 416–422. doi:10.1112/jlms/s1-42.1.416
- [10] K. Moneyhun, *Isoclinisms in Lie algebras*, Algebras Groups Geom. **11** (1994) 9–22.
- [11] F. Saeedi, H. Arabyani and P. Niroomand, *On dimension of Schur multiplier of nilpotent Lie algebra II*, Asian-Eur. J. Math. **10** (4) (2017) 1750076 (8 pages). doi:10.1142/S1793557117500760
- [12] F. Saeedi, A.R. Salemkar and B. Edalatzadeh, *The commutator subalgebra and Schur multiplier of a pair of nilpotent Lie algebras*, J. Lie Theory **21** (2011) 491–498.
- [13] H. Safa and H. Arabyani, *On c -nilpotent multiplier and c -covers of a pair of Lie algebras*, Comm. Algebra **45** (2017) 4429–4434. doi:10.1080/00927872.2016.1265125
- [14] A.R. Salemkar and S. Alizadeh Niri, *Bounds for the dimension of the Schur multiplier of a pair of nilpotent Lie algebras*, Asian-Eur. J. Math. **5** (2012) 1250059 (9 pages). doi:10.1142/S1793557112500593
- [15] A.R. Salemkar, B. Edalatzadeh and M. Araskhan, *Some inequalities for the dimension of the c -nilpotent multiplier of Lie algebras*, J. Algebra **322** (2009) 1575–1585. doi:10.1016/j.jalgebra.2009.05.036
- [16] I. Schur, *Über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen*, J. Reine Angew. Math. **127** (1904) 20–50. doi:10.1515/crll.1904.127.20

Received 31 August 2019
 Revised 9 December 2019
 Accepted 10 December 2019