Discussiones Mathematicae General Algebra and Applications 41 (2021) 23–31 doi:10.7151/dmgaa.1347

ON THE GENUS OF THE IDEMPOTENT GRAPH OF A FINITE COMMUTATIVE RING

G. Gold Belsi¹, S. Kavitha²

AND

K. Selvakumar¹

Reg. No. 18114012092031

¹Department of Mathematics

Manonmaniam Sundaranar University

Tirunelveli 627 012, Tamil Nadu, India

²Department of Mathematics Gobi Arts and Science College Gobichettipalayam 638 476, Tamilnadu

e-mail: goldbelsi@gmail.com kavithaashmi@gmail.com selva_158@yahoo.co.in

Abstract

Let R be a finite commutative ring with identity. The *idempotent graph* of R is the simple undirected graph I(R) with vertex set, the set of all nontrivial idempotents of R and two distinct vertices x and y are adjacent if and only if xy=0. In this paper, we have determined all isomorphism classes of finite commutative rings with identity whose I(R) has genus one or two. Also we have determined all isomorphism classes of finite commutative rings with identity whose I(R) has crosscap one. Also we study the the book embedding of toroidal idempotent graphs and classify finite commutative rings whose I(R) is a ring graph.

Keywords: idempotent graph, planar, genus, crosscap.

2010 Mathematics Subject Classification: 13A15, 13M05, 05C25, 05C75.

1. Introduction

The study on linking commutative ring theory with graph theory has been started with the concept of the zero-divisor graph of a commutative ring which was first launched by I. Beck [3]. Recall that the *idempotent graph* of a commutative ring R, is a simple undirected graph I(R) with vertex set, the set of all non-trivial idempotents of R and two distinct vertices x and y are adjacent if and only if xy = 0. The concept was first initiated by, Akbari, Habibi, Majidinya and Manaviyat [1]. They obtained some basic results of $I(M_n(R))$, for a division ring R. Influenced by the ideas of the above authors, we try to classify the finite commutative rings with unity whose idempotent graph is planar, ring graph, has genus 1 or 2 and crosscap 1.

2. Preliminaries

In this section, we recollect some definitions and theorems which are required for the subsequent sections.

Let G be a graph with n vertices and q edges. Let C be a cycle of G. We say C is a primitive cycle if it has no chords. Also a graph G has the primitive cycle property (PCP) if any two primitive cycles intersect in at most one edge. The number frank(G) is called the free rank of G and it is the number of primitive cycles of G. Also the number rank(G) = q + n - r, is called the cycle rank of G, where F is the number of connected components of F0. A graph F1 is called a F2 if it satisfies one of the following equivalent conditions:

1. rank(G) = frank(G);

2. F3 satisfies the PCP and F4 does not contain a subdivision of F4 as a subgraph. A F4 split F4 graph is one whose vertex set can be partitioned as the disjoint union of an independent set and a clique (either of which may be empty).

A graph is said to be planar if it can be drawn in the plane so that its edges intersect only at their ends. A planar graph, which has all the vertices in the outer face of the embedding, is called an outerplanar graph. For non-negative integers g and k, let S_g denote the sphere with g handles and N_k denote the sphere with g crosscaps attached to it. It is well-known that every connected compact surface is homeomorphic to S_g or N_k for some non-negative integers g and g. The genus of a graph g, denoted by g(g), is the minimum integer g such that g can be embedded in g. Similarly the crosscap (nonorientable genus) g(g) is the minimum g such that g can be embedded in g and g is toroidal if g(g) = 1. For details on the notion of embedding of graphs in surface, one can refer to White [14] and for graph theory definitions one can refer [4]. Also for ring theory definitions we refer [2].

The following results are useful for further reference in this paper.

Theorem 1 [14, Kuratowski's]. A graph G is planar if and only if it contains no subdivision of K_5 or $K_{3,3}$.

Theorem 2 [8, Theorem 1]. A graph G is outerplanar if and only if it contains no subgraph homeomorphic to $K_{2,3}$ or K_4 .

Lemma 3 [14, Theorem 4.4.7]. $g(K_{m,n}) = \left\lceil \frac{(m-2)(n-2)}{4} \right\rceil$ if $m, n \ge 2$. In particular, $g(K_{4,4}) = g(K_{3,n}) = 1$ if n = 3, 4, 5, 6. Also $g(K_{5,4}) = g(K_{6,4}) = g(K_{m,3}) = 2$ if m = 7, 8, 9, 10.

Lemma 4 [14, Theorem 4.4.7]. Let m,n be positive integers. Then we have the following $\overline{g}(K_{m,n}) = \lceil \frac{1}{2}(m-2)(n-2) \rceil$ if $m,n \geq 2$.

3. The idempotent graph with $g(I(R)) \leq 2$

In this section, we characterize all finite commutative rings R with identity whose I(R) has genus at most two. Using the Euler characteristic formula and a technique of deletion and insertion, we are able to successfully exclude some cases of higher genus.

Remark 5 [1]. Let R be a finite commutative ring. Then

- (i) I(R) is a null graph if and only if R is a field or a local ring.
- (ii) I(R) is a complete graph if and only if I(R) is a complete graph of order 2.

In view of Remark 5, throughout this paper we assume that R is a finite commutative nonlocal ring with nonzero identity. Recall that every Artinian (finite) ring R is decomposed into Artinian local rings, i.e., $R = R_1 \times \cdots \times R_n$, where each (R_i, \mathfrak{m}_i) is a local ring.

We are now in a position to classify all finite nonlocal rings such that the idempotent graph is planar. Note that, a_i 's notates the non-trivial idempotents of R.

Theorem 6. Let R be a finite commutative nonlocal ring. Then I(R) planar (outerplanar) if and only if $n \leq 3$, where n is the number of distinct maximal ideals of R.

Proof. If n=2, then $I(R)\cong K_2$. When n=3, then the proof follows from Figure 1.




Figure 1. $I(R_1 \times R_2 \times R_3)$.

Conversely, assume that I(R) is planar. Since R is finite, $R \cong R_1 \times R_2 \times \cdots \times R_n$, where each R_i is a local ring. Suppose n > 3. Let $\Omega = \{a_1, a_2, \ldots, a_6\}$ where, $a_1 = (1, 0, 0, \ldots, 0), \ a_2 = (0, 1, 0, \ldots, 0), \ a_3 = (1, 1, 0, \ldots, 0), \ a_4 = (0, 0, 1, \ldots, 0), \ a_5 = (0, 0, 0, 1, 0, \ldots, 0), \ a_6 = (0, 0, 1, 1, 0, \ldots, 0)$. Then the subgraph induced by Ω in I(R) contains $K_{3,3}$ as a subgraph and by Theorem 1, we get a contradiction. Hence $n \leq 3$. The proof for outerplanarity can be easily obtained, using a similar argument and Theorem 2.




Figure 2. An embedding of $I(R_1 \times R_2 \times R_3 \times R_4)$ in S_1 .

Now we characterize all finite commutative nonlocal rings R such that the idempotent graph is toroidal.

Theorem 7. Let R be a finite commutative nonlocal ring. Then g(I(R)) = 1 if and only if n = 4, where n is the number of distinct maximal ideals of R.

Conversely, assume that n=4. Let $C=\{a_1,a_2,\ldots,a_{14}\}$ where, $a_1=(1,0,0,0),\ a_2=(0,1,0,0),\ a_3=(1,1,0,0),\ a_4=(1,0,1,0),\ a_5=(1,0,0,1),\ a_6=(0,0,1,0),\ a_7=(0,0,0,1),\ a_8=(0,0,1,1),\ a_9=(0,1,1,0),\ a_{10}=(0,1,0,1),\ a_{11}=(0,1,1,1),\ a_{12}=(1,0,1,1),\ a_{13}=(1,1,0,1),\ a_{14}=(1,1,1,0).$ Then the subgraph induced by C in I(R) contains $K_{3,3}$ as a subgraph. By Theorem 1, $g(I(R))\geq 1$, whereas an embedding of I(R) given in Figure 2 explicitly shows that g(I(R))=1.

Theorem 8. There is no finite commutative nonlocal ring R for which g(I(R)) = 2.

Proof. Since R is finite, $R \cong R_1 \times R_2 \times \cdots \times R_n$, where each R_i is a local ring. By Theorems 6 and 7, $n \geq 5$. Let $E = \{a_1, a_2, \dots, a_{30}\}$, where $a_1 = (1, 0, 0, 0, \dots, 0)$, $a_2 = (0, 1, 0, 0, \dots, 0)$, $a_3 = (0, 0, 1, 0, \dots, 0)$, $a_4 = (0, 0, 0, 1, 0, \dots, 0)$, $a_5 = (0, 0, 0, 0, 1, 0, \dots, 0)$, $a_6 = (1, 1, 0, 0, \dots, 0)$, $a_7 = (1, 0, 1, 0, \dots, 0)$, $a_8 = (1, 0, 0, 1, 0, \dots, 0)$, $a_9 = (1, 0, 0, 0, 1, 0, \dots, 0)$, $a_{10} = (0, 1, 1, 0, \dots, 0)$, $a_{11} = (0, 1, 0, 1, 0, \dots, 0)$, $a_{12} = (0, 1, 0, 0, 1, 0, \dots, 0)$, $a_{13} = (0, 0, 1, 1, 0, \dots, 0)$, $a_{14} = (0, 0, 0, 1, 1, 0, \dots, 0)$, $a_{15} = (0, 0, 1, 0, 1, 0, \dots, 0)$, $a_{16} = (1, 1, 1, 0, \dots, 0)$, $a_{17} = (0, 1, 1, 1, 0, \dots, 0)$, $a_{18} = (0, 0, 1, 1, 1, 0, \dots, 0)$, $a_{19} = (0, 1, 0, 1, 1, 0, \dots, 0)$, $a_{20} = (0, 1, 1, 1, 0, \dots, 0)$, $a_{21} = (1, 0, 1, 1, 0, \dots, 0)$, $a_{22} = (1, 0, 0, 1, 1, 0, \dots, 0)$, $a_{23} = (1, 0, 1, 0, 1, 0, \dots, 0)$, $a_{24} = (1, 1, 0, 0, 1, 0, \dots, 0)$, $a_{25} = (1, 1, 0, 1, 0, \dots, 0)$, $a_{26} = (0, 1, 1, 1, 1, 0, \dots, 0)$, $a_{27} = (1, 0, 1, 1, 1, 0, \dots, 0)$, $a_{28} = (1, 1, 0, 1, 1, 0, \dots, 0)$, $a_{29} = (1, 1, 1, 1, 0, \dots, 0)$, $a_{30} = (1, 1, 1, 1, 0, \dots, 0)$. Then the subgraph induced by E in I(R) contains a subdivision of $K_{3,12}$ as a subgraph and by Lemma 3, $g(I(R)) \geq 3$.




Figure 3. The subgraph induced by B.

We are now in a point to classify all finite nonlocal rings such that the idempotent graph is projective.

Theorem 9. Let R be a finite commutative ring. Then $\overline{g}(I(R)) = 1$ if and only if n = 4, where n is the number of distinct maximal ideals of R.

Proof. Suppose $\overline{g}(I(R)) = 1$. We have, $R \cong R_1 \times R_2 \times \cdots \times R_n$, where each R_i is a local ring. When $n \geq 5$, let $H = \{a_1, a_2, \dots, a_{10}\}$ where $a_1 = (1, 0, 0, \dots, 0), a_2 = (0, 1, 0, \dots, 0), a_3 = (1, 1, 0, \dots, 0), a_4 = (0, 0, 1, 0, \dots, 0), a_5 = (0, 0, 0, 1, 0, \dots, 0), a_6 = (0, 0, 0, 1, 0, \dots, 0), a_7 = (0, 0, 1, 1, 0, \dots, 0), a_8 = (0, 0, 0, 1, 1, 0, \dots, 0), a_9 = (0, 0, 1, 0, 1, 0, \dots, 0), a_{10} = (0, 0, 1, 1, 1, 0, \dots, 0)$. Then the subgraph induced by H in I(R) contains $K_{3,7}$ as a subgraph. By Lemma 4, $\overline{g}(I(R)) \geq 3$, a contradiction. Hence n = 4.

Conversely, assume that n = 4. Let $J = \{a_1, a_2, \dots, a_{14}\}$ where, $a_1 = (1, 0, 0, 0), a_2 = (0, 1, 0, 0), a_3 = (1, 1, 0, 0), a_4 = (1, 0, 1, 0), a_5 = (1, 0, 0, 1),$

 $a_6 = (0,0,1,0), \ a_7 = (0,0,0,1), \ a_8 = (0,0,1,1), \ a_9 = (0,1,1,0), \ a_{10} = (0,1,0,1), \ a_{11} = (0,1,1,1), \ a_{12} = (1,0,1,1), \ a_{13} = (1,1,0,1), \ a_{14} = (1,1,1,0).$ Then the subgraph induced by J in I(R) contains $K_{3,3}$ as a subgraph. By Theorem 4, $\overline{g}(I(R)) \geq 1$, whereas an embedding of I(R) given in Figure 4, explicitly shows that $\overline{g}(I(R)) = 1$.




Figure 4. An embedding of $I(R_1 \times R_2 \times R_3 \times R_4)$ in N_1 .

Theorem 10. There is no finite commutative ring R for which $\overline{g}(I(R)) = 2$.

Proof. Since R is finite, $R \cong R_1 \times R_2 \times \cdots \times R_n$, where each R_i is a local ring. By Theorem 9, $n \geq 5$. Let $K = \{a_1, a_2, \dots, a_{10}\}$ where $a_1 = (1, 0, 0, \dots, 0), a_2 = (0, 1, 0, \dots, 0), a_3 = (1, 1, 0, \dots, 0), a_4 = (0, 0, 1, 0, \dots, 0), a_5 = (0, 0, 0, 1, 0, \dots, 0), a_6 = (0, 0, 0, 0, 1, 0, \dots, 0), a_7 = (0, 0, 1, 1, 0, \dots, 0), a_8 = (0, 0, 0, 1, 1, 0, \dots, 0), a_9 = (0, 0, 1, 0, 1, 0, \dots, 0), a_{10} = (0, 0, 1, 1, 1, 0, \dots, 0).$ Then the subgraph induced by K in I(R) contains $K_{3,7}$ as a subgraph and by Lemma 4, $\overline{g}(I(R)) \geq 3$.

Theorem 11. Let R be a finite commutative ring. Then I(R) is not a split graph for $n \geq 4$, where n is the number of distinct maximal ideals of R.

Proof. By the structure of I(R), I(R) contains K_n for $n \geq 4$. Consider an vertex a of I(R) that is not in $V(K_n)$, which has 1 in the i, j th places and 0 in the remaining places. This vertex must adjacent with the vertex b that has 0 in the i, j th places and 1 in the remaining places. Hence the remaining vertices cannot form an independent set. Hence the theorem.

Theorem 12. Let R be a finite commutative ring. Then I(R) is a ring graph if and only if n = 3, where n is the number of distinct maximal ideals of R.

Proof. Assume that I(R) is a ring graph. Since every ring graph is planar it is enough to consider whether I(R) is a ring graph for each ring in Theorem 6.

When n = 3, I(R) has rank(I(R)) = frank(I(R)). Hence I(R) is a ring graph. The converse is obvious.

4. Book thickness of I(R)

A standard n-book is formed by joining n half-planes, called pages, together at a common line, called spine. When embedding a graph in a book, the vertices are placed along the spine. Each edge is embedded on a single page of the book so that no two edges cross each on a page. The book thickness of a graph G is the smallest n, for which G has an n-book embedding. Yannakakis [10] has shown that all planar graphs have book thickness at most 4.




Figure 5. Book embedding of $I(R_1 \times R_2 \times R_3)$.

Theorem 13 [5, Theorem 3.4]. The book thickness of the complete graph K_n is equal to $\lceil n/2 \rceil$, when $n \geq 4$.

Theorem 14 [5, Theorem 2.5]. A graph has book thickness one if and only if it is outer planar.

Theorem 15 [5, Theorem 2.5]. The book thickness of a graph is at most two if and only if it is a subgraph of a planar graph that has a Hamiltonian cycle.

Now we classify all rings such that the book thickness of idempotent graph is 1.

Theorem 16. Let R be a finite commutative ring. Then the book thickness of I(R) is 1 if and only if $n \leq 3$, where n is the number of distinct maximal ideals of R.

Proof. The proof is clear by Theorem 6 and Theorem 14.

Now we characterize all finite commutative rings R such that the book thickness of the idempotent graph is 3.

Theorem 17. Let R be a finite commutative ring. The book thickness of I(R) is 3 if and only if R has exactly 4 distinct maximal ideals.

Proof. Suppose R has exactly 4 distinct maximal ideals, then by Theorem 7, we know that $R = R_1 \times R_2 \times R_3 \times R_4$, which is toroidal. By Theorem 15, one can note that, a two page book embedding is corresponding to a planar structure. Hence a toroidal graph has book thickness at least three. But the three pages of I(R) are represented by the sets of dashed and solid edges above and below the spine in Figure 6 (For a_i , $1 \le i \le 14$ refer Figure 2). Hence the book thickness of I(R) is 3.

Conversely assume that the book thickness of I(R) is 3. Suppose that R has at least 5 distinct maximal ideals. Let $J = \{a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_{10}, a_{11}, a_{13}, a_{18}\}$ be a subset of E in Theorem 8. Then, the subgraph induced by J in I(R) must contain a subdivision (each edge should be subdivided at most once) of K_7 . By [11, Theorem 3.1], we come to know that, book thickness of K_n does not change, even though we subdivide its edges at most once. Hence by Theorem 13, the book thickness of I(R) is at least 4, a contradiction.

Corollary 18. There exists no finite commutative ring R, for which book thickness of I(R) is 2.




Figure 6. Book embedding of $I(R_1 \times R_2 \times R_3 \times R_4)$.

Acknowledgment

The support for this research work is provided by MANF programme (201718-MANF-2017-18-TAM-82372) of University Grants Commission, Government of India for the first author.

References

- [1] S. Akbari, M. Habibi, A. Majidinya and R. Manaviyat, On the Idempotent Graph of a Ring, J. Algebra 12 (2013) 1–14. doi:10.1142/S0219498813500035
- [2] M.F. Atiyah and I.G. Macdonald, Introduction to Commutative Algebra (Addison Wiley Publishing Company, London, 1969).
- [3] I. Beck, Coloring of commutative rings, J. Algebra 116 (1988) 208–226. doi:10.1016/0021-8693(88)90202-5
- [4] J.A. Bondy and U.S.R. Murty, Graph Theory and its Applications (American Elsevier, New York, 1976).
- [5] F.R. Bernhart and P.C. Kainen, The book thickness of a graph, J. Comb. Theory, Ser. B 27 (1979) 320–331.
 doi:10.1016/0095-8956(79)90021-2
- [6] S. Kavitha and R. Kala, On the genus of graphs from commutative rings, AKCE Int. J. Graphs Combin. 14 (2017) 27–34. doi:10.1016/j.akcej.2016.11.006
- [7] T.Y. Lam, A First Course in Non-commutative Rings (Springer-Verlag, New York, 2001).
- [8] M.M. Sysło, Characterizations of Outer planar graphs, Discrete Math. 26 (1979) 47–53.
 doi:10.1016/0012-365X(79)90060-8
- [9] A. Mallika and R. Kala, Nilpotent graphs with crosscap at most two, AKCE Int. J. Graphs Combin. 15 (2018) 229–237.
 doi:10.1016/j.akcej.2017.11.006
- [10] M. Yannakakis, Embedding planar graphs in four pages, J. Comput. Syst. Sci. 38 (1989) 36–67. doi:10.1016/0022-0000(89)90032-9
- [11] R. Blankenship and B. Oporowski, Drawing subdivisions of complete and complete bipartite graphs on books (Tech Rep. 1994-4, Department of Mathematics, Louisiana State University, 1999).
- [12] T. McKenzie and S. Overbay, Book thickness of toroidal zero-divisor graphs, Africa Matematika 28 (2017) 823–830. doi:10.35834/mjms/1449161362
- [13] T. Endo, The page number of toroidal graph is at most seven, Discrete Math. 175 (1997) 87–96. doi:10.1016/S0012-365X(96)00144-6
- [14] A.T. White, Graphs, Groups and Surfaces (North-Holland-Amsterdam, 1984).

Received 9 March 2020 Revised 16 March 2020 Accepted 23 April 2020