Discussiones Mathematicae General Algebra and Applications 42 (2022) 51–96 https://doi.org/10.7151/dmgaa.1380

STONE COMMUTATOR LATTICES AND BAER RINGS

Claudia Mureşan

University of Bucharest Faculty of Mathematics and Computer Science

e-mail: c.muresan@yahoo.com cmuresan@fmi.unibuc.ro claudia.muresan@unibuc.ro

Abstract

In this paper, we transfer Davey's characterization for κ -Stone bounded distributive lattices to lattices with certain kinds of quotients, in particular to commutator lattices with certain properties, and obtain related results on prime, radical, complemented and compact elements, annihilators and congruences of these lattices. We then apply these results to certain congruence lattices, in particular to those of semiprime members of semidegenerate congruence-modular varieties, and use this particular case to transfer Davey's Theorem to commutative unitary rings.

Keywords: (strongly) Stone lattice, commutator lattice, annihilator, modular commutator, Baer ring.

2010 Mathematics Subject Classification: Primary: 06B10; Secondary: 06D22, 08A30, 08B10.

1. INTRODUCTION

We shall refer to [6, Theorem 1] as *Davey's Theorem*. Given an arbitrary infinite cardinality κ , Davey's Theorem provides a characterization for κ -Stone bounded distributive lattices: those bounded distributive lattices with the property that the annihilators of their subsets of cardinality at most κ are principal ideals generated by elements from their Boolean center.

It turns out that commutator lattices with certain properties, in particular congruence lattices of semiprime algebras from semi-degenerate congruencemodular varieties, satisfy the equivalences from Davey's Theorem; moreover, changing the cardinalities in those equivalent conditions to any nonzero value produces more properties equivalent to those conditions; furthermore, by restricting the values of these cardinalities, we obtain a generalization of this equivalence result to a generalization of commutator lattices.

To prove this, we first transfer Davey's Theorem to bounded lattices from certain quotients of theirs which are distributive, or frames, or a generalization of frames, and then apply this transfer to commutator lattices satisfying certain conditions and a certain quotient of such commutator lattices.

Then we apply the previous result to the ideal lattices of commutative unitary rings, from which we transfer this result to the elements of these rings.

2. Definitions and notations

We shall denote by \mathbb{N} the set of the natural numbers and by $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$. For any set S, |S| will denote the cardinality of S.

Throughout this paper, all algebras shall be nonempty and, unless there is danger of confusion, they will be designated by their underlying sets. By trivial algebra we mean one-element algebra. Recall that a variety \mathcal{V} is said to be semidegenerate iff no nontrivial algebra in \mathcal{V} has trivial subalgebras. For any algebra A, $(\operatorname{Con}(A), \vee, \cap, \Delta_A, \nabla_A)$ shall be the bounded lattice of the congruences of A, with the exception of the case when A is a lattice-ordered algebra, mentioned below, which produces no danger of confusion in what follows; for any $X \subseteq A^2$ and any $a, b \in A$, $Cg_A(X)$ shall be the congruence of A generated by X and we will denote $Cg_A(a,b) = Cg_A(\{(a,b)\})$; the set of the principal congruences of A, that is the compact elements of the lattice $\operatorname{Con}(A)$, are exactly the finitely generated congruences of A. For any $\theta \in \operatorname{Con}(A)$, $p_{\theta} : A \to A/\theta$ will be the canonical surjection. For any algebra L having a lattice reduct (in particular for any commutator lattice L), $\operatorname{Con}(L)$ will denote the set of the congruences of the lattice reduct of L.

Now let L be an arbitrary lattice. We denote by $\operatorname{Cp}(L)$, $\operatorname{Mi}(L)$ and $\operatorname{Smi}(L)$ the sets of the compact, the meet-irreducible and the strictly meet-irreducible elements of L, respectively. Recall that L is said to be *compact* iff $\operatorname{Cp}(L) = L$ and L is said to be *algebraic* iff each of its elements is a join of compact elements. Note that, if L is compact, then the join of any nonempty $U \subseteq L$ equals the join of a finite subset of U, and that, if L has finite length, then L is compact, thus Lis algebraic. Note that, if L has a 1, then $1 \notin \operatorname{Smi}(L)$, because $1 = \bigwedge \emptyset = \bigwedge \{x \in L \mid 1 < x\}$. For each $a \in \operatorname{Smi}(L)$, we shall denote by $a^+ = \bigwedge \{x \in L \mid a < x\}$ the unique successor of a in L. If L has a 1, then we shall denote by Max_L the set of the maximal elements of the ordered set $(L \setminus \{1\}, \leq)$. For any algebra A, $\operatorname{Max}_{\operatorname{Con}(A)}$ will simply be denoted by $\operatorname{Max}(A)$. Filt(L) and $\operatorname{Id}(L)$ shall be the bounded lattices of the filters and ideals of L, respectively, and $\operatorname{PId}(L)$ will be the bounded sublattice of $\operatorname{Id}(L)$ of the principal ideals of L. Note that a filter of L is principal iff it has a minimum and an ideal of L is principal iff it has a maximum. Recall, also, that any class of a congruence of L is a convex sublattice of L, so it has a unique writing as the intersection between a filter and an ideal of L.

Let $U \subseteq L$ and $a, b \in L$, arbitrary. We denote by $(U]_L$ and $[U)_L$ the ideal and the filter of L generated by U, respectively, by $(a]_L = (\{a\}]_L$ and $[a)_L = [\{a\})_L$ and, to avoid overlapping with the classical notation for the commutator operation in commutator lattices (see Section 3), $[a, b]_L = [a)_L \cap (b]_L$ will be the notation for intervals; if L is the chain of natural numbers with the natural order, then we denote $\overline{a, b} = [a, b]_L$. Note that, for any lattice M and any surjective lattice morphism $f : L \to M$, the map $I \mapsto f(I)$ is a complete lattice morphism from $\mathrm{Id}(L)$ to $\mathrm{Id}(M)$ that satisfies $f((U]_L) = (f(U)]_M$; in particular, for any $\theta \in \mathrm{Con}(L)$, we have $(U]_L/\theta = (U/\theta]_{L/\theta}$, so $(a]_L/\theta = (a/\theta)_{L/\theta}$.

Ann_L(a) and Ann_L(U) shall be the annihilator of a and U in L, respectively: Ann_L(a) = { $x \in L \mid x \land a = 0$ } and Ann_L(U) = $\bigcap_{u \in U} \operatorname{Ann}_L(u)$. We will denote by $\mathcal{A}nn(L) = \{\operatorname{Ann}_L(U) \mid U \subseteq L\}$, PAnn(L) = { $\operatorname{Ann}_L(a) \mid a \in L\}$, P2Ann(L) = { $\operatorname{Ann}_L(\operatorname{Ann}_L(a)) \mid a \in L\}$ and 2Ann(L) = { $\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \mid U \subseteq L\}$. The following notations will also be useful: let $\mathcal{A}nn_{<\infty}(L) = \{\operatorname{Ann}_L(U) \mid U \subseteq L, |U| < \aleph_0\}$ and 2Ann_{< ∞}(L) = { $\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \mid U \subseteq L, |U| < \aleph_0\}$ and, if κ is a cardinality, let $\mathcal{A}nn_{\kappa}(L) = \{\operatorname{Ann}_L(U) \mid U \subseteq L, |U| \leq \kappa\}$ and 2Ann_{κ}(L) = { $\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \mid U \subseteq L, |U| \leq \kappa\}$.

 $\mathcal{B}(L)$ will denote the set of the complemented elements of the bounded lattice L, regardless of whether L is distributive. Unless mentioned otherwise, we shall denote by \neg the complementation in every Boolean algebra.

Recall that the bounded lattice L is said to be *Stone*, respectively *strongly Stone*, iff, for all $a \in L$, respectively all $U \subseteq L$, there exists an $e \in \mathcal{B}(L)$ such that $\operatorname{Ann}_{L}(a) = (e]_{L}$, respectively $\operatorname{Ann}_{L}(U) = (e]_{L}$. Also, for any cardinality κ , L is said to be κ -*Stone* iff, for all $U \subseteq L$ with $|U| \leq \kappa$, there exists an $e \in \mathcal{B}(L)$ such that $\operatorname{Ann}_{L}(U) = (e]_{L}$.

Remember that L is called a *frame* iff L is complete and the meet in L is completely distributive w.r.t. the join.

3. The theorem we are going to transfer to commutator Lattices, then to commutative unitary rings

Throughout this section, L will be a bounded lattice. We shall use the following notations for these conditions on L, where κ is an arbitrary cardinality:

- $(1)_{\kappa,L}$ L is a κ -Stone lattice; $(1)_{<\infty,L} \quad \mathcal{A}nn_{<\infty}(L) \subseteq \{(e|_L \mid e \in \mathcal{B}(L)\};\$ L is a strongly Stone lattice; $(1)_{L}$ L is a Stone lattice and $\mathcal{B}(L)$ is a κ -complete Boolean sublattice of L; $(2)_{\kappa,L}$ $(2)_{<\infty,L}$ L is a Stone lattice and $\mathcal{B}(L)$ is a Boolean sublattice of L; L is a Stone lattice and $\mathcal{B}(L)$ is a complete Boolean sublattice of L; $(2)_{L}$ P2Ann(L) is a κ -complete Boolean sublattice of Id(L) such that $(3)_{\kappa,L}$ $a \mapsto \operatorname{Ann}_L(\operatorname{Ann}_L(a))$ is a lattice morphism from L to P2Ann(L); $(3)_{<\infty,L}$ P2Ann(L) is a Boolean sublattice of Id(L) such that $a \mapsto \operatorname{Ann}_L(\operatorname{Ann}_L(a))$ is a lattice morphism from L to P2Ann(L); $(3)_{L}$ P2Ann(L) is a complete Boolean sublattice of Id(L) such that $a \mapsto \operatorname{Ann}_L(\operatorname{Ann}_L(a))$ is a lattice morphism from L to P2Ann(L); for all $a, b \in L$, $\operatorname{Ann}_L(a \wedge b) = (\operatorname{Ann}_L(a) \cup \operatorname{Ann}_L(b)]_L$, and $(4)_{\kappa,L}$ $2\operatorname{Ann}_{\kappa}(L) \subseteq \operatorname{PAnn}(L);$ $(4)_{<\infty,L}$ for all $a, b \in L$, $\operatorname{Ann}_L(a \wedge b) = (\operatorname{Ann}_L(a) \cup \operatorname{Ann}_L(b)]_L$, and $2\operatorname{Ann}_{<\infty}(L) \subseteq \operatorname{PAnn}(L);$ for all $a, b \in L$, $\operatorname{Ann}_L(a \wedge b) = (\operatorname{Ann}_L(a) \cup \operatorname{Ann}_L(b)]_L$, and $(4)_{L}$ $2\operatorname{Ann}(L) \subseteq \operatorname{PAnn}(L);$ $(iv)_L$ for all $a, b \in L$, $\operatorname{Ann}_L(a \wedge b) = (\operatorname{Ann}_L(a) \cup \operatorname{Ann}_L(b)]_L$; for each $U \subseteq L$ with $|U| \leq \kappa$, $(\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L(\operatorname{Ann}_L(U))]_L = L$; $(5)_{\kappa,L}$
- $(5)_{\kappa,L}$ for each finite $U \subseteq L$, $(\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L(\operatorname{Ann}_L(U))]_L = L;$
- (5)_L for each $U \subseteq L$, $(\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L(\operatorname{Ann}_L(U))]_L = L$.

Of course, annihilators are nonempty, since each of them contains 0. Note also that, for any $U \subseteq V \subseteq L$, we have $\operatorname{Ann}_L(V) \subseteq \operatorname{Ann}_L(U)$, hence $\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \subseteq \operatorname{Ann}_L(\operatorname{Ann}_L(V))$.

Since $0, 1 \in \mathcal{B}(L)$, we have $\operatorname{Ann}_L(\emptyset) = L = (1]_L = \operatorname{Ann}_L(0) \in \operatorname{PAnn}(L) \cap \{(e]_L \mid e \in \mathcal{B}(L)\} \text{ and } \operatorname{Ann}_L(\operatorname{Ann}_L(\emptyset)) = \operatorname{Ann}_L(L) = \{0\} = (0]_L = \operatorname{Ann}_L(1) \in \operatorname{PAnn}(L) \cap \{(e]_L \mid e \in \mathcal{B}(L)\}, \text{ hence conditions } (1)_{0,L} \text{ and } (5)_{0,L} \text{ are trivially sat$ $isfied; we may also note that <math>\operatorname{PAnn}(L) = \mathcal{Ann}_1(L) \text{ and } \operatorname{P2Ann}(L) = 2\operatorname{Ann}_1(L).$

Clearly, if L is distributive, then every annihilator of L is an ideal of L.

Remark 3.1. If *L* is a frame, then every annihilator of *L* is a principal ideal of *L*. Indeed, if *L* is a frame, then, clearly, for all $U \subseteq L$, $\bigvee \operatorname{Ann}_L(U) \in \operatorname{Ann}_L(U)$, hence the ideal $\operatorname{Ann}_L(U)$ is principal.

Of course, if $\mathcal{B}(L)$ is a distributive sublattice of L, in particular if L is distributive, then $\mathcal{B}(L)$ is a Boolean sublattice of L.

Note that, for any bounded lattice L, any cardinalities $\kappa \leq \mu$ and any $i \in \overline{1, 5}$:

- $(4)_{\kappa,L}$ implies $(iv)_L$;
- $(i)_{\mu,L}$ implies $(i)_{\kappa,L}$, hence, if the converse implication holds, as well, then $(i)_{\kappa,L}$ is equivalent to $(i)_{\nu,L}$ for any cardinality ν with $\kappa \leq \nu \leq \mu$;

- $(i)_{<\infty,L}$ is equivalent to $(i)_{\nu,L}$ being valid for all finite cardinalities ν ;
- $(i)_L$ is equivalent to $(i)_{\nu,L}$ being valid for all cardinalities ν .

For any nonempty family $(U_i)_{i\in I}$ of subsets of L, clearly $\operatorname{Ann}_L(\bigcup_{i\in I} U_i) = \bigcap_{i\in I} \operatorname{Ann}_L(U_i)$. For any family $(a_i)_{i\in I} \subseteq L$ having a meet in L, we have $\bigcap_{i\in I} (a_i]_L = (\bigwedge_{i\in I} a_i]_L$. Trivially, if L is strongly Stone, then L is Stone, and, by the above, the converse holds if $\operatorname{Ann}(L) = \operatorname{PAnn}(L)$ or $\mathcal{B}(L)$ is closed w.r.t. arbitrary meets; in particular, $(2)_L$ implies $(1)_L$.

If L is distributive, then, for any $n \in \mathbb{N}^*$ and any $u_1, \ldots, u_n \in L$, $\operatorname{Ann}_L(\{u_1, \ldots, u_n\}) = \operatorname{Ann}_L(u_1 \lor \cdots \lor u_n) \in \operatorname{PAnn}(L)$, so $\operatorname{Ann}_{<\infty}(L) = \operatorname{PAnn}(L)$, hence $(1)_{1,L}$ is equivalent to $(1)_{<\infty,L}$, that is L is a Stone lattice iff it satisfies $(1)_{<\infty,L}$, and it immediately follows that $\operatorname{Ann}_L(U) = \operatorname{Ann}_L((U]_L)$ for all $U \subseteq L$ and thus, for any family $(I_k)_{k \in K}$ of ideals of L, $\operatorname{Ann}_L(\bigvee_{k \in K} I_k) = \operatorname{Ann}_L((\bigcup_{k \in K} I_k]_L) = \operatorname{Ann}_L(\bigcup_{k \in K} I_k) = \bigcap_{k \in K} \operatorname{Ann}_L(I_k)$.

Remark 3.2. Let κ be a nonzero cardinality.

If $(x_i)_{i \in I} \subseteq L$ such that $\bigvee_{i \in I} x_i \in L$ and $x \land (\bigvee_{i \in I} x_i) = \bigvee_{i \in I} (x \land x_i)$ for all $x \in L$, then clearly $\operatorname{Ann}_L(\{x_i \mid i \in I\}) = \operatorname{Ann}_L(\bigvee_{i \in I} x_i) \in \operatorname{PAnn}(L)$.

Thus, if L is closed w.r.t. the joins of all families of elements of cardinality at most κ and has the meet distributive w.r.t. the joins of families of cardinalities at most κ , then $\operatorname{Ann}_{L}(U) = \operatorname{Ann}_{L}(\bigvee U)$ for all $U \subseteq L$ with $|U| \leq \kappa$, thus $\operatorname{Ann}_{\kappa}(L) = \operatorname{PAnn}(L)$.

Also, if L is a frame, then $\operatorname{Ann}_L(U) = \operatorname{Ann}_L(\bigvee U)$ for all $U \subseteq L$, thus $\mathcal{Ann}(L) = \operatorname{PAnn}(L)$.

If $\mathcal{A}nn_{\kappa}(L) = \operatorname{PAnn}(L)$, in particular if L is closed w.r.t. the joins of families of elements of cardinality at most κ and has the meet distributive w.r.t. such joins, then $(1)_{1,L}$ is equivalent to $(1)_{\kappa,L}$ (thus to $(1)_{\lambda,L}$ for any nonzero cardinality $\lambda \leq \kappa$), and $\operatorname{PAnn}_{\kappa}(L) = \operatorname{P2Ann}(L)$, hence $\operatorname{P2Ann}(L) \subseteq \operatorname{PAnn}(L)$ iff $\operatorname{PAnn}_{\kappa}(L) \subseteq \operatorname{PAnn}(L)$, thus $(4)_{1,L}$ is equivalent to $(4)_{\kappa,L}$ (thus to $(4)_{\lambda,L}$ for any nonzero cardinality $\lambda \leq \kappa$).

If $\mathcal{A}nn(L) = \operatorname{PAnn}(L)$, in particular if L is a frame, then $(1)_{1,L}$ is equivalent to $(1)_L$ (thus to $(1)_{\lambda,L}$ for any nonzero cardinality λ), and $2\operatorname{Ann}(L) = \operatorname{P2Ann}(L) \subseteq \mathcal{A}nn(L) = \operatorname{PAnn}(L)$, thus the second part of condition $(4)_L$ is satisfied, which means that $(4)_L$ is equivalent to $(iv)_L$ and thus to $(4)_{\lambda,L}$ for any nonzero cardinality λ .

As an example, note that any Boolean lattice is Stone, because, if L is Boolean, then $\operatorname{Ann}_L(e) = (\neg e]_L$ for all $e \in L$, thus, by the above, any complete Boolean lattice is strongly Stone.

Theorem 3.3. (i) If L is a bounded distributive lattice, then the conditions $(1)_{\kappa,L}, (2)_{\kappa,L}, (3)_{\kappa,L}, (4)_{\kappa,L}$ and $(5)_{\kappa,L}$ are equivalent for any nonzero cardinality κ .

- (ii) If L is a bounded distributive lattice, then conditions $(1)_L$, $(2)_L$, $(3)_L$, $(4)_L$ and $(5)_L$ are equivalent.
- (iii) Let m be a nonzero cardinality. If L is a bounded distributive lattice having $\mathcal{A}nn_m(L) = \text{PAnn}(L)$, in particular if the bounded lattice L is closed w.r.t. the joins of all families of elements of cardinality at most m and has the meet distributive w.r.t. such joins then, for any $h, i \in \overline{1,5}$ and any nonzero cardinality $\kappa \leq m$, conditions $(h)_{\kappa,L}$ and $(i)_{<\infty,L}$ are equivalent, in particular the Boolean center of L is m-complete and L is Stone iff it is m-Stone.
- (iv) If L is a bounded distributive lattice with Ann(L) = PAnn(L), in particular if L is a frame, then, for any $h, i, j \in \overline{1,5}$ and any nonzero cardinality κ , conditions $(iv)_L$, $(h)_{\kappa,L}$, $(i)_{<\infty,L}$ and $(j)_L$ are equivalent, in particular the Boolean center of L is complete and L is Stone iff it is strongly Stone.

Proof. (i) is [6, Theorem 1] for κ infinite and part of Remark 3.2 for κ finite, and it clearly implies (ii).

(iii), (iv) By (i) and Remark 3.2.

Definition 3.4 [2, 5, 18]. Let $[\cdot, \cdot]$ be a binary operation on L. The algebra $(L, \lor, \land, [\cdot, \cdot], 0, 1)$ (which we shall also denote, simply, by $(L, [\cdot, \cdot])$) is called a *commutator lattice* and the operation $[\cdot, \cdot]$ is called *commutator* iff $(L, \lor, \land, 0, 1)$ is a complete lattice with lattice bounds 0 and 1 and, for all $x, y \in L$ and any family $(y_i)_{i \in I} \subseteq L$:

- $[x,y] = [y,x] \le x \land y$ ([\cdot, \cdot] is commutative and smaller than its arguments);
- $[x, \bigvee_{i \in I} y_i] = \bigvee_{i \in I} [x, y_i] ([\cdot, \cdot] \text{ is completely distributive w.r.t. the join}).$

The latter condition in Definition 3.4 implies that the commutator $[\cdot, \cdot]$ is distributive w.r.t. the (binary) join (in each argument, since it is commutative), thus $[\cdot, \cdot]$ is order-preserving in each argument.

Remark 3.5. For any complete lattice L, we have the equivalence: $(L, [\cdot, \cdot])$ is a commutator lattice with $[\cdot, \cdot] = \wedge$ iff L is a frame.

Let $(L, \lor, \land, [\cdot, \cdot], 0, 1)$ be a commutator lattice. We call p a prime element of L iff $p \in L \setminus \{1\}$ and, for all $a, b \in L$, if $[a, b] \leq p$, then $a \leq p$ or $b \leq p$. We denote by Spec_L the set of the prime elements of L. Note that, if $[\cdot, \cdot] = \land$, then Spec_L is the set of the meet-prime elements of L and L is distributive (actually a frame, by Remark 3.5), hence $\operatorname{Spec}_L = \operatorname{Mi}(L) \setminus \{1\} \supseteq \operatorname{Smi}(L)$.

For any $x \in L$, we denote by $V(x) = [x)_L \cap \operatorname{Spec}_L$, by $\rho(x) = \bigwedge V(x) = \bigwedge \{p \in \operatorname{Spec}_L \mid x \leq p\}$ and by $R(L) = \{\rho(x) \mid x \in L\}$. We call $\rho(x)$ the radical of x, and the elements of R(L) radical elements of L. Clearly, $\operatorname{Spec}_L \subseteq R(L) = \{x \in L \mid \rho(x) = x\}$.

Recall that all elements of an algebraic lattice are meets of strictly meetirreducible elements, thus, if L is algebraic and $[\cdot, \cdot] = \wedge$, then R(L) = L; see also Remarks 5.3 and 5.11 and Proposition 5.15, (iii), below.

Example 3.6 [1, 7]. If \mathcal{V} is a congruence-modular variety, A is a member of \mathcal{V} and $[\cdot, \cdot]_A$ is the (modular) commutator of A, then $(\operatorname{Con}(A), \lor, \cap, [\cdot, \cdot]_A, \Delta_A, \nabla_A)$ is a commutator lattice.

Let A be an arbitrary member of a congruence-modular variety \mathcal{V} . Then we will denote the modular commutator of A as above and the set $\operatorname{Spec}_{\operatorname{Con}(A)}$ of the prime elements of the commutator lattice $(\operatorname{Con}(A), \lor, \cap, [\cdot, \cdot]_A, \Delta_A, \nabla_A)$, called *prime congruences* of A, by $\operatorname{Spec}(A)$. The elements of $R(\operatorname{Con}(A))$ are called *radical congruences* of A. Recall that A is said to be *semiprime* iff Δ_A is a radical congruence of A.

Recall that, if \mathcal{V} is congruence-distributive, then \mathcal{V} has no skew congruences and the commutator $[\cdot, \cdot]_A$ coincides to the intersection of congruences. If $[\cdot, \cdot]_A$ equals the intersection, then $[\theta, \nabla_A]_A = \theta \cap \nabla_A = \theta$ for all $\theta \in \text{Con}(A)$, and, by the above, A is congruence-distributive and, moreover, Con(A) is a frame, and we have $\text{Smi}(\text{Con}(A)) \subseteq \text{Mi}(\text{Con}(A)) \setminus {\nabla_A} = \text{Spec}(A)$, so that R(Con(A)) =Con(A) since the lattice Con(A) is algebraic, in particular A is semiprime.

If \mathcal{V} is semi-degenerate, then \mathcal{C} has no skew congruences [7, Theorem 8.5, p. 85], ∇_A is a compact congruence of A, $[\theta, \nabla_A]_A = \theta$ for all $\theta \in \text{Con}(A)$, any maximal congruence of A is prime and each proper congruence of A is included in a prime congruence [1, Theorem 5.3]; see also Lemma 5.2 below.

4. TRANSFERRING CONDITIONS $(i)_{\kappa,\cdot}$ BETWEEN BOUNDED LATTICES AND THEIR QUOTIENTS, AND RELATED RESULTS

Throughout this section, M shall be an arbitrary bounded lattice and $\theta \in Con(M)$, unless mentioned otherwise.

Remark 4.1. Let $x \in M$ and $U \subseteq M$. Then, clearly, $x \in \operatorname{Ann}_M(U)$ implies $x/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$, hence $\operatorname{Ann}_M(U)/\theta \subseteq \operatorname{Ann}_{M/\theta}(U/\theta)$.

Lemma 4.2. If M is a bounded lattice and a $\theta \in Con(M)$ has $0/\theta = \{0\}$, then, for all $x \in M$ and all $U, V \subseteq M$:

- (i) $x/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$ iff $x \in \operatorname{Ann}_M(U)$ iff $x/\theta \subseteq \operatorname{Ann}_M(U)$, and $x/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta))$ iff $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ iff $x/\theta \subseteq \operatorname{Ann}_M(\operatorname{Ann}_M(U))$;
- (ii) $\operatorname{Ann}_{M}(U)/\theta = \operatorname{Ann}_{M/\theta}(U/\theta)$ and $\operatorname{Ann}_{M}(\operatorname{Ann}_{M}(U))/\theta = \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta));$

(iii) $U/\theta \subseteq \operatorname{Ann}_M(V)/\theta$ iff $U \subseteq \operatorname{Ann}_M(V)$, and $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_M(V)/\theta$ iff $\operatorname{Ann}_M(U) = \operatorname{Ann}_M(V)$.

Proof. (i) and (ii) If $x/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$, then, for all $u \in U$, we have $x \wedge u \in 0/\theta = \{0\}$, thus $x \in \operatorname{Ann}_M(U)$, so $x/\theta \in \operatorname{Ann}_M(U)/\theta$, hence $\operatorname{Ann}_{M/\theta}(U/\theta) \subseteq \operatorname{Ann}_M(U)/\theta$. We have the converse implication and inclusion from Remark 4.1, therefore $\operatorname{Ann}_{M/\theta}(U/\theta) = \operatorname{Ann}_M(U)/\theta$ and $x \in \operatorname{Ann}_M(U)$ iff $x/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$, so that: $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ iff $x/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_M(U)/\theta) = \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_M(U)/\theta)$.

Clearly, if $x/\theta \subseteq \operatorname{Ann}_M(U)$, then $x \in \operatorname{Ann}_M(U)$, while, if $x/\theta \subseteq \operatorname{Ann}_M(\operatorname{Ann}_M(U))$, then $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$. By the above, for any $y \in x/\theta$, we have: $x \in \operatorname{Ann}_M(U)$ iff $x/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$ iff $y/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$ iff $y \in \operatorname{Ann}_M(U)$, and, similarly, $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ iff $x/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta))$ iff $y/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta))$ iff $y \in \operatorname{Ann}_M(U)$. Therefore $x \in \operatorname{Ann}_M(U)$ implies $x/\theta \subseteq \operatorname{Ann}_M(U)$, while $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ implies $x/\theta \subseteq \operatorname{Ann}_M(U)$, while $x \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ implies $x/\theta \subseteq \operatorname{Ann}_M(U)$.

(iii) By (ii), for all $u \in U$, we have: $u/\theta \in \operatorname{Ann}_M(V)/\theta = \operatorname{Ann}_{M/\theta}(V/\theta)$ iff $u \in \operatorname{Ann}_M(V)$, hence the first equivalence, therefore: $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_M(V)/\theta$ iff $\operatorname{Ann}_M(U)/\theta \subseteq \operatorname{Ann}_M(V)/\theta$ and $\operatorname{Ann}_M(V)/\theta \subseteq \operatorname{Ann}_M(U)/\theta$ iff $\operatorname{Ann}_M(U) \subseteq \operatorname{Ann}_M(V)$ and $\operatorname{Ann}_M(V) \subseteq \operatorname{Ann}_M(U)$ iff $\operatorname{Ann}_M(V)$.

Lemma 4.3. Let M be a bounded lattice, $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and κ a cardinality. Then:

- (i) the maps $P \mapsto P/\theta$ from: Ann(M) to $Ann(M/\theta)$, $Ann_{\kappa}(M)$ to $Ann_{\kappa}(M/\theta)$, PAnn(M) to PAnn(M/ θ), 2Ann(M) to 2Ann(M/ θ), 2Ann_{κ}(M) to 2Ann_{κ} (M/ θ), respectively P2Ann(M) to P2Ann(M/ θ), are order isomorphisms;
- (ii) $\mathcal{A}nn_{\kappa}(M/\theta) = \operatorname{PAnn}(M/\theta) \text{ iff } \mathcal{A}nn_{\kappa}(M) = \operatorname{PAnn}(M); \mathcal{A}nn(M/\theta) = \operatorname{PAnn}(M/\theta) \text{ iff } \mathcal{A}nn(M) = \operatorname{PAnn}(M); \operatorname{2Ann}_{\kappa}(M/\theta) \subseteq \operatorname{PAnn}(M/\theta) \text{ iff } \operatorname{2Ann}_{\kappa}(M) \subseteq \operatorname{PAnn}(M); \operatorname{2Ann}(M/\theta) \subseteq \operatorname{PAnn}(M) \subseteq \operatorname{PAnn}(M);$
- (iii) for all $U \subseteq M$: $\operatorname{Ann}_M(U) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(U/\theta) \in \operatorname{Id}(M/\theta)$, and $\operatorname{Ann}_M(\operatorname{Ann}_M(U)) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta)) \in \operatorname{Id}(M/\theta)$;
- (iv) $\mathcal{A}nn(M) \subseteq \mathrm{Id}(M)$ iff $\mathcal{A}nn(M/\theta) \subseteq \mathrm{Id}(M/\theta)$; $\mathrm{PAnn}(M) \subseteq \mathrm{Id}(M)$ iff $\mathrm{PAnn}(M/\theta) \subseteq \mathrm{Id}(M/\theta)$; $\mathrm{P2Ann}(M) \subseteq \mathrm{Id}(M)$ iff $\mathrm{P2Ann}(M/\theta) \subseteq \mathrm{Id}(M/\theta)$;
- (v) for all $U, V \subseteq M$ such that $\operatorname{Ann}_M(U), \operatorname{Ann}_M(V) \in \operatorname{Id}(M)$, we have, in $\operatorname{Id}(M)$ and $\operatorname{Id}(M/\theta)$: $\operatorname{Ann}_M(U \cap V) = \operatorname{Ann}_M(U) \vee \operatorname{Ann}_M(V) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(U/\theta \cap V/\theta) = \operatorname{Ann}_{M/\theta}(U/\theta) \vee \operatorname{Ann}_{M/\theta}(V/\theta) \in \operatorname{Id}(M/\theta);$
- (vi) if $Ann(M) \subseteq Id(M)$ and $Ann_M(U \cap V) = Ann_M(U) \vee Ann_M(V)$ for all $U, V \subseteq M$, then Ann(M) and $Ann(M/\theta)$ are sublattices of Id(M) and $Id(M/\theta)$, respectively, and the map $P \mapsto P/\theta$ from Ann(M) to $Ann(M/\theta)$ is a lattice isomorphism;

- (vii) for all $a, b \in M$ such that $\operatorname{Ann}_M(a), \operatorname{Ann}_M(b) \in \operatorname{Id}(M)$, we have, in $\operatorname{Id}(M)$ and $\operatorname{Id}(M/\theta)$: $\operatorname{Ann}_M(a \lor b) = \operatorname{Ann}_M(a) \cap \operatorname{Ann}_M(b) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(a/\theta \lor b/\theta) = \operatorname{Ann}_{M/\theta}(a/\theta) \cap \operatorname{Ann}_{M/\theta}(b/\theta) \in \operatorname{Id}(M/\theta)$, and $\operatorname{Ann}_M(a \land b) = \operatorname{Ann}_M(a) \lor \operatorname{Ann}_M(b) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(a/\theta \land b/\theta) = \operatorname{Ann}_{M/\theta}(a/\theta) \lor \operatorname{Ann}_{M/\theta}(b/\theta) \in \operatorname{Id}(M/\theta)$;
- (viii) for all $a, b \in M$ such that $\operatorname{Ann}_M(\operatorname{Ann}_M(a)), \operatorname{Ann}_M(\operatorname{Ann}_M(b)) \in \operatorname{Id}(M)$, we have, in $\operatorname{Id}(M)$ and $\operatorname{Id}(M/\theta)$: $\operatorname{Ann}_M(\operatorname{Ann}_M(a \lor b)) = \operatorname{Ann}_M(\operatorname{Ann}_M(a)) \lor$ $\operatorname{Ann}_M(\operatorname{Ann}_M(b)) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(a/\theta \lor b/\theta)) = \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(a/\theta)) \lor \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(b/\theta)) \in \operatorname{Id}(M/\theta)$, and $\operatorname{Ann}_M(\operatorname{Ann}_M(a \land b)) = \operatorname{Ann}_M(\operatorname{Ann}_M(a)) \cap \operatorname{Ann}_M(\operatorname{Ann}_M(b)) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(a/\theta)) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(a/\theta)) \in \operatorname{Id}(M/\theta)$;
 - (ix) PAnn(M) is a sublattice of Id(M) such that the map $x \mapsto \text{Ann}_M(x)$ is a lattice anti-morphism from M to PAnn(M) iff PAnn(M/ θ) is a sublattice of Id(M/ θ) such that the map $y \mapsto \text{Ann}_{M/\theta}(y)$ is a lattice anti-morphism from M/θ to PAnn(M/ θ), and, if so, then the map $P \mapsto P/\theta$ from PAnn(M) to PAnn(M/ θ) is a lattice isomorphism;
 - (x) P2Ann(M) is a sublattice of Id(M) such that the map $x \mapsto \operatorname{Ann}_M(\operatorname{Ann}_M(x))$ is a lattice morphism from M to P2Ann(M) iff P2Ann(M/ θ) is a sublattice of Id(M/ θ) such that the map $y \mapsto \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(y))$ is a lattice morphism from M/ θ to P2Ann(M/ θ), and, if so, then the map $P \mapsto P/\theta$ from P2Ann(M) to P2Ann(M/ θ) is a lattice isomorphism.

Proof. (i) By Lemma 4.2, (ii), these maps are well defined and surjective; by Lemma 4.2, (iii), they are also injective, hence they are bijective. By Lemma 4.2, (ii), these maps, as well as their inverses, preserve inclusion. Therefore they are order isomorphisms.

(ii) By (i) and Lemma 4.2, (iii).

(iii) From (ii) and the clear fact that $I/\theta \in \mathrm{Id}(M/\theta)$ for any $I \in \mathrm{Id}(M)$, we get the direct implications.

Now assume that $\operatorname{Ann}_{M/\theta}(U/\theta) \in \operatorname{Id}(M/\theta)$, and let $x, y, z \in M$ such that $x, y \in \operatorname{Ann}_M(U)$ and $x \geq z$, so that $x/\theta, y/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$ and $x/\theta \geq z/\theta$, thus $(x \vee y)/\theta, z/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$, hence $x \vee y, z \in \operatorname{Ann}_M(U)$ by Lemma 4.2, (i), therefore $\operatorname{Ann}_M(U) \in \operatorname{Id}(M)$.

Thus $\operatorname{Ann}_M(U) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(U/\theta) \in \operatorname{Id}(M/\theta)$. By Lemma 4.2, (ii), from this we also get that $\operatorname{Ann}_M(\operatorname{Ann}_M(U)) \in \operatorname{Id}(M)$ iff $\operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta)) = \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_M(U)/\theta) \in \operatorname{Id}(M/\theta)$.

(iv) By (iii).

(v) If $\operatorname{Ann}_M(U)$, $\operatorname{Ann}_M(V) \in \operatorname{Id}(M)$, then $\operatorname{Ann}_{M/\theta}(U/\theta)$, $\operatorname{Ann}_{M/\theta}(V/\theta) \in \operatorname{Id}(M/\theta)$ by (iii), so the equivalences in the enunciation follow from Lemma 4.2, (ii), and the fact that the map $I \mapsto I/\theta$ is a lattice morphism from $\operatorname{Id}(M)$ to $\operatorname{Id}(M/\theta)$.

(vi) By (i), (iii) and the fact that, for all $U, V \subseteq M$, $\operatorname{Ann}_M(U \cup V) = \operatorname{Ann}_M(U) \cap \operatorname{Ann}_M(V)$ and the same goes for $U/\theta, V/\theta$ in M/θ .

(vii) and (viii) Similar to the proof of (v).

(ix) By (i), (iii), (iv) and (vii).

(x) By (i), (iii), (iv) and (viii).

Proposition 4.4. Let M be a bounded lattice and $\theta \in Con(M)$ such that $0/\theta = \{0\}$.

- (i) If M/θ is distributive, then $Ann(M) \subseteq Id(M)$ and $Ann_M(U) = Ann_M(U)_M$, so $Ann_M(\bigvee_{k \in K} I_k) = \bigcap_{k \in K} Ann_M(I_k)$ for any $(I_k)_{k \in K} \subseteq Id(M)$.
- (ii) Let κ be a nonzero cardinality. If Ann_κ(M/θ) = PAnn(M/θ), in particular if M/θ is closed w.r.t. the joins of families of elements of cardinality at most κ and has the meet distributive w.r.t. the joins of families of elements of cardinality at most κ, then Ann_κ(M) = PAnn(M), so M is Stone iff M is κ-Stone.

If M and M/θ are closed w.r.t. the joins of families of elements of cardinality at most κ , M/θ has the meet distributive w.r.t. such joins and θ preserves such joins, then $\operatorname{Ann}_M(U) = \operatorname{Ann}_M(\bigvee U)$ for any $U \subseteq M$ with $|U| \leq \kappa$.

(iii) If $Ann(M/\theta) = PAnn(M/\theta)$, in particular if M/θ is a frame, then $Ann(M) = PAnn(M) \subseteq PId(M)$, so M is Stone iff M is strongly Stone.

If M is complete, M/θ is a frame and θ preserves arbitrary joins, then Ann_M(U) = Ann_M($\bigvee U$) for any $U \subseteq M$.

Proof. (i) By Lemma 4.2, (ii) and (iii), $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_{M/\theta}(U/\theta) = \operatorname{Ann}_{M/\theta}((U/\theta)_{M/\theta}) = \operatorname{Ann}_{M/\theta}((U)_M/\theta) = \operatorname{Ann}_M((U)_M)/\theta$, thus $\operatorname{Ann}_M(U) = \operatorname{Ann}_M((U)_M)$, hence the equality for the family of ideals of M.

Also, $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_{M/\theta}(U/\theta) \in \operatorname{Id}(M/\theta)$, so that $\operatorname{Ann}_{M/\theta}(U/\theta) = (\operatorname{Ann}_{M/\theta}(U/\theta)]_{M/\theta} = (\operatorname{Ann}_M(U)/\theta]_{M/\theta} = (\operatorname{Ann}_M(U)]_M/\theta$, thus $(\operatorname{Ann}_M(U)]_M/\theta$ $\subseteq \operatorname{Ann}_M(U)/\theta$, hence $(\operatorname{Ann}_M(U)]_M \subseteq \operatorname{Ann}_M(U)$, therefore $\operatorname{Ann}_M(U) = (\operatorname{Ann}_M(U))_M \in \operatorname{Id}(M)$.

(ii) By Remark 3.2 and Lemma 4.3, (ii), $Ann_{\kappa}(M) = PAnn(M)$.

If an $U \subseteq M$ has $|U| \leq \kappa$, then $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_{M/\theta}(\bigvee(U/\theta)) = \operatorname{Ann}_{M/\theta}(\bigvee(U)/\theta) = \operatorname{Ann}_M(\bigvee U)/\theta$, hence $\operatorname{Ann}_M(U) = \operatorname{Ann}_M(\bigvee U)$ by Lemma 4.2, (iii). (iii) By Remark 3.2 and Lemma 4.3, (ii), $\mathcal{Ann}(M) = \operatorname{PAnn}(M)$.

Additionally, $(\bigvee \operatorname{Ann}_M(U))/\theta = \bigvee (\operatorname{Ann}_M(U)/\theta) = \bigvee \operatorname{Ann}_{M/\theta}(U/\theta) \in \operatorname{Ann}_{M/\theta}(U/\theta) = \operatorname{Ann}_M(U)/\theta$ by Remark 3.1, thus $\bigvee \operatorname{Ann}_M(U) \in \operatorname{Ann}_M(U)$, hence the ideal $\operatorname{Ann}_M(U)$ of M is principal.

As in the proof in (ii), here we obtain that, for any $U \subseteq M$, $\operatorname{Ann}_M(U) = \operatorname{Ann}_M(\bigvee U)$.

60

Remark 4.5. Let $e \in M$. Then $e = \max(e/\theta)$ iff, for all $x \in M$, we have the equivalence: $x/\theta \le e/\theta$ iff $x \le e$.

Indeed, the latter equivalence and the fact that $e \in e/\theta$ imply that $e = \max(e/\theta)$, while, if the latter equality holds and $x/\theta \leq e/\theta$, then $(x \vee e)/\theta = e/\theta$, that is $x \vee e \in e/\theta$, so that $x \vee e \leq \max(e/\theta) = e$, thus $x \leq e$.

Hence, if $e = \max(e/\theta)$, then, for all $U \subseteq M$, we have $U/\theta \subseteq (e]_M/\theta = (e/\theta)_{M/\theta}$ iff $U \subseteq (e]_M$.

Note that Theorem 3.3, (i), relies on the fact that $\mathcal{B}(\mathrm{Id}(D)) = \{(e]_D \mid e \in \mathcal{B}(D)\}$ for any bounded distributive lattice D. Let us see that we can transfer this property from M/θ to M.

Remark 4.6. Clearly, $\mathcal{B}(M)/\theta \subseteq \mathcal{B}(M/\theta)$, thus the map $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is well defined.

Recall from [9] that, by definition, θ has the Boolean Lifting Property (BLP) iff $\mathcal{B}(M)/\theta = \mathcal{B}(M/\theta)$, that is iff the map above is surjective.

Remark 4.7. If $0/\theta = \{0\}$ and $1/\theta = \{1\}$, then, clearly, for any $e, f \in M$: e is a complement of f iff e/θ is a complement of f/θ , thus $e \in \mathcal{B}(M)$ iff $e/\theta \in \mathcal{B}(M/\theta)$, hence $\mathcal{B}(M/\theta) = \mathcal{B}(M)/\theta$ (that is θ has the BLP).

Remark 4.8. If $e = \max(e/\theta)$ for all $e \in \mathcal{B}(M)$, then, by Remark 4.5, for all $e, f \in \mathcal{B}(M)$, we have: $e/\theta = f/\theta$ iff $e/\theta \leq f/\theta$ and $f/\theta \leq e/\theta$ iff $e \leq f$ and $f \leq e$ iff e = f, hence the map $p_{\theta}|_{\mathcal{B}(M)} \colon \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is injective.

Remark 4.9. Clearly, if $\mathcal{B}(M)$ is a sublattice, respectively a Boolean sublattice of M, then $\mathcal{B}(M)/\theta$ is a sublattice, respectively a Boolean sublattice of M/θ .

Since $p_{\theta} : M \to M/\theta$ is a bounded lattice morphism, it follows that, if $\mathcal{B}(M)$ and $\mathcal{B}(M/\theta)$ are sublattices, thus bounded sublattices, of M and M/θ , respectively, then $p_{\theta} \mid_{\mathcal{B}(M)} : \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a bounded lattice morphism, hence, if $\mathcal{B}(M)$ and $\mathcal{B}(M/\theta)$ are Boolean sublattices of M and M/θ , respectively, then $p_{\theta} \mid_{\mathcal{B}(M)} : \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean morphism, which is surjective iff $\mathcal{B}(M)/\theta = \mathcal{B}(M/\theta)$ and is injective iff $0/\theta \cap \mathcal{B}(M) = (p_{\theta} \mid_{\mathcal{B}(M)})^{-1}(\{0/\theta\}) = \{0\}$ iff $1/\theta \cap \mathcal{B}(M) = (p_{\theta} \mid_{\mathcal{B}(M)})^{-1}(\{1/\theta\}) = \{1\}$.

Therefore, if $\mathcal{B}(M)$ is a Boolean sublattice of M and $\mathcal{B}(M)/\theta = \mathcal{B}(M/\theta)$, then $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ and $p_{\theta}|_{\mathcal{B}(M)} \colon \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a surjective Boolean morphism.

Remark 4.10. If the map $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is injective and $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ , then $\mathcal{B}(M)$ is a Boolean sublattice of M and $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is an injective Boolean morphism.

Indeed, if this restriction of the bounded lattice morphism $p_{\theta} : M \to M/\theta$ is injective and its codomain $\mathcal{B}(M/\theta)$ is a distributive sublattice of M/θ , then its domain $\mathcal{B}(M)$ is a distributive and thus a Boolean sublattice of M and hence $p_{\theta} \mid_{\mathcal{B}(M)} : \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean embedding. **Lemma 4.11.** Let M be a bounded lattice, $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and $e \in M$.

- (i) If $(e]_M \in Ann(M)$ or $e \in \mathcal{B}(M)$ is the unique complement of an $f \in \mathcal{B}(M)$, then $e = \max(e/\theta)$.
- (ii) If $\{(g]_M \mid g \in \mathcal{B}(M)\} \subseteq \mathcal{A}nn(M)$ or M is uniquely complemented, in particular if $\mathcal{B}(M)$ is a Boolean sublattice of M, then $g = \max(g/\theta)$ for all $g \in \mathcal{B}(M)$.
- (iii) If $e = \max(e/\theta)$, in particular if $(e]_M \in Ann(M)$ or $e \in \mathcal{B}(M)$ is the unique complement of an $f \in \mathcal{B}(M)$, then, for all $U \subseteq M$: $(e]_M/\theta = \operatorname{Ann}_M(U)/\theta$ iff $(e]_M = \operatorname{Ann}_M(U)$.
- (iv) If M/θ is distributive, $e \in \mathcal{B}(M)$ and $f \in \mathcal{B}(M)$ is a complement of e, then: $(e]_M \in \mathcal{A}nn(M)$ iff $(e]_M \in \text{PAnn}(M)$ iff $(e]_M = \text{Ann}_M(f)$ iff $e = \max(e/\theta)$.
- (v) Assume that $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ . If $g = \max(g/\theta)$ for all $g \in \mathcal{B}(M)$, in particular if $\{(g]_M \mid g \in \mathcal{B}(M)\} \subseteq \mathcal{A}nn(M)$ or Mis uniquely complemented, then $\mathcal{B}(M)$ is a Boolean sublattice of M and $p_{\theta} \mid_{\mathcal{B}(M)} : \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean embedding.
- (vi) If M/θ is distributive and M is uniquely complemented, then $\mathcal{B}(M)$ is a Boolean sublattice of M, $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean embedding and, for all $g \in \mathcal{B}(M)$, $g = \max(g/\theta)$ and $(g]_M = \operatorname{Ann}_M(\neg g)$.

Proof. (i) If $(e]_M = \operatorname{Ann}_M(U)$ for some $U \subseteq M$, then, by Lemma 4.2, (i), $e \in e/\theta \subseteq \operatorname{Ann}_M(U) = (e]_M$, thus $e = \max(e/\theta)$.

Now assume that $e \in \mathcal{B}(M)$ is the unique complement of an $f \in \mathcal{B}(M)$, and assume by absurdum that there exists an $x \in e/\theta$ such that $x \nleq e$. Then, if we denote by $a = x \lor e$, it follows that a > e and thus $a \lor f = 1$. But $a/\theta = x/\theta \lor e/\theta = e/\theta \lor e/\theta = e/\theta$, so $(a \land f)/\theta = a/\theta \land f/\theta = e/\theta \land f/\theta =$ $(e \land f)/\theta = 0/\theta = \{0\}$, hence $a \land \neg e = 0$. Since $a \neq e$, we have a contradiction to the uniqueness of the complement of f in M. Therefore $e = \max(e/\theta)$.

(ii) By (i).

(iii) Trivially, if $(e]_M = \operatorname{Ann}_M(U)$, then $(e]_M/\theta = \operatorname{Ann}_M(U)/\theta$.

If $e = \max(e/\theta)$ and $(e]_M/\theta = \operatorname{Ann}_M(U)/\theta$, that is $(e]_M/\theta \subseteq \operatorname{Ann}_M(U)/\theta$ and $\operatorname{Ann}_M(U)/\theta \subseteq (e]_M/\theta$, then $(e]_M \subseteq \operatorname{Ann}_M(U)$ by Lemma 4.2, (iii), and $\operatorname{Ann}_M(U) \subseteq (e]_M$ by Remark 4.5, so the converse of the implication above also holds.

We get the particular cases from (i).

(iv) By (i), if $(e|_M \in Ann(M)$, then $e = \max(e/\theta)$.

Of course, $(e]_M = \operatorname{Ann}_M(f)$ implies $(e]_M \in \operatorname{PAnn}(M)$, which in turn implies $(e]_M \in \operatorname{Ann}(M)$.

Now assume that $e = \max(e/\theta)$. Since e is a complement of f in M and M/θ is distributive, it follows that e/θ is the unique complement of f/θ in M/θ .

We have $e \in \operatorname{Ann}_M(f)$, thus $(e]_M \subseteq \operatorname{Ann}_M(f)$. Assume by absurdum that there exists an $x \in \operatorname{Ann}_M(f)$ such that $x \nleq e$, and let $a = x \lor e$. Then a > e, thus $a \lor f = 1$, so $a/\theta \lor f/\theta = 1/\theta$, and $a/\theta \neq e/\theta$ since $e = \max(e/\theta)$. Since M/θ is distributive, we have: $a/\theta \land f/\theta = (x/\theta \lor e/\theta) \land f/\theta = (x/\theta \land f/\theta) \lor (e/\theta \land f/\theta) = (x \land f)/\theta \lor (e \land f)/\theta = 0/\theta \lor 0/\theta = 0/\theta$, which gives us a contradiction to the uniqueness of the complement of f/θ . Therefore $(e]_M = \operatorname{Ann}_M(f)$.

(v) By Remarks 4.8 and 4.10, with the particular cases given by (ii).

(vi) By (ii), (v) and (iv).

Proposition 4.12. Let M be a bounded lattice. Then:

- (i) $\{(e]_M \mid e \in \mathcal{B}(M)\} \subseteq \mathcal{B}(\mathrm{Id}(M));$
- (ii) let κ be a nonzero cardinality; if, for all e, f ∈ B(M) such that e is a complement of f in M, we have (e]_M = Ann_M(f), in particular if, for some θ ∈ Con(M) such that 0/θ = {0} and M/θ is distributive, we have e = max(e/θ) for all e ∈ B(M), or, equivalently, {(e]_M | e ∈ B(M)} ⊆ Ann(M), in particular if M is uniquely complemented and has a θ ∈ Con(M) with 0/θ = {0} such that M/θ is distributive, then:
 - $\mathcal{B}(\mathrm{Id}(M)) = \{(e|_M \mid e \in \mathcal{B}(M)\} \subseteq \mathrm{PAnn}(M) \subseteq \mathcal{A}nn_{\kappa}(M) \subseteq \mathcal{A}nn(M);$
 - M is a Stone lattice iff $PAnn(M) = \mathcal{B}(Id(M));$
 - M is a κ -Stone lattice iff $Ann_{\kappa}(M) = \mathcal{B}(\mathrm{Id}(M));$
 - M is a strongly Stone lattice iff $Ann(M) = \mathcal{B}(Id(M))$.

Proof. (i) For all $e \in \mathcal{B}(M)$, if f is a complement of e in M, then $(e]_M \vee (f]_M = (e \vee f]_M = (1]_M = M$ and $(e]_M \cap (f]_M = (e \wedge f]_M = (0]_M = \{0\}$, so $(e]_M \in \mathcal{B}(\mathrm{Id}(M))$.

(ii) Let $I \in \mathcal{B}(\mathrm{Id}(M))$, so that $I \cap J = \{0\}$ and $I \vee J = M = (1]_M$ for some $J \in \mathrm{Id}(M)$, hence $e \vee f = 1$ for some $e \in I$ and $f \in J$, thus $e \wedge f \in I \cap J = \{0\}$, so $e \wedge f = 0$, hence $e, f \in \mathcal{B}(M)$ and f is a complement of e, thus $(e]_M = \mathrm{Ann}_M(f)$ by the hypothesis. Since $e \in I$, we have $(e]_M \subseteq I$. For all $x \in I$ and all $y \in J$, we have $x \wedge y \in I \cap J = \{0\}$, so $x \wedge y = 0$, hence $I \subseteq \mathrm{Ann}_M(J) \subseteq \mathrm{Ann}_M(f) = (e]_M$. Therefore $I = (e]_M = \mathrm{Ann}_M(f) \in \mathrm{PAnn}(M)$. Hence the converse of the inclusion in (i), thus $\mathcal{B}(\mathrm{Id}(M)) = \{(g]_M \mid g \in \mathcal{B}(M)\} \subseteq \mathrm{PAnn}(M) \subseteq \mathcal{Ann}(M)$, hence the last two statements.

We get the particular cases from Lemma 4.11, (ii) and (iv).

Lemma 4.13. Let M be a bounded lattice and $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and $1/\theta = \{1\}$.

- (i) If $\mathcal{B}(M)$ is a Boolean sublattice of M, then $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ and $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean isomorphism.
- (ii) If $e = \max(e/\theta)$ for all $e \in \mathcal{B}(M)$, in particular if $\{(e]_M \mid e \in \mathcal{B}(M)\} \subseteq \mathcal{A}nn(M)$ or M is uniquely complemented, then: $\mathcal{B}(M)$ is a Boolean sublat-

tice of M iff $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ , and, if so, then $p_{\theta}|_{\mathcal{B}(M)}: \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a Boolean isomorphism.

Proof. (i) By Remark 4.7, $\mathcal{B}(M)/\theta = \mathcal{B}(M/\theta)$, thus, according to Remark 4.9, $\mathcal{B}(M/\theta)$ is a Boolean sublattice of M/θ and $p_{\theta} \mid_{\mathcal{B}(M)} : \mathcal{B}(M) \to \mathcal{B}(M/\theta)$ is a surjective Boolean morphism. Since $0/\theta \cap \mathcal{B}(M) = \{0\} \cap \mathcal{B}(M) = \{0\}$, Remark 4.9 ensures us that this Boolean morphism is also injective, so it is a Boolean isomorphism.

(ii) By (i) and Remarks 4.10 and 4.8, with Lemma 4.11, (ii), for the particular cases.

Proposition 4.14. If M is a bounded lattice and $\theta \in Con(M)$ such that $0/\theta = \{0\}$, then, for any cardinality κ :

- (i) $(1)_{\kappa,M}$ implies $(1)_{\kappa,M/\theta}$, thus: if M is Stone, respectively κ -Stone, respectively strongly Stone, then M/θ is Stone, respectively κ -Stone, respectively strongly Stone;
- (ii) if 1/θ = {1}, then: if e = max(e/θ) for all e ∈ B(M), in particular if (e]_M ∈ Ann(M) for all e ∈ B(M), in particular if M is uniquely complemented, in particular if B(M) is a Boolean sublattice of M, then (1)_{κ,M} is equivalent to (1)_{κ,M/θ}, thus: M is Stone, respectively κ-Stone, respectively strongly Stone iff M/θ is Stone, respectively κ-Stone, respectively strongly Stone.

Proof. (i) Let $V \subseteq M/\theta$ such that $|V| \leq \kappa$, so that $V = U/\theta$ for some $U \subseteq M$ with $|U| = |V| \leq \kappa$. If $(1)_{\kappa,M}$ is fulfilled, then there exists an $e \in \mathcal{B}(M)$ such that $\operatorname{Ann}_M(U) = (e]_M$, so that $e/\theta \in \mathcal{B}(M)/\theta \subseteq \mathcal{B}(M/\theta)$, and $(e/\theta]_{M/\theta} = (e]_M/\theta =$ $\operatorname{Ann}_M(U)/\theta = \operatorname{Ann}_{M/\theta}(U/\theta) = \operatorname{Ann}_{M/\theta}(V)$ by Lemma 4.2, (ii), hence $(1)_{\kappa,M/\theta}$ is fulfilled.

(ii) By (i), we have the direct implication. For the converse, let $U \subseteq M$ such that $|U| \leq \kappa$, so that $|U/\theta| \leq |U| \leq \kappa$, and thus, if $(1)_{\kappa,M/\theta}$ is fulfilled, then, for some $e \in M$ such that $e/\theta \in \mathcal{B}(M/\theta)$, so that $e \in \mathcal{B}(M)$ by Remark 4.7, and thus $e = \max(e/\theta)$ by the hypothesis, we have $(e]_M/\theta = (e/\theta)_{M/\theta} = \operatorname{Ann}_{M/\theta}(U/\theta) = \operatorname{Ann}_M(U)/\theta$, hence $(e]_M = \operatorname{Ann}_M(U)$ by Lemma 4.11, (iii). For the particular cases, see Lemma 4.11, (i) and (ii).

Proposition 4.15. If M is a bounded lattice and $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and $1/\theta = \{1\}$, then, for any cardinality κ :

- (i) $(2)_{\kappa,M}$ implies $(2)_{\kappa,M/\theta}$;
- (ii) if $e = \max(e/\theta)$ for all $e \in \mathcal{B}(M)$, in particular if $(e]_M \in \mathcal{A}nn(M)$ for all $e \in \mathcal{B}(M)$, in particular if M is uniquely complemented, in particular if $\mathcal{B}(M)$ is a Boolean sublattice of M, then $(2)_{\kappa,M}$ is equivalent to $(2)_{\kappa,M/\theta}$.

64

Proof. By Proposition 4.14 and Lemma 4.13, (ii), with Lemma 4.11, (i) and (ii), for the particular cases in (ii).

Proposition 4.16. If M is a bounded lattice and $\theta \in \text{Con}(M)$ such that $0/\theta = \{0\}$, then, for any cardinality κ , $(3)_{\kappa,M}$ is equivalent to $(3)_{\kappa,M/\theta}$.

Proof. By Lemma 4.2, (ii), Lemma 4.3, (x), and the fact that the following diagram is commutative:

$$\begin{array}{c} M \xrightarrow{x \mapsto \operatorname{Ann}_{M}(\operatorname{Ann}_{M}(x))} \operatorname{P2Ann}(M) \\ \downarrow x \mapsto x/\theta \qquad \qquad \downarrow \forall x \in M, \operatorname{Ann}_{M}(\operatorname{Ann}_{M}(x)) \mapsto \\ M/\theta \xrightarrow{x/\theta \mapsto \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(x/\theta))} \operatorname{P2Ann}(M/\theta) \operatorname{Ann}_{M}(\operatorname{Ann}_{M}(x))/\theta = \\ \operatorname{P2Ann}(M/\theta) \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(x/\theta)) \end{array}$$

Proposition 4.17. If M is a bounded lattice and $\theta \in \text{Con}(M)$ such that $0/\theta = \{0\}$, then, for any cardinality κ , $(4)_{\kappa,M}$ is equivalent to $(4)_{\kappa,M/\theta}$.

Proof. By Lemma 4.2, (ii), Lemma 4.3, (x), and the surjectivity of the map $x \mapsto x/\theta$ from M to M/θ .

Proposition 4.18. For any bounded lattice M, any $\theta \in Con(M)$ and any cardinality κ :

- (i) $(5)_{\kappa,M}$ implies $(5)_{\kappa,M/\theta}$;
- (ii) if $0/\theta = \{0\}$ and $1/\theta = \{1\}$, then $(5)_{\kappa,M}$ is equivalent to $(5)_{\kappa,M/\theta}$.

Proof. (i) If $(5)_{\kappa,M}$ if fulfilled and $V \subseteq M/\theta$ with $|V| \leq \kappa$, then $V = U/\theta$ for some $U \subseteq M$ with $|U| \leq \kappa$, so that $(\operatorname{Ann}_M(U) \cup \operatorname{Ann}_M(\operatorname{Ann}_M(U))]_M = M$, thus $1 = a_1 \lor \cdots \lor a_n \lor b_1 \lor \cdots \lor b_k$ for some $n, k \in \mathbb{N}^*$, $a_1, \ldots, a_n \in \operatorname{Ann}_M(U)$ and $b_1, \ldots, b_k \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$. Then $a_1/\theta, \ldots, a_n/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta) =$ $\operatorname{Ann}_{M/\theta}(V)$ and $b_1/\theta, \ldots, b_k/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta)) = \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(V))$, therefore $1/\theta = a_1/\theta \lor \cdots \lor a_n/\theta \lor b_1/\theta \lor \cdots \lor b_k/\theta \in (\operatorname{Ann}_{M/\theta}(V) \cup \operatorname{Ann}_{M/\theta}(V))$ $(\operatorname{Ann}_{M/\theta}(V))]_{M/\theta}$, hence $(\operatorname{Ann}_{M/\theta}(V) \cup \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(V))]_{M/\theta} = M/\theta$.

(ii) Assume that $0/\theta = \{0\}$ and $1/\theta = \{1\}$. If $(5)_{\kappa,M/\theta}$ is fulfilled and $U \subseteq M$ with $|U| \leq \kappa$, then $|U/\theta| \leq |U| \leq \kappa$, so that $(\operatorname{Ann}_{M/\theta}(U/\theta) \cup \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta))]_{M/\theta} = M/\theta$, thus $1/\theta = a_1/\theta \vee \cdots \vee a_n/\theta \vee b_1/\theta \vee \cdots \vee b_k/\theta$ for some $n, k \in \mathbb{N}^*$ and $a_1, \ldots, a_n, b_1, \ldots, b_k \in M$ such that $a_1/\theta, \ldots, a_n/\theta \in \operatorname{Ann}_{M/\theta}(U/\theta)$ and $b_1/\theta, \ldots, b_k/\theta \in \operatorname{Ann}_{M/\theta}(\operatorname{Ann}_{M/\theta}(U/\theta))$. But then $a_1, \ldots, a_n \in \operatorname{Ann}_M(U)$ and $b_1, \ldots, b_k \in \operatorname{Ann}_M(\operatorname{Ann}_M(U))$ by Lemma 4.2, (i), and $\{1\} = 1/\theta = (a_1 \vee \cdots \vee a_n \vee b_1 \vee \cdots \vee b_k)/\theta$, thus $1 = a_1 \vee \cdots \vee a_n \vee b_1 \vee \cdots \vee b_k \in (\operatorname{Ann}_M(U) \cup \operatorname{Ann}_M(\operatorname{Ann}_M(U))]_M$ hence $(\operatorname{Ann}_M(U) \cup \operatorname{Ann}_M(\operatorname{Ann}_M(U))]_M = M$. We have the converse from (i), so the equivalence holds. **Theorem 4.19.** Let M be a bounded lattice, $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and $1/\theta = \{1\}$ and m be a nonzero cardinality. If $e = \max(e/\theta)$ for all $e \in \mathcal{B}(M)$, in particular if $(e]_M \in Ann(M)$ for all $e \in \mathcal{B}(M)$, in particular if M is uniquely complemented, in particular if $\mathcal{B}(M)$ is a Boolean sublattice of M, then:

- (i) if M/θ is distributive, then, for any nonzero cardinality κ, conditions (1)_{κ,M}, (2)_{κ,M}, (3)_{κ,M}, (4)_{κ,M} and (5)_{κ,M} are equivalent;
- (ii) if M/θ is distributive and Ann_m(M) = PAnn(M), in particular if M/θ is closed w.r.t. the joins of all families of elements of cardinality at most m and has the meet distributive w.r.t. the joins of families of cardinalities at most m, then, for any h, i ∈ 1,5 and any nonzero cardinality κ ≤ m, conditions (h)_{κ,M} and (i)_{<∞,M} are equivalent;
- (iii) if M/θ is distributive and Ann(M) = PAnn(M), in particular if M/θ is a frame, then, for any $h, i, j \in \overline{1,5}$ and any nonzero cardinality κ , conditions $(iv)_M, (h)_{\kappa,M}, (i)_{<\infty,M}$ and $(j)_M$ are equivalent.

Proof. (i) By Theorem 3.3, (i), Proposition 4.14, (ii), Proposition 4.15, (ii), Propositions 4.16 and 4.17 and Proposition 4.18, (ii).

(ii) By Theorem 3.3, (iii), Proposition 4.14, (ii), Proposition 4.15, (ii), Propositions 4.16 and 4.17 and Proposition 4.18, (ii).

(iii) By Theorem 3.3, (iv), Proposition 4.14, (ii), Proposition 4.15, (ii), Propositions 4.16 and 4.17 and Proposition 4.18, (ii).

Let us also note:

Proposition 4.20. Let M be a bounded lattice, $\theta \in Con(M)$ such that $0/\theta = \{0\}$ and m be a nonzero cardinality.

- If M/θ is distributive, then, for any nonzero cardinality κ, condition (3)_{κ,M} is equivalent to (4)_{κ,M}.
- If $1/\theta = \{1\}$ and M/θ is distributive, then, for any nonzero cardinality κ , conditions $(3)_{\kappa,M}$, $(4)_{\kappa,M}$ and $(5)_{\kappa,M}$ are equivalent.
- If M/θ is distributive and $Ann_m(M/\theta) = PAnn(M/\theta)$, in particular if M/θ is closed w.r.t. the joins of all families of elements of cardinality at most m and has the meet distributive w.r.t. such joins, then, for any nonzero cardinalities $\kappa \leq m$ and $\mu \leq m$, condition (3)_{κ,M} is equivalent to (4)_{μ,M}.
- If $1/\theta = \{1\}$, then: if M/θ is distributive and $Ann_m(M/\theta) = PAnn(M/\theta)$, in particular if M/θ is closed w.r.t. the joins of all families of elements of cardinality at most κ and has the meet distributive w.r.t. the joins of families of cardinalities at most κ , then, for any nonzero cardinalities $\kappa \leq m$, $\lambda \leq m$ and $\mu \leq m$, conditions $(3)_{\kappa,M}$, $(4)_{\lambda,M}$ and $(5)_{\mu,M}$ are equivalent.

- If M/θ is distributive and Ann(M/θ) = PAnn(M/θ), in particular if M/θ is a frame, then, for any nonzero cardinalities κ and μ, conditions (3)_{κ,M}, (4)_{μ,M} and (iv)_M are equivalent.
- If $1/\theta = \{1\}$, then: if M/θ is distributive and $Ann(M/\theta) = PAnn(M/\theta)$, in particular if M/θ is a frame, then, for any nonzero cardinalities κ , λ and μ , conditions $(3)_{\kappa,M}$, $(4)_{\lambda,M}$, $(iv)_M$ and $(5)_{\mu,M}$ are equivalent.

Proof. By Theorem 3.3 and Propositions 4.16, 4.17 and 4.18.

Recall from Proposition 4.4 that, if $0/\theta = \{0\}$ and M/θ is distributive, then $\mathcal{A}nn(M) \subseteq \mathrm{Id}(M)$, while, if $0/\theta = \{0\}$ and M/θ is a frame, then $\mathcal{A}nn(M) = \mathrm{PAnn}(M) \subseteq \mathrm{Id}(M)$.

If we eliminate the nontrivial implications from Theorem 4.19, (ii) and (iii), along with those that immediately follow from Proposition 4.4, (ii) and (iii), and Remark 3.2, then we obtain the following.

Corollary 4.21. Let M be a bounded lattice and $\theta \in \text{Con}(M)$ such that $0/\theta = \{0\}$ and $1/\theta = \{1\}$ and m be a nonzero cardinality. If $e = \max(e/\theta)$ for all $e \in \mathcal{B}(M)$, in particular if $(e]_M \in Ann(M)$ for all $e \in \mathcal{B}(M)$, in particular if M is uniquely complemented, in particular if $\mathcal{B}(M)$ is a Boolean sublattice of M, then:

- (i) if M/θ is distributive and Ann_m(M) = PAnn(M), in particular if M/θ is closed w.r.t. the joins of all families of elements of cardinality at most m and has the meet distributive w.r.t. such joins, then the following are equivalent:
 - M is Stone;
 - M is m-Stone and $\mathcal{B}(M)$ is an m-complete Boolean sublattice of M;
 - P2Ann(M) is a Boolean sublattice of Id(M) such that a → Ann_M(Ann_M
 (a)) is a lattice morphism from M to P2Ann(M);
 - P2Ann(M) is an m-complete Boolean sublattice of Id(M) such that $a \mapsto$ Ann_M(Ann_M(a)) is a lattice morphism from M to P2Ann(M);
 - for all $a, b \in M$, $\operatorname{Ann}_M(a \wedge b) = \operatorname{Ann}_M(a) \vee \operatorname{Ann}_M(b)$ and $\operatorname{P2Ann}(M) \subseteq \operatorname{PAnn}(M)$;
 - for all $a \in M$, $\operatorname{Ann}_M(a) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(a)) = M$;
- (ii) if M/θ is distributive and Ann(M) = PAnn(M), in particular if M/θ is a frame, then the following are equivalent:
 - M is Stone;
 - M is strongly Stone and $\mathcal{B}(M)$ is a complete Boolean sublattice of M;
 - P2Ann(M) is a Boolean sublattice of Id(M) such that a → Ann_M(Ann_M
 (a)) is a lattice morphism from M to P2Ann(M);

- P2Ann(M) is a complete Boolean sublattice of Id(M) such that $a \mapsto Ann_M(Ann_M(a))$ is a lattice morphism from M to P2Ann(M);
- for all $a, b \in M$, $\operatorname{Ann}_M(a \wedge b) = \operatorname{Ann}_M(a) \vee \operatorname{Ann}_M(b)$;
- for all $a \in M$, $\operatorname{Ann}_M(a) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(a)) = M$.

5. A CERTAIN CONGRUENCE OF A COMMUTATOR LATTICE

Throughout this section, unless mentioned otherwise, $(L, \lor, \land, [\cdot, \cdot], 0, 1)$ will be a commutator lattice. Some of the results that follow in this paper generalize properties obtained in [10] for the particular case of congruence lattices endowed with the term-condition commutator, under the condition that this commutator operation is commutative and distributive in both arguments (of course, one suffices, by the commutativity) w.r.t. arbitrary joins, in particular for congruence lattices of members of congruence-modular varieties, endowed with the modular commutator.

Remark 5.1. Since $[x, y] \leq x \wedge y$ for all $x, y \in L$, we clearly have $\text{Spec}_L \subseteq \text{Mi}(L) \setminus \{1\}$. If L has finite length, then $\text{Mi}(L) \setminus \{1\} = \text{Smi}(L)$, so that $\text{Spec}_L \subseteq \text{Smi}(L)$.

Lemma 5.2. Let $(L, [\cdot, \cdot])$ be a commutator lattice. Then:

- (i) $\{x \in \operatorname{Smi}(L) \mid [x^+, x^+] \nleq x\} \subseteq \operatorname{Spec}_L; \text{ if } L \text{ has finite length, then } \operatorname{Spec}_L = \{x \in \operatorname{Smi}(L) \mid [x^+, x^+] \nleq x\};$
- (ii) Spec_L = { $x \in Mi(L) \setminus \{1\} \mid (\forall a \in L) ([a, a] \le x \Rightarrow a \le x)$ };
- (iii) if [1,1] = 1, then $\operatorname{Max}_L \subseteq \operatorname{Spec}_L$;
- (iv) if $1 \in Cp(L)$, then, for each $x \in L \setminus \{1\}$, there exists a $p \in Max_L$ such that $x \leq p$;
- (v) if $1 \in Cp(L)$ and [1,1] = 1, then, for each $x \in L \setminus \{1\}$, there exists a $p \in Spec_L$ such that $x \leq p$.

Proof. (i) Take $x \in \text{Smi}(L) \subseteq L \setminus \{1\}$ with $[x^+, x^+] \nleq x$, so that $[x^+, x^+] = x^+$ since $[x^+, x^+] \le x^+$, and let $a, b \in L$ such that $[a, b] \le x$. Assume by absurdum that $a \nleq x$ and $b \nleq x$, which means that $a \lor x \ne x$ and $b \lor x \ne x$, hence $a \lor x > x$ and $b \lor x > x$, so that $a \lor x \ge x^+$ and $b \lor x \ge x^+$. Then $x < x^+ = [x^+, x^+] \le$ $[a \lor x, b \lor x] = [a, b] \lor [a, x] \lor [x, b] \lor [x, x] \le x$, so we have a contradiction. Thus $x \in \text{Spec}_L$.

By the definition of Spec_L , if $x \in \operatorname{Smi}(L)$ is such that $[x^+, x^+] \leq x$, then $x \notin \operatorname{Spec}_L$. If L has finite length, then $\operatorname{Spec}_L \subseteq \operatorname{Smi}(L)$, hence the equality for this case.

(ii) By the proof of [1, Proposition 1.2], which we reproduce here for the sake of completeness. The left-to-right inclusion is clear. Now let $x \in Mi(L) \setminus \{1\}$

such that, for all $a \in L$, $[a, a] \leq x$ implies $a \leq x$. Let $a, b \in L$ such that $[a, b] \leq x$ and assume by absurdum that $a \nleq x$ and $b \nleq x$, so that $a \lor x > x$ and $b \lor x > x$ and thus $(a \lor x) \land (b \lor x) > x$ since x is meet-irreducible. Then $[(a \lor x) \land (b \lor x), (a \lor x) \land (b \lor x)] \leq [a \lor x, b \lor x] = [a, b] \lor [a, x] \lor [x, b] \lor [x, x] \leq x$, hence $(a \lor x) \land (b \lor x) \leq x$ by the choice of x, and we have a contradiction. Hence $x \in \operatorname{Spec}_L$.

(iii) Clearly, each $x \in Max_L$ is strictly meet-irreducible, with $x^+ = 1$. Now apply (i).

(iv) Assume that $1 \in \operatorname{Cp}(L)$ and let $x \in L \setminus \{1\}$. We prove that the set $([x)_L \setminus \{1\}, \leq)$ is inductively ordered. Let $C \subseteq [x)_L \setminus \{1\}$ such that (C, \leq) is a chain, and let $t = \bigvee C$. We can not have t = 1, because then, since $1 \in \operatorname{Cp}(L)$, there would exist an $n \in \mathbb{N}^*$ and elements $c_1, \ldots, c_n \in C$ such that $1 = \bigvee_{i=1}^n c_i = \max\{c_1, \ldots, c_n\} \in \{c_1, \ldots, c_n\} \subseteq C \subseteq L \setminus \{1\}$, which gives us a contradiction. Hence $t \in L \setminus \{1\}$. But $x \leq t$, thus $t \in [x)_L \setminus \{1\}$, so indeed $([x)_L \setminus \{1\}, \leq)$ is inductively ordered, therefore it has maximal elements by Zorn's Lemma, and clearly its maximal elements are also maximal elements of $L \setminus \{1\}$, that is they belong to Max_L , and they are greater than x.

(v) By (iii) and (iv).

Remark 5.3. By Lemma 5.2, (ii), if $[\cdot, \cdot] = \wedge$, then $\text{Spec}_L = \text{Mi}(L) \setminus \{1\}$.

Remark 5.4. Let $x, y \in L$, $M = \{a \in L \mid [a, x] \leq y\}$ and $N = \{b \in L \mid [b, x] = y\}$. Then $\bigvee \emptyset = 0 \in M$, in particular M is nonempty, and, for any nonempty family $(a_i)_{i \in I} \subseteq M$, we have $[a_i, x] \leq y$ for all $i \in I$ and thus $[\bigvee_{i \in I} a_i, x] = \bigvee_{i \in I} [a_i, x] \leq y$, hence $\bigvee_{i \in I} a_i \in M$. Therefore $\bigvee_{i \in I} a_i \in M$ for any family $(a_i)_{i \in I} \subseteq M$, hence the set M has a maximum, namely $\max(M) = \bigvee M$.

If N is nonempty, then, for any nonempty family $(b_j)_{j\in J} \subseteq N$, we have $[b_j, x] = y$ for all $j \in J$ and thus $[\bigvee_{j\in J} b_j, x] = \bigvee_{j\in J} [b_j, x] = y$, hence $\bigvee_{j\in I} b_j \in N$, and thus N has a maximum, namely $\max(N) = \bigvee N$.

Remark 5.5. The radical elements of L are exactly the meets of the families of prime elements of L, hence $\operatorname{Spec}_L \subseteq R(L)$ and the map $x \mapsto \rho(x)$ is a closure operator on L with associated closure system $R(L) = \{\rho(x) \mid x \in L\} = \{x \in L \mid \rho(x) = x\}$, so that $1 \in R(L)$ since $\rho(1) = \bigwedge \emptyset = 1$ and the following hold for all $a, b \in L, p \in \operatorname{Spec}_L$ and $r \in R(L)$:

- $a \leq \rho(a);$
- $a \leq \rho(b)$ iff $\rho(a) \leq \rho(b)$, so that $a \leq r$ iff $\rho(a) \leq r$; in particular, $a \leq p$ iff $\rho(a) \leq p$, thus $V(a) = V(\rho(a))$;
- $a \leq b$ implies $V(b) \subseteq V(a)$, which implies $\rho(a) \leq \rho(b)$, which in turn implies $V(b) = V(\rho(b)) \subseteq V(\rho(a)) = V(a)$, hence: $\rho(a) \leq \rho(b)$ iff $V(b) \subseteq V(a)$, and thus: $\rho(a) = \rho(b)$ iff V(a) = V(b).

Throughout the rest of this section, unless mentioned otherwise, \equiv will be the following equivalence on the set L: $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$. By Remark 5.5, $\equiv = \{(a, b) \in L^2 \mid V(a) = V(b)\}$.

Remark 5.6. $1 \in 1/\equiv$, thus the equality $1/\equiv \{1\}$ means that, for all $a \in L$: a = 1 iff $\rho(a) = \rho(1) = 1$ iff $V(a) = V(1) = \emptyset$, which in turn is equivalent to $\{a \in L \mid \rho(a) = 1\} = \{1\}$ and to $V(a) \neq \emptyset$ for each $a \in L \setminus \{1\}$.

Lemma 5.7. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $1 \in Cp(L)$ and [1, 1] = 1 and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$, then $1/\equiv = \{1\}$.

Proof. By Lemma 5.2, (v), and Remark 5.6.

Remark 5.8. We use Remark 5.5 in what follows. Let $a, b \in L$ and $(a_i)_{i \in I} \subseteq L$. Then $V(a) = V(\rho(a)), V(a \land b) = V([a, b]) = V(a) \cup V(b) = V(\rho(a) \land \rho(b)) = V([\rho(a), \rho(b)]) = V(\rho(a)) \cup V(\rho(b))$ and $V(\bigvee_{i \in I} a_i) = \bigcap_{i \in I} V(a_i) = V(\bigvee_{i \in I} \rho(a_i))$, so that $\rho(a \land b) = \rho([a, b]) = \rho(a) \land \rho(b) = \rho(\rho(a) \land \rho(b)) = \rho([\rho(a), \rho(b)])$ and $\rho(\bigvee_{i \in I} a_i) = \rho(\bigvee_{i \in I} \rho(a_i))$, otherwise written: $a \land b \equiv [a, b] \equiv \rho(a) \land \rho(b) \equiv [\rho(a), \rho(b)]$ and $\bigvee_{i \in I} a_i \equiv \bigvee_{i \in I} \rho(a_i)$.

Indeed, $\rho(a) = \rho(\rho(a))$, hence the first equality. Since $[a, b] \leq a \wedge b \leq a, b$, we have $V(a) \cup V(b) \subseteq V(a \wedge b) \subseteq V([a, b])$. By the definition of prime elements, $V([a, b]) \subseteq V(a) \cup V(b)$. Thus, by also using the first equality, we get the second set of equalities. Finally, $V(\bigvee_{i \in I} a_i) = [\bigvee_{i \in I} a_i)_L \cap \operatorname{Spec}_L = \bigcap_{i \in I} [a_i)_L \cap \operatorname{Spec}_L = \bigcap_{i \in I} ([a_i)_L \cap \operatorname{Spec}_L) = \bigcap_{i \in I} V(a_i) = \bigcap_{i \in I} V(\rho(a_i)) = V(\bigvee_{i \in I} \rho(a_i)).$

For any $x \in L$ and any $n \in \mathbb{N}^*$, we shall denote: $x^1 = x$ and $x^{n+1} = [x, x^n]$.

Proposition 5.9. Let $(L, [\cdot, \cdot])$ be a commutator lattice and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$. Then:

- (i) \equiv is a lattice congruence of L which preserves arbitrary joins and the commutator operation $[\cdot, \cdot]$ and satisfies $[a, b] \equiv a \wedge b$ for all $a, b \in L$, $R(L) = \{\max(x/\equiv) \mid x \in L\} = \{x \in L \mid x = \max(x/\equiv)\}, 0/\equiv = (\rho(0)]_L,$ and, for all $x \in L, \rho(x) = \max(x/\equiv) = \max(\rho(x)/\equiv) = \min([x)_L \cap R(L));$
- (ii) $\equiv Cg_L(\{(x, \rho(x)) \mid x \in L\}) \supseteq Cg_L(\{(x \land y, [x, y]) \mid x, y \in L\}) \supseteq Cg_L(\{(x, [x, x]) \mid x \in L\});$
- (iii) for all $x \in L$ such that x/\equiv has a minimum and all $a \in [\min(x/\equiv))_L \supseteq x/\equiv$, $[a, \min(x/\equiv)] = \min(x/\equiv);$
- (iv) if, for each $x \in L$, there exists an $n_x \in \mathbb{N}^*$ such that $\rho(x)^{n_x} = \min(x/\equiv)$, then: $\equiv -Cg_L(\{(x, \rho(x)) \mid x \in L\}) = Cg_L(\{(x \land y, [x, y]) \mid x, y \in L\})$ and, for all $a \in x/\equiv$ and all $n \in \mathbb{N}$ with $n \ge n_x$, $a^n = \min(x/\equiv)$.

Proof. We denote by $\sigma = Cg_L(\{(x, \rho(x)) \mid x \in L\}), \sim = Cg_L(\{(x \land y, [x, y]) \mid x, y \in L\})$ and $\approx = Cg_L(\{(x, [x, x]) \mid x \in L\})$. We will repeatedly use Remark 5.8. (i) For all $a, b, x, y \in L$ such that $a \equiv x$ and $b \equiv y$, so that $\rho(a) = \rho(x)$ and $\rho(b) = \rho(y)$, we have $[a, b] \equiv a \land b \equiv \rho(a) \land \rho(b) = \rho(x) \land \rho(y) \equiv x \land y \equiv [x, y]$. For all $(a_i)_{i \in I}, (b_i)_{i \in I} \subseteq L$ such that, for all $i \in I$, $a_i \equiv b_i$, so that $\rho(a_i) = \rho(b_i)$, we have $\bigvee_{i \in I} \rho(a_i) = \bigvee_{i \in I} \rho(b_i) \equiv \bigvee_{i \in I} b_i$.

For each $x \in L$, we have $\rho(\rho(x)) = \rho(x)$, so that $\rho(x) \in x/\equiv$, and each $y \in x/\equiv$ fulfills $y \leq \rho(y) = \rho(x)$, thus $\rho(x) = \max(x/\equiv) = \max(\rho(x)/\equiv)$ since $x/\equiv = \rho(x)/\equiv$. Therefore $\max(0/\equiv) = \rho(0)$, thus $0/\equiv = (\rho(0)]_L$, since $0/\equiv \in \operatorname{Id}(L)$. It also follows that $R(L) = \{\rho(x) \mid x \in L\} = \{\max(x/\equiv) \mid x \in L\}$ and, also, $R(L) = \{x \in L \mid x = \rho(x)\} = \{x \in L \mid x = \max(x/\equiv)\}$. Thus, for any $x \in L$, $\rho(x) \in [x)_L \cap R(L)$ and, for any $r \in [x)_L \cap R(L)$, we have $r \geq x$, thus $r = \rho(r) \geq \rho(x)$, hence $\rho(x) = \min([x)_L \cap R(L))$.

(ii) By (i), $\sim \subseteq \equiv$. Obviously, $\sim \supseteq Cg_L(\{(x \land x, [x, x]) \mid x \in L\}) = Cg_L(\{(x, [x, x]) \mid x \in L\})$. Each $x \in L$ fulfills $x \equiv \rho(x)$, hence $\sigma \subseteq \equiv$. If $x, y \in L$ fulfill $x \equiv y$, then $x \sigma \rho(x) = \rho(y) \sigma y$, hence $x \sigma y$, therefore $\sigma \subseteq \equiv$. Hence $\equiv = \sigma$.

(iii) Let $x \in L$ such that x/\equiv has a minimum and $a \in [\min(x/\equiv))_L$. Then $\min(x/\equiv) \leq a$, $(x, \min(x/\equiv)) \in \equiv$ and $(\min(x/\equiv), [\min(x/\equiv), \min(x/\equiv)]) \in \approx \subseteq \equiv$ by (ii), so that $\min(x/\equiv) = \min(\min(x/\equiv)/\equiv) \leq [\min(x/\equiv), \min(x/\equiv)] \leq [a, \min(x/\equiv)] \leq \min(x/\equiv)$, hence $[\min(x/\equiv), \min(x/\equiv)] = [a, \min(x/\equiv)] = \min(x/\equiv)$.

(iv) Let $a \in L$, arbitrary, and $x \in L$ such that $\min(x/\equiv)$ exists, so that $x/\equiv = \lfloor \min(x/\equiv), \rho(x) \rceil_L$ by (i).

For all $n \in \mathbb{N}^*$, $(a, a^2) \in \mathbb{R} \subseteq \mathbb{C}$ and, if $(a, a^n) \in \mathbb{C}$, then $a = a \wedge a \sim a \wedge a^n \sim [a, a^n] = a^{n+1}$. Thus, by (ii), $(a^n, a) \in \mathbb{C} \subseteq \mathbb{E}$ for all $n \in \mathbb{N}^*$, hence $(\min(x/\mathbb{E}), \max(x/\mathbb{E})) = (\rho(x)^{nx}, \rho(x)) \in \mathbb{C}$, therefore $\mathbb{E} \subseteq \mathbb{C}$, so $\mathbb{E} = \mathbb{C}$.

If $a \in x/\equiv$ and $n \in \mathbb{N}$ with $n \ge n_x$, then, by the above, $a^n \equiv a \equiv x$, hence, by (ii) and (iii), $\min(x/\equiv) \le a \le \rho(x)$, thus $\min(x/\equiv) \le a^n \le a^{n_x} \le \rho(x)^{n_x} = \min(x/\equiv)$, hence $a^n = \min(x/\equiv)$.

Remark 5.10. Of course, for all $a \in L$, since $a \leq \rho(a)$, $\rho(a) = 0$ implies a = 0. By Proposition 5.9, (i), $0 \in R(L)$ iff $\rho(0) = 0$ iff $0/\equiv = \{0\}$ iff $(\forall a \in L) (\rho(a) = \rho(0) \Leftrightarrow a = 0)$ iff $(\forall a \in L) (\rho(a) = 0 \Leftrightarrow a = 0)$.

Remark 5.11. Note from Proposition 5.9, (i), that each class of \equiv contains exactly one element of R(L), namely the maximum of that class.

 $= \cap R(L)^2 = \Delta_{R(L)}, \text{ because, for all } x, y \in L, \ \rho(x) \equiv \rho(y) \text{ iff } \rho(x) = \rho(\rho(x)) = \rho(\rho(y)) = \rho(y). \text{ Hence, if } R(L) = L, \text{ then } \equiv \Delta_L. \text{ Moreover:} R(L) = L \text{ iff } x = \rho(x) \text{ for all } x \in L \text{ iff } Cg_L(\{(x, \rho(x)) \mid x \in L\}) = \Delta_L \text{ iff } \equiv \Delta_L. \text{ Obviously, if } \equiv \Delta_L, \text{ then we also have } Cg_L(\{(x \land y, [x, y]) \mid x, y \in L\}) = Cg_L(\{(x, [x, x]) \mid x \in L\}) = \Delta_L, \text{ thus } [\cdot, \cdot] = \wedge.$

If L is algebraic and $\operatorname{Smi}(L) \subseteq \operatorname{Spec}_L$, in particular if L is algebraic and $[\cdot, \cdot] = \wedge$ (see Remark 5.3) or L has finite length and all $x \in \operatorname{Smi}(L) = \operatorname{Mi}(L)$ fulfill $[x^+, x^+] \not\leq x$ (see Lemma 5.2, (i))), in particular if L has finite length and $[\cdot, \cdot] = \wedge$, then R(L) = L, thus $\equiv = \Delta_L$.

Now recall that: [x, x] = x for all $x \in L$ iff $[x, y] = x \land y$ for all $x, y \in L$. Indeed, the converse implication is trivial, while, if [x, x] = x for all $x \in L$, then we have, for all $x, y \in L$: $x \land y \ge [x, y] \ge [x \land y, x \land y] = x \land y$, therefore $[x, y] = x \land y$. Hence: $Cg_L(\{(x, [x, x]) \mid x \in L\}) = \Delta_L$ iff $Cg_L(\{(x \land y, [x, y]) \mid x, y \in L\}) = \Delta_L$.

Remark 5.12. Obviously, all congruences of a lattice of finite length are complete. Note, also, that a distributive lattice of finite length is a frame.

Let M be a lattice and $\theta \in \operatorname{Con}(M)$. Then, for any $x, y \in M$ such that $x/\theta, y/\theta, (x \lor y)/\theta$ and $(x \land y)/\theta$ have minima and maxima: $\min((x \lor y)/\theta) = \min(x/\theta) \lor \min(y/\theta)$, $\max((x \land y)/\theta) = \max(x/\theta) \land \max(y/\theta)$ and: $x/\theta \le y/\theta$ iff $\min(x/\theta) \le \min(y/\theta)$ iff $\max(x/\theta) \le \max(y/\theta)$. Indeed, the argument below for the family $(x_i)_{i \in I} \subseteq M$ holds, without further hypotheses, for the finite family $\{x, y\}$, hence the first two equalities. Now, if $x/\theta \le y/\theta$, that is $(x \lor y)/\theta = x/\theta \lor y/\theta = y/\theta$, then $\min(y/\theta) = \min((x \lor y)/\theta) = \min(x/\theta) \lor \min(y/\theta)$, hence $\min(x/\theta) \le \min(y/\theta)$. Conversely, since $(x, \min(x/\theta)), (y, \min(y/\theta)) \in \theta$, if $\min(x/\theta) \le \min(y/\theta)$, then $x/\theta = \min(x/\theta)/\theta \le \min(y/\theta)/\theta = y/\theta$. Therefore $x/\theta \le y/\theta$ iff $\min(x/\theta) \le \min(y/\theta)$. Similarly, $x/\theta \le y/\theta$ iff $\max(x/\theta) \le \max(y/\theta)$.

If M is complete and θ preserves arbitrary meets, then each class of θ has a minimum, because, for all $x \in M$, $\bigwedge(x/\theta) \in x/\theta$, thus $\bigwedge(x/\theta) = \min(x/\theta)$. Dually for joins and maxima of classes. Hence, if M is a complete lattice and θ is a complete congruence, then all classes of θ have minima and maxima, so that $x/\theta = |\min(x/\theta), \max(x/\theta)|_M$ for all $x \in M$.

Note, also, that, if M is complete and θ preserves arbitrary joins, then, for any family $(x_i)_{i \in I} \subseteq M$, there exists in $M/\theta \bigvee_{i \in I} x_i/\theta = (\bigvee_{i \in I} x_i)/\theta$, thus M/θ is a complete lattice. Similarly if θ preserves arbitrary meets.

If M is a complete lattice and θ is a complete congruence, then, for any nonempty family $(x_i)_{i\in I} \subseteq M$, we have, in the complete lattice M/θ : $\min(\bigvee_{i\in I} x_i/\theta) = \bigvee_{i\in I} \min(x_i/\theta)$ and $\max(\bigwedge_{i\in I} x_i/\theta) = \bigwedge_{i\in I} \max(x_i/\theta)$. Indeed, if we denote by $a_i = \min(x_i/\theta)$ for all $i \in I$ and by $a = \min((\bigvee_{i\in I} x_i)/\theta) = \min(\bigvee_{i\in I} x_i/\theta)$, then, since $a_i \in x_i/\theta$ for all $i \in I$, we have $\bigvee_{i\in I} a_i \in (\bigvee_{i\in I} x_i)/\theta = \bigvee_{i\in I} x_i/\theta$, hence $\bigvee_{i\in I} a_i \ge a$. For all $k \in I$, $a \land a_k \in (\bigvee_{i\in I} x_i)/\theta \land x_k/\theta = ((\bigvee_{i\in I} x_i) \land x_k)/\theta =$ $((x_k \lor \bigvee_{i\in I \setminus \{k\}} x_i) \land x_k)/\theta = x_k/\theta$, thus $a \ge a \land a_k \ge a_k$, hence $a \ge \bigvee_{i\in I} a_i$. Therefore $a = \bigvee_{i\in I} a_i$, that is $\min(\bigvee_{i\in I} x_i/\theta) = \bigvee_{i\in I} \min(x_i/\theta)$. By duality, it follows that we also have $\max(\bigwedge_{i\in I} x_i/\theta) = \bigwedge_{i\in I} \max(x_i/\theta)$.

For any lattice M and any $\theta \in \operatorname{Con}(M)$ such that all classes of θ have minima, let us denote by $[\cdot, \cdot]_{\theta}$ the binary operation on $\operatorname{Con}(M)$ defined by: $[x, y]_{\theta} = \min((x \wedge y)/\theta)$ for all $x, y \in M$. **Proposition 5.13.** If M is a complete lattice and $\theta \in Con(M)$ is a complete congruence such that M/θ is a frame, then:

- $(M, [\cdot, \cdot]_{\theta})$ is a commutator lattice;
- $\theta = Cg_M(\{(x \land y, [x, y]_{\theta}) \mid x, y \in M\}) = Cg_M(\{(x, [x, x]_{\theta}) \mid x \in M\}) = Cg_M(\{(\min(x/\theta), \max(x/\theta)) \mid x \in M\}) = Cg_M(\{(x, \max(x/\theta)) \mid x \in M\});$
- w.r.t. the commutator operation $[\cdot, \cdot]_{\theta}$, $\operatorname{Spec}_M = \operatorname{Mi}(M) \cap \{\max(x/\theta) \mid x \in M \setminus 1/\theta\}$ and $R(M) \subseteq \{\max(x/\theta) \mid x \in M\}$;
- if M has finite length, then $R(M) = \{\max(x/\theta) \mid x \in M\}, \rho(y) = \max(y/\theta)$ for all $y \in M$, and $\theta = Cg_M(\{(x, \rho(x)) \mid x \in M\}).$

Proof. We are using Remark 5.12.

For each $x \in M$, x/θ has a minimum and a maximum, so that $x/\theta = \lfloor \min(x/\theta), \max(x/\theta) \rfloor_M$. For all $x, y \in M$, $[x, y]_\theta = \min((x \land y)/\theta) = [y, x]_\theta \leq x \land y$ and $[x, y]_\theta \in (x \land y)/\theta$, thus $[x, y]_\theta/\theta = (x \land y)/\theta$. Trivially, $[\cdot, \cdot]_\theta$ distributes over $\bigvee \emptyset = 0$. Now let us consider a nonempty family $(x_i)_{i \in I} \subseteq M$ and and $a \in M$. Since M/θ is a frame, we get: $[\bigvee_{i \in I} x_i, a]_\theta = \min(((\bigvee_{i \in I} x_i) \land a)/\theta) = \min((\bigvee_{i \in I} (x_i \land a))/\theta) = \bigvee_{i \in I} \min((x_i \land a)/\theta) = \bigvee_{i \in I} [x_i, a]_\theta$. Therefore $(M, [\cdot, \cdot]_\theta)$ is a commutator lattice.

Let us denote by $\approx = Cg_M(\{(x, [x, x]_{\theta}) \mid x \in M\})$ and by $\sim = Cg_M(\{(x \land y, [x, y]_{\theta}) \mid x, y \in M\})$. Then $\approx = Cg_M(\{(x \land x, [x, x]_{\theta}) \mid x \in M\}) \subseteq \sim \subseteq \theta$ since $(x \land y, [x, y]_{\theta}) \in \theta$ for all $x, y \in M$. But, for all $x \in M$, $[\max(x/\theta), \max(x/\theta)]_{\theta} = \min((\max(x/\theta) \land \max(x/\theta))/\theta) = \min(\max(x/\theta)/\theta) = \min(x/\theta)$ since $(x, \max(x/\theta)) \in \theta$. Therefore $(\max(x/\theta), \min(x/\theta)) = (\max(x/\theta), [\max(x/\theta), \max(x/\theta), \max(x/\theta)]_{\theta}) \in \approx$. Hence, for all $y, z \in M$ such that $(y, z) \in \theta$, we have $\min(y/\theta) = \min(z/\theta) \leq y, z \leq \max(y/\theta) = \max(z/\theta)$, and $(\min(y/\theta), \max(y/\theta)) \in \approx$, hence $(y, z) \in \approx$. Thus $\theta \subseteq \approx$. Therefore $\theta = \sim = \approx = Cg_M(\{(\min(x/\theta), \max(x/\theta)) \mid x \in M\}) = Cg_M(\{(x, \max(x/\theta)) \mid x \in M\})$, since, as above, all inclusions hold.

By Lemma 5.2, (ii), $\operatorname{Spec}_M = \{x \in \operatorname{Mi}(M) \setminus \{1\} \mid (\forall a \in M) ([a, a]_{\theta} \leq x \Rightarrow a \leq x)\}$. Thus, if $x \in \operatorname{Spec}_M$ and $a = \max(x/\theta)$, then $[a, a]_{\theta} = \min(x/\theta) \leq x$, thus $\max(x/\theta) = a \leq x \leq \max(x/\theta)$, hence $x = \max(x/\theta)$. For all $x \in M$ such that $x = \max(x/\theta)$ and all $a \in M$, if $\min(a/\theta) = [a, a]_{\theta} \leq x$, then $a/\theta = \min(a/\theta)/\theta \leq x/\theta$, hence $a \leq \max(a/\theta) \leq \max(x/\theta) = x$, thus, if $x \in \operatorname{Mi}(M) \setminus \{1\}$, then $x \in \operatorname{Spec}_M$. Therefore $\operatorname{Spec}_M = \{x \in \operatorname{Mi}(M) \setminus \{1\} \mid x = \max(x/\theta)\}$, otherwise written $\operatorname{Spec}_M = \operatorname{Mi}(M) \cap \{\max(x/\theta) \mid x \in M \setminus 1/\theta\}$, since, clearly, $\{x \in M \mid x = \max(x/\theta)\} = \{\max(x/\theta) \mid x \in M\}$ and $1 = \max(1/\theta)$.

The set $\{\max(x/\theta) \mid x \in M\}$ is closed w.r.t. arbitrary meets, in particular all meets of prime elements of M belong to $\{\max(x/\theta) \mid x \in M\}$, that is $R(M) \subseteq \{\max(x/\theta) \mid x \in M\}$. $R(M) \supseteq \{1\} \cup \operatorname{Spec}_M = \{1\} \cup (\operatorname{Mi}(M) \cap \{\max(x/\theta) \mid x \in M \setminus 1/\theta\})$ by the above. Let $x \in M$ such that $x = \max(x/\theta)$, but $x \neq 1$ and $x \notin \operatorname{Mi}(M)$, so that $x \notin 1/\theta$ and $x = a \wedge b$ for some $a, b \in M$ with x < a and x < b. Then $x = \max(x/\theta) = \max((a \wedge b)/\theta) = \max(a/\theta) \wedge \max(b/\theta)$, so x is a meet of two elements of $\{\max(y/\theta) \mid y \in M\}$, both of which are strictly greater than x, so that none of them equals 1, thus none of them belongs to $1/\theta$. If the lattice M has finite length, it follows that x is a finite meet of elements of $\operatorname{Mi}(M) \cap \{\max(y/\theta) \mid y \in M \setminus 1/\theta\} = \operatorname{Spec}_M$, thus $x \in R(M)$. Therefore, if Mhas finite length, then $R(M) = \{\max(x/\theta) \mid x \in M\}$, hence $\rho(x) = \max(x/\theta)$ for all $x \in M$, and thus $\theta = Cg_M(\{(x, \max(x/\theta)) \mid x \in M\}) = Cg_M(\{(x, \rho(x)) \mid x \in M\})$.

Proposition 5.14. Let M be a lattice, $\langle \cdot, \cdot \rangle$ be a binary operation on M, $\gamma = Cg_M(\{(x \land y, \langle x, y \rangle) \mid x, y \in M\})$ and $\theta, \zeta \in Con(M)$ such that all classes of γ , θ and ζ have minima. Then:

- (i) for all $x \in M$, $\min(x/(\theta \cap \zeta)) = \min(x/\theta) \vee \min(x/\zeta)$;
- (ii) $[\cdot, \cdot]_{\theta \cap \zeta} = [\cdot, \cdot]_{\theta} \vee [\cdot, \cdot]_{\zeta};$
- (iii) if $\zeta \subseteq \theta$, then $[\cdot, \cdot]_{\theta} \leq [\cdot, \cdot]_{\zeta}$;
- (iv) $[\cdot, \cdot]_{\gamma} \leq \langle \cdot, \cdot \rangle$.

Proof. (i) Let $a = \min(x/\theta)$, $b = \min(x/\zeta)$ and $c = \min(x/(\theta \cap \zeta)) = \min(x/\theta \cap x/\zeta)$ since $x/(\theta \cap \zeta) = x/\theta \cap x/\zeta$. Then $a \lor b \in x/\theta \cap x/\zeta$, hence $a \lor b \ge c$. But $c \land a \in x/\theta$, hence $c \ge c \land a \ge a$ and, analogously, $c \ge b$, thus $c \ge a \lor b$. Therefore $c = a \lor b$.

(ii) By (i).

(iii) By (ii), or simply noticing that $\zeta \subseteq \theta$ means that, for all $x \in M$, $x/\zeta \subseteq x/\theta$, so that $\min(x/\zeta) \ge \min(x/\theta)$, hence the inequality in the enunciation. (iv) For all $x, y \in M$, $\langle x, y \rangle \in (x \land y)/\gamma$, thus $\langle x, y \rangle \ge \min((x \land y)/\gamma) = [x, y]_{\gamma}$.

Recall that a lattice M with 0 is said to be 0-regular iff, for any $\theta, \zeta \in \text{Con}(M)$, $0/\theta = 0/\zeta$ implies $\theta = \zeta$.

Proposition 5.15. If $(L, [\cdot, \cdot])$ is a commutator lattice and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$, then:

- (i) R(L) = L iff $\equiv = \Delta_L$;
- (ii) if R(L) = L, then $[\cdot, \cdot] = \wedge$ in L, in particular L is a frame;
- (iii) if L is algebraic, in particular if L has finite length, then: R(L) = L iff $[\cdot, \cdot] = \wedge$ in L;
- (iv) if L is 0-regular and $\rho(0) = 0$, then R(L) = L, so $[\cdot, \cdot] = \wedge$ in L and L is a frame.

Proof. Recall from Proposition 5.9, (i), that $\equiv \in \text{Con}(L)$.

(i), (ii), (iii) By Remark 5.11.

(iv) If L is 0-regular, then, by (ii) and Remarks 5.10 and 5.11: if $\rho(0) = 0$, which means that $0/\equiv = \{0\} = 0/\Delta_L$, then $\equiv = \Delta_L$, so that R(L) = L, thus $[\cdot, \cdot] = \wedge$ in L and L is a frame.

Remark 5.16. Let $(L, [\cdot, \cdot])$ be a commutator lattice and consider a $\theta \in Con(L)$ that preserves arbitrary joins and the commutator. If we define $[\cdot, \cdot]_{(\theta)} : L/\theta \times$ $L/\theta \to L/\theta$ by $[x/\theta, y/\theta]_{(\theta)} = [x, y]/\theta$ for all $x, y \in L$, then it is straightforward that $(L/\theta, [\cdot, \cdot]_{(\theta)})$ is a commutator lattice. In particular, L/\equiv is a commutator lattice in which $[\cdot, \cdot]_{(\equiv)} = \wedge$. See also the proof of Proposition 5.17.

Proposition 5.17. If $(L, [\cdot, \cdot])$ is a commutator lattice and $\equiv = \{(a, b) \in L^2 \mid \rho(a)\}$ $= \rho(b)$, then L/\equiv is a frame, thus:

- $\mathcal{A}nn(L/\equiv) = \operatorname{PAnn}(L/\equiv) \subseteq \operatorname{PId}(L/\equiv);$
- $L \equiv is Stone iff L \equiv is strongly Stone.$

Proof. For all $x \in L$ and any family $(y_i)_{i \in I} \subseteq L, x/\equiv \wedge (\bigvee_{i \in I} y_i/\equiv) =$ $x/\equiv \wedge \; (\bigvee_{i\in I} y_i)/\equiv = \; (x \wedge (\bigvee_{i\in I} y_i))/\equiv = \; [x,\bigvee_{i\in I} y_i]/\equiv = \; (\bigvee_{i\in I} [x,y_i])/\equiv = \; (\bigvee_{i\in I} y_i)/\equiv = : (\bigvee_{i\in I} y_i)/\equiv : (\bigvee_{i\in I} y_i)/\cong : (\bigvee_{i\in I} y_i)/\equiv : (\bigvee_{i\in I} y_i)/\cong : (\bigvee_{i\in I} y_i)/\cong$ $\bigvee_{i \in I} [x, y_i] / \equiv = \bigvee_{i \in I} (x \land y_i) / \equiv = \bigvee_{i \in I} (x / \equiv \land y_i / \equiv).$ This also followed from Remarks 3.5 and 5.16.

Lemma 5.18. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$, then:

- (i) for all $a, b \in L$, $a \wedge b = 0$ iff [a, b] = 0;
- (ii) for any $U \subseteq L$, $\operatorname{Ann}_L(U) = \{x \in L \mid (\forall u \in U) ([x, u] = 0)\}.$

Proof. (i) Let $\equiv = \{(a,b) \in L^2 \mid \rho(a) = \rho(b)\}$. By Remarks 5.10 and 5.8, if $\rho(0) = 0$, then, for all $a, b \in L$: $a \wedge b = 0$ iff $a \wedge b \in 0 / \equiv$ iff $[a, b] \in 0 / \equiv$ iff [a, b] = 0. (ii) By (i) and the definition of $\operatorname{Ann}_L(U)$.

Proposition 5.19. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$, then:

- for all $U \subseteq L$, $\operatorname{Ann}_L(U) = \operatorname{Ann}_L((U|_L)) = \operatorname{Ann}_L(\bigvee U);$
- for any family $(I_k)_{k \in K} \subseteq \mathrm{Id}(L)$, $\mathrm{Ann}_L(\bigvee_{k \in K} I_k) = \bigcap_{k \in K} \mathrm{Ann}_L(I_k)$;
- $\mathcal{A}nn(L) = \operatorname{PAnn}(L) \subset \operatorname{PId}(L);$
- L is Stone iff L is strongly Stone.

Proof. By Propositions 5.17 and 4.4 and Proposition 5.9, (i), we have the equalities. By Proposition 4.4, (iii), L is Stone iff it is strongly Stone.

Remark 5.20. By Proposition 5.17, the second part of condition $(4)_{L/\equiv}$ is trivially fulfilled, so that $(4)_{L/\equiv}$ is equivalent to $(iv)_{L/\equiv}$. By Proposition 5.19, if $\rho(0) = 0$, then the second part of condition $(4)_L$ is trivially fulfilled, so that $(4)_L$ is equivalent to $(iv)_L$.

6. TRANSFERRING DAVEY'S THEOREM TO COMMUTATOR LATTICES AND RELATED RESULTS

Throughout this section, unless mentioned otherwise, $(L, \lor, \land, [\cdot, \cdot], 0, 1)$ will be a commutator lattice and, as in Section 5, we will denote by $\equiv = \{(x, y) \in L^2 \mid \rho(x) = \rho(y)\} \in \text{Con}(L)$. Recall from Remark 5.10 that $0/\equiv = \{0\}$ iff $\rho(0) = 0$.

Let us define $\stackrel{\bullet}{\vee}: R(L)^2 \to R(L)$ by $\rho(x) \stackrel{\bullet}{\vee} \rho(y) = \rho(\rho(x) \lor \rho(y))$ for all $x, y \in L$, and, for any family $(x_i)_{i \in I} \subseteq L$, let us denote by $\bigvee_{i \in I} \rho(x_i) = \rho(\bigvee_{i \in I} \rho(x_i)) \in R(L)$.

Remark 6.1. Note from the definitions of \checkmark and \equiv and Proposition 5.9, (i), that \equiv preserves \checkmark over arbitrary families of elements of R(L).

Proposition 6.2. If $(L, [\cdot, \cdot])$ is a commutator lattice and \checkmark is the binary operation defined on R(L) as above, then:

- (i) $(R(L), \lor, \land, \rho(0), 1)$ is a frame, isomorphic to L/\equiv ;
- (ii) in the commutator lattice $(R(L), \checkmark, \land, \land, \rho(0), 1)$, $\operatorname{Spec}_{R(L)} = \operatorname{Spec}_L$ and R(R(L)) = R(L), in particular $\rho(0) \in R(R(L))$;
- (iii) in the commutator lattice $(L/\equiv, \lor, \land, \land, 0/\equiv, 1/\equiv)$, $\operatorname{Spec}_{L/\equiv} = \{p/\equiv \mid p \in V(\rho(0)) = [\rho(0))_L \cap \operatorname{Spec}_L\}$ and $R(L/\equiv) = L/\equiv$, in particular $\rho(0/\equiv) = 0/\equiv$.

Proof. (i) $1 = \rho(1) \in R(L)$ and, for all $a, b \in L$, $\rho(a) \land \rho(b) = \rho(a \land b) \in R(L)$, and $\rho(a) \lor \rho(b) = \rho(\rho(a) \lor \rho(b)) = \rho(a \lor b)$, from which it is straightforward that $(R(L), \lor, \land, \rho(0), 1)$ is a bounded lattice.

Let $f: L \to R(L)$, for all $x \in L$, $f(x) = \rho(x)$. Then f is surjective and, by the above, for all $a, b \in L$, $f(a \wedge b) = f(a) \wedge f(b)$ and $f(a \vee b) = f(a) \checkmark f(b)$, hence f is a surjective lattice morphism. By the definition of \equiv , $\operatorname{Ker}(f) = \equiv$. Hence the Main Isomorphism Theorem gives us a lattice isomorphism $h: L/\equiv \to R(L)$, defined by $h(x/\equiv) = \rho(x)$ for all $x \in L$. By Proposition 5.17, it follows that R(L)is a frame and h is a frame isomorphism.

(ii) Since $(R(L), \stackrel{\bullet}{\vee}, \wedge, \rho(0), 1)$ is a frame by (i), $(R(L), \stackrel{\bullet}{\vee}, \wedge, \wedge, \rho(0), 1)$ is a commutator lattice. Spec_L $\subseteq R(L) = \{\rho(u) \mid u \in L\}$, and, for any $a, b, x \in L$: $\rho(a) \wedge \rho(b) \leq \rho(x)$ iff $\rho(a \wedge b) \leq \rho(x)$ iff $\rho([a, b]) = \rho(a \wedge b) \leq \rho(x)$ iff $[a, b] \leq \rho(x)$, and $a \leq \rho(x)$ iff $\rho(a) \leq \rho(x)$ and similarly for b, hence: $\rho(x) \in \text{Spec}_{R(L)}$ iff $\rho(x) \in \text{Spec}_L$, therefore $\text{Spec}_{R(L)} = \text{Spec}_{R(L)}$. Thus, in R(L), for any $x \in L$, the radical of $\rho(x)$ is $\Lambda\{p \in \text{Spec}_{R(L)} \mid \rho(x) \leq p\} = \Lambda\{p \in \text{Spec}_L \mid \rho(x) \leq p\} = \rho(\rho(x)) = \rho(x)$, which means that every element of the commutator lattice R(L) is a radical element, in particular the first element of this lattice, $\rho(0)$, is a radical element.

(iii) By (i) and (ii) and the definition of the frame isomorphism h.

Remark 6.3. By Remarks 4.1 and 5.10, we have $\operatorname{Ann}_L(U)/\equiv \subseteq \operatorname{Ann}_{L/\equiv}(U/\equiv)$ for any $U \subseteq L$ and, if $\rho(0) = 0$, then the properties of Lemma 4.2 hold for M and θ replaced by L and \equiv , respectively.

For all $x, y \in L$, let us define $x \to y = \bigvee \{a \in L \mid [x, a] \le y\}$ and $\neg x = x \to 0 = \bigvee \{a \in L \mid [x, a] = 0\}.$

Remark 6.4. Let $x, y \in L$. By Remark 5.4, $x \to y = \max\{a \in L \mid [x, a] \leq y\}$ and $\neg x = \max\{a \in L \mid [x, a] = 0\}$, so that, by Lemma 5.18, (ii), if $\rho(0) = 0$, then $\neg x = \bigvee_{a \in \operatorname{Ann}_L(x)} a = \max(\operatorname{Ann}_L(x)).$

Lemma 6.5. If $(L, [\cdot, \cdot])$ is a commutator lattice, then, for all $x, y, z \in L$:

- (i) $[x, y] \leq z \text{ iff } x \leq y \rightarrow z;$
- (ii) if [y,1] = y, then: $y \to z = 1$ iff $y \le z$.

Proof. (i) $y \to z = \max\{a \in L \mid [a, y] \le z\}$, so both implications hold. (ii) If [y, 1] = y, then, by (i): $y \to z = 1$ iff $1 \le y \to z$ iff $y = [1, y] \le z$.

Lemma 6.6. If $(L, [\cdot, \cdot])$ is a commutator lattice, then, for all $e \in \mathcal{B}(L)$ and all $a \in L$ such that $[1, e \land a] = e \land a$, we have $e \land a = [e, a]$.

Proof. Since $e \in \mathcal{B}(L)$, we have $e \lor f = 1$ and $e \land f = 0$ for some $f \in L$. Then $[e, a] \le e \land a = [1, e \land a] = [e \lor f, e \land a] = [e, e \land a] \lor [f, e \land a] \le [e, e \land a] \lor (f \land e \land a) = [e, e \land a] \lor 0 = [e, e \land a] \le [e, a]$, hence $e \land a = [e, a]$.

Proposition 6.7. If $(L, [\cdot, \cdot])$ is a commutator lattice such that [x, 1] = x for all $x \in L$, then $e \wedge a = [e, a]$ for all $e \in \mathcal{B}(L)$ and all $a \in L$, in particular $[\cdot, \cdot] = \wedge$ in $\mathcal{B}(L)$, and $\mathcal{B}(L)$ is a Boolean sublattice of L whose complementation is defined by $\neg e = e \rightarrow 0 = \max\{a \in L \mid [e, a] = 0\} = \max(\operatorname{Ann}_L(e))$ for all $e \in \mathcal{B}(L)$.

Proof. By Lemma 6.6, $e \wedge a = [e, a]$ for all $e \in \mathcal{B}(L)$ and all $a \in L$, in particular for all $e, a \in \mathcal{B}(L)$.

Obviously, $0, 1 \in \mathcal{B}(L)$. Now let $x, y \in \mathcal{B}(L)$, so that $x \vee \overline{x} = y \vee \overline{y} = 1$ and $x \wedge \overline{x} = y \wedge \overline{y} = 0$ for some $\overline{x}, \overline{y} \in \mathcal{B}(L)$, so that $[x, \overline{x}] = [y, \overline{y}] = 0$, as well. Then $[x \vee y, \overline{x} \wedge \overline{y}] = [x, \overline{x} \wedge \overline{y}] \vee [y, \overline{x} \wedge \overline{y}] \leq [x, \overline{x}] \vee [y, \overline{y}] = 0$, so $[x \vee y, \overline{x} \wedge \overline{y}] = 0$. By Lemma 6.6, $[x, x \vee y] = x \wedge (x \vee y) = x$, hence $x \vee y \vee (\overline{x} \wedge \overline{y}) = x \vee y \vee [\overline{x}, \overline{y}] = [1, x \vee y] \vee [\overline{x}, \overline{y}] = [x \vee \overline{x}, x \vee y] \vee [\overline{x}, \overline{y}] = [x, x \vee y] \vee [\overline{x}, \overline{y}] = x \vee [\overline{x}, x \vee y \vee \overline{y}] = x \vee [\overline{x}, 1] = x \vee \overline{x} = 1$, hence $x \vee y \in \mathcal{B}(L)$ and $\overline{x} \wedge \overline{y} \in \mathcal{B}(L)$, thus also $x \wedge y \in \mathcal{B}(L)$, since we can interchange x and \overline{x} , respectively y and \overline{y} in the above.

Therefore $\mathcal{B}(L)$ is a bounded sublattice of L in which the meet coincides to $[\cdot, \cdot]$, thus $\mathcal{B}(L)$ is a bounded distributive lattice, and it is clearly complemented, so $\mathcal{B}(L)$ is a Boolean sublattice of L. Let $e \in \mathcal{B}(L)$. As in every Boolean algebra, the complement of e in $\mathcal{B}(L)$ is max $\{a \in \mathcal{B}(L) \mid e \land a = 0\} \leq \max\{a \in L \mid e \land a = 0\} = \max\{a \in L \mid [e, a] = 0\} = \neg e$, hence $e \lor \neg e = 1$. But, by the above, $\neg e = \max(\operatorname{Ann}_{L}(e)) \in \operatorname{Ann}_{L}(e)$, thus $e \land \neg e = 0$. Hence $\neg e \in \mathcal{B}(L)$ and $\neg e$ is the complement of e in $\mathcal{B}(L)$.

Remark 6.8. If [1,1] = x < 1, then no $y \in [x)_L$ can be prime, thus $\rho(x) = \bigwedge \emptyset = 1 = \rho(1)$, hence $1 \neq x \in 1/\Xi$. Therefore $1/\Xi = \{1\}$ implies [1,1] = 1.

Proposition 6.9. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$, then:

- (i) for all $x \in L$, $\operatorname{Ann}_L(x) = (\neg x]_L$;
- (ii) L is a Stone lattice iff, for all $x \in L$, $\neg x \in \mathcal{B}(L)$;
- (iii) if [x,1] = x for all $x \in L$, then, for all $e \in \mathcal{B}(L)$, $(e]_L = \operatorname{Ann}_L(\neg e) \in \operatorname{PAnn}(L)$.

Proof. (i) By Proposition 5.19 and Remark 6.4.

(ii) By (i) and the definition of a Stone lattice.

(iii) By (i) and Proposition 6.7, for all $e \in \mathcal{B}(L)$, $(e]_L = (\neg \neg e]_L = \operatorname{Ann}_L(\neg e) \in \operatorname{PAnn}(L)$.

Proposition 6.10. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$ and [x, 1] = x for all $x \in L$, then:

- $\mathcal{B}(\mathrm{Id}(L)) = \{(e|_L \mid e \in \mathcal{B}(L)\} \subseteq \mathrm{PAnn}(L) \subseteq \mathcal{A}nn(L);$
- L is a Stone lattice iff $PAnn(L) = \mathcal{B}(Id(L));$
- L is a strongly Stone lattice iff $Ann(L) = \mathcal{B}(Id(L))$.

Proof. By Proposition 4.12, (ii), and Proposition 6.9, (iii).

Let us see, in the following proposition, some side results on compact elements.

Proposition 6.11. Let $(L, [\cdot, \cdot])$ be a commutator lattice and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}.$

- (i) If $1 \in \operatorname{Cp}(L)$, then $\{x \in \mathcal{B}(L) \mid [x, 1] = x\} \subseteq \operatorname{Cp}(L)$.
- (ii) If $1 \in Cp(L)$ and [x, 1] = x for all $x \in \mathcal{B}(L)$, then $\mathcal{B}(L) \subseteq Cp(L)$.
- (iii) If $1 \in \operatorname{Cp}(L)$ and $1/\equiv = \{1\}$, then $1/\equiv \in \operatorname{Cp}(L/\equiv)$, $\mathcal{B}(L/\equiv) \subseteq \operatorname{Cp}(L/\equiv)$ and, in L/\equiv , $V(x/\equiv) \neq \emptyset$ for all $x \in L \setminus \{1\}$.

_

Proof. (i) Let $x \in \mathcal{B}(L)$, so that $x \lor y = 1$ and $x \land y = 0$ for some $y \in L$, so we also have [x, y] = 0. Assume that [x, 1] = x and $1 \in \operatorname{Cp}(L)$, and let $\emptyset \neq M \subseteq L$ such that $x \leq \bigvee M$, so that $1 = x \lor y \leq \bigvee M \lor y$, therefore, since $1 \in \operatorname{Cp}(L)$, $1 = x \lor y = \bigvee_{i=1}^{n} x_i \lor y$ for some $n \in \mathbb{N}^*$ and some $x_1, \ldots, x_n \in M$. Then $x = [x, 1] = [x, \bigvee_{i=1}^{n} x_i \lor y] = [x, \bigvee_{i=1}^{n} x_i] \lor [x, y] = [x, \bigvee_{i=1}^{n} x_i] \leq \bigvee_{i=1}^{n} x_i$, hence $x \leq \bigvee_{i=1}^{n} x_i$.

(ii) By (i).

(iii) Let $\emptyset \neq M \subseteq L$ such that $(\bigvee_{x \in M} x) / \equiv = \bigvee_{x \in M} x / \equiv = 1 / \equiv$. If $1/\equiv = \{1\}$, then it follows that $\bigvee_{x \in M} x = 1$. If, furthermore, $1 \in \operatorname{Cp}(L)$, then we obtain that $1 = \bigvee_{i=1}^{n} x_i = 1$ for some $n \in \mathbb{N}^*$ and some $x_1, \ldots, x_n \in M$, hence $1/\equiv = (\bigvee_{i=1}^{n} x_i) / \equiv = \bigvee_{i=1}^{n} x_i / \equiv$, therefore $1/\equiv \in \operatorname{Cp}(L/\equiv)$.

Since L/\equiv is a commutator lattice with $[\cdot, \cdot] = \wedge$, L/\equiv fulfills $[x/\equiv, 1/\equiv] = x/\equiv \wedge 1/\equiv = x/\equiv$ for all $x \in L$. Now apply (ii) to obtain that $\mathcal{B}(L/\equiv) \subseteq \operatorname{Cp}(L/\equiv)$, and Lemma 5.2, (v), to obtain that, in L/\equiv , $V(x/\equiv) \neq \emptyset$ for all $x \in L \setminus 1/\equiv = L \setminus \{1\}$.

Proposition 6.12. If $(L, [\cdot, \cdot])$ is a commutator lattice such that [x, 1] = x for all $x \in L$ and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$, then:

- (i) $p_{\equiv} \mid_{\mathcal{B}(L)} : \mathcal{B}(L) \to \mathcal{B}(L/\equiv)$ is a Boolean morphism, which is injective iff $1/\equiv \cap \mathcal{B}(L) = \{1\}$ iff $0/\equiv \cap \mathcal{B}(L) = \{0\}$;
- (ii) if $\rho(0) = 0$ or $1/\equiv \{1\}$, in particular if $\rho(0) = 0$ or $1 \in Cp(L)$, then the Boolean morphism $p_{\equiv}|_{\mathcal{B}(L)}: \mathcal{B}(L) \to \mathcal{B}(L/\equiv)$ is injective;
- (iii) if $\rho(0) = 0$ and $1/\equiv \{1\}$, in particular if $\rho(0) = 0$ and $1 \in Cp(L)$, then:
 - for all $x \in L$: $x \equiv \mathcal{B}(L)$ iff $x \in \mathcal{B}(L)$;
 - $p_{\equiv} \mid_{\mathcal{B}(L)} : \mathcal{B}(L) \to \mathcal{B}(L/\equiv)$ is a Boolean isomorphism.

Proof. Assume that [x, 1] = x for all $x \in L$, so that $\mathcal{B}(L)$ is a Boolean sublattice of L by Proposition 6.7.

(i) By the above and Remark 4.9.

(ii) By (i) and Remark 5.10, with Lemma 5.7 for the particular case.

(iii) By Lemma 4.13, (i) and Remark 5.10, with Lemma 5.7 for the particular case. $\hfill\blacksquare$

Remark 6.13. By Proposition 6.12, (iii), if $1/\equiv \{1\}$ and $\rho(0) = 0$, then \equiv has the BLP (see [9]).

Proposition 6.14. Let $(L, [\cdot, \cdot])$ be a commutator lattice such that $\rho(0) = 0$ and [x, 1] = x for all $x \in L$, $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}, U \subseteq L, a \in L$ and $e \in \mathcal{B}(L)$. Then:

(i)
$$a/\equiv \leq e/\equiv iff \ a \leq e; \ e = \max(e/\equiv) = \rho(e); \ \mathcal{B}(L) \subseteq R(L);$$

- (ii) if L is complemented, then R(L) = L, $[\cdot, \cdot] = \wedge$ in L and L is a complete Boolean algebra;
- (iii) $(\rho(a)]_L \equiv \operatorname{Ann}_L(U) \equiv iff(\rho(a)]_L = \operatorname{Ann}_L(U); (e]_L \equiv \operatorname{Ann}_L(U) = iff(e]_L = \operatorname{Ann}_L(U).$

Proof. (i) By Lemma 4.11, (ii), Proposition 6.7, Proposition 5.9, (i), and Remark 4.5, $e = \max(e/\equiv) = \rho(e) \in R(L)$, so that $\mathcal{B}(L) \subseteq R(L)$ and $a/\equiv \leq e/\equiv$ iff $a \leq e$. (ii) By (i), if L is complemented, then $L = \mathcal{B}(L) \subseteq R(L) \subseteq L$, so that $L = \mathcal{B}(L) = R(L)$, thus L is a complete Boolean algebra and has $[\cdot, \cdot] = \wedge$ by Proposition 6.7 and the fact that L is a complete lattice.

(iii) By (i), Lemma 4.11, (iii), and Proposition 5.9, (i).

Proposition 6.15. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$ and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$, then, for any cardinality κ :

- (i) (1)_{κ,L} implies (1)_{$\kappa,L/\equiv$} (that is, if L is κ -Stone, then L/\equiv is κ -Stone);
- (ii) if $1/\equiv = \{1\}$ and [x,1] = x for all $x \in L$, in particular if $1 \in \operatorname{Cp}(L)$ and [x,1] = x for all $x \in L$, then properties $(1)_{\kappa,L}$ and $(1)_{\kappa,L/\equiv}$ are equivalent (that is L is κ -Stone iff L/\equiv is κ -Stone).

Proof. (i) By Remark 5.10 and Proposition 4.14, (i).

(ii) By Remark 5.10, Proposition 6.14, (i), and Proposition 4.14, (ii), with Lemma 5.7 for the particular case. ■

Corollary 6.16. Let $(L, [\cdot, \cdot])$ be a commutator lattice such that $\rho(0) = 0$ and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}.$

- If L is Stone, respectively strongly Stone, then L/\equiv is Stone, respectively strongly Stone.
- If $1/\equiv = \{1\}$ and [x, 1] = x for all $x \in L$, in particular if $1 \in Cp(L)$ and [x, 1] = x for all $x \in L$, then L is Stone, respectively strongly Stone, iff L/\equiv is Stone, respectively strongly Stone.

Proof. By Proposition 6.15 applied for $\kappa = 1$, then for all cardinalities κ .

Proposition 6.17. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$ and $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$, then, for any cardinality κ :

- (i) if $1/\equiv = \{1\}$, in particular if $1 \in Cp(L)$ and [1,1] = 1, then $(2)_{\kappa,L}$ implies $(2)_{\kappa,L/\equiv}$;
- (ii) if $1/\equiv = \{1\}$ and [x, 1] = x for all $x \in L$, in particular if $1 \in \operatorname{Cp}(L)$ and [x, 1] = x for all $x \in L$, then properties $(2)_{\kappa,L}$ and $(2)_{\kappa,L/\equiv}$ are equivalent.

Proof. We get the particular cases from Lemma 5.7.

- (i) By Remark 5.10 and Proposition 4.15, (i).
- (ii) By Remark 5.10, Proposition 6.14, (i), and Proposition 4.15, (ii).

Proposition 6.18. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$ and $\equiv = \{(a,b) \in L^2 \mid \rho(a) = \rho(b)\}, \text{ then, for any cardinality } \kappa, \text{ the properties } (3)_{\kappa,L}$ and $(3)_{\kappa,L/\equiv}$ are equivalent.

Proof. By Remark 5.10 and Proposition 4.16.

Proposition 6.19. If $(L, [\cdot, \cdot])$ is a commutator lattice such that $\rho(0) = 0$ and $\equiv = \{(a,b) \in L^2 \mid \rho(a) = \rho(b)\}, \text{ then, for any cardinality } \kappa, \text{ the properties } (iv)_L,$ $(4)_{\kappa,L}$ and $(4)_{\kappa,L/\equiv}$ are equivalent.

Proof. By Remarks 5.10 and 5.20 and Proposition 4.17.

Proposition 6.20. If $(L, [\cdot, \cdot])$ is a commutator lattice and $\equiv = \{(a, b) \in L^2 \mid \rho(a)\}$ $= \rho(b)$, then, for any cardinality κ :

- (i) $(5)_{\kappa,L}$ implies $(5)_{\kappa,L/\equiv}$;
- (ii) if $\rho(0) = 0$ and $1/\equiv = \{1\}$, in particular if $\rho(0) = 0, 1 \in Cp(L)$ and [1,1] = 1, then $(5)_{\kappa,L}$ is equivalent to $(5)_{\kappa,L/\equiv}$.

Proof. (i) By Proposition 4.18, (i).

(ii) By Remark 5.10 and Proposition 4.18, (ii), with Lemma 5.7 for the particular case.

Theorem 6.21. Let $(L, [\cdot, \cdot])$ be a commutator lattice with $\rho(0) = 0$, and consider the congruence $\equiv = \{(a, b) \in L^2 \mid \rho(a) = \rho(b)\}$ of L. If $1/\equiv = \{1\}$ and [x, 1] = xfor all $x \in L$, in particular if $1 \in Cp(L)$ and [x, 1] = x for all $x \in L$, then:

- (i) for any $h, i, j \in \overline{1,5}$ and any nonzero cardinality κ , conditions $(iv)_L$, $(h)_{\kappa,L}$, $(i)_{<\infty,L}$ and $(j)_L$ are equivalent, thus L satisfies the equivalences from Corollary 4.21, (ii);
- (ii) let m be a nonzero cardinality; if, in Definition 3.4, we replace the conditions that L is complete and $[\cdot, \cdot]$ is completely distributive w.r.t. the join by L being closed w.r.t. the joins of all families of elements of cardinalities at most m and $[\cdot, \cdot]$ being distributive w.r.t. such joins, then we get that: for any $h, i \in \overline{1,5}$ and any nonzero cardinality $\kappa \leq m$, conditions $(h)_{\kappa,L}$ and $(i)_{<\infty}$ L are equivalent, thus L satisfies the equivalences from Corollary 4.21, (i).

Proof. We get the particular case from Lemma 5.7.

(i) By Theorem 3.3, (iv), Proposition 6.15, (ii), Proposition 6.17, (ii), Propositions 6.18 and 6.19, Proposition 6.20, (ii), and Proposition 5.17.

(ii) By Theorem 4.19, (ii), Proposition 6.15, (ii), Proposition 6.17, (ii), Propositions 6.18 and 6.19, Proposition 6.20, (ii), and the fact that, in this case, L/\equiv is closed w.r.t. the joins of all families of elements of cardinalities at most κ and has the meet distributive w.r.t. the joins of families of elements of cardinalities at most κ , which follows imediately through an argument analogous to the first proof of Proposition 5.17.

Corollary 6.22. Let $(L, [\cdot, \cdot])$ be a commutator lattice with $\rho(0) = 0$. Then:

- for any nonzero cardinalities κ and μ , conditions $(3)_{\kappa,L}$, $(4)_{\mu,L}$ and $(iv)_L$ are equivalent;
- if $\{a \in L \mid \rho(a) = 1\} = \{1\}$, in particular if $1 \in Cp(L)$ and [1,1] = 1, then, for any nonzero cardinalities κ , λ and μ , conditions $(3)_{\kappa,L}$, $(4)_{\lambda,L}$, $(iv)_L$ and $(5)_{\mu,L}$ are equivalent.

Let m be a nonzero cardinality. If, in Definition 3.4, we replace the conditions that L is complete and $[\cdot, \cdot]$ is completely distributive w.r.t. the join by L being closed w.r.t. the joins of all families of elements of cardinalities at most m and $[\cdot, \cdot]$ being distributive w.r.t. such joins, then:

- for any nonzero cardinalities $\kappa \leq m$ and $\mu \leq m$, conditions $(3)_{\kappa,L}$ and $(4)_{\mu,L}$ are equivalent;
- if $\{a \in L \mid \rho(a) = 1\} = \{1\}$, in particular if $1 \in Cp(L)$ and [1,1] = 1, then, for any nonzero cardinalities $\kappa \leq m$, $\lambda \leq m$ and $\mu \leq m$, conditions $(3)_{\kappa,L}$, $(4)_{\lambda,L}$, and $(5)_{\mu,L}$ are equivalent.

Proof. By Remark 5.10 and Propositions 4.20 and 5.17, with Lemma 5.7 for the particular cases.

7. TRANSFERRING DAVEY'S THEOREM TO CONGRUENCE LATTICES AND PRESERVATION OF THE CONDITIONS FROM DAVEY'S THEOREM BY DIRECT PRODUCTS AND SUBLATTICES

Throughout this section, unless mentioned otherwise, A will be a member of a congruence-modular variety \mathcal{V} . Following [10], we use these notations for the radical of a congruence of A in the commutator lattice $(\operatorname{Con}(A), \lor, \cap, [\cdot, \cdot]_A, \Delta_A, \nabla_A)$ and the lattice congruence \equiv associated to the same commutator lattice: $\rho_A(\alpha) = \bigcap(\operatorname{Spec}(A) \cap [\alpha)_{\operatorname{Con}(A)})$ for all $\alpha \in \operatorname{Con}(A)$ and $\equiv_A = \{(\theta, \zeta) \in \operatorname{Con}(A)^2 \mid \rho_A(\theta) = \rho_A(\zeta)\}.$

Recall from the end of Section 3 that A is semiprime, that is $\rho_A(\Delta_A) = \Delta_A$, if the commutator $[\cdot, \cdot]_A$ of A equals the intersection, in particular if \mathcal{V} is congruencedistributive. Recall, also, that ∇_A is a compact congruence of A if \mathcal{V} is semidegenerate, and that $[\theta, \nabla_A]_A = \theta$ for all $\theta \in \operatorname{Con}(A)$ if \mathcal{V} is semi-degenerate or the commutator of A equals the intersection, in particular if \mathcal{V} is congruencedistributive. Of course, ∇_A is a compact congruence of A in other particular cases such as the case when $\operatorname{Con}(A)$ is compact, in particular when $\operatorname{Con}(A)$ has finite height, in particular when $\operatorname{Con}(A)$ is finite, in particular when A is finite.

Remark 7.1. By Lemma 5.7 and the above, the class of ∇_A w.r.t. \equiv_A is a singleton if \mathcal{V} is semi-degenerate, or if, for instance, $\operatorname{Con}(A)$ is compact and \mathcal{V} is congruence-distributive. Actually, compactness is not necessary and, moreover, all congruence classes of \equiv_A are singletons if the commutator of A equals the intersection (see the remarks after Example 3.6 and Corollary 7.3 below for the converse).

 \equiv_A satisfies the properties from Proposition 5.9, in particular, by (i), \equiv_A is a lattice congruence of Con(A) that preserves arbitrary joins and the commutator of A and satisfies $[\alpha, \beta]_A \equiv_A \alpha \cap \beta$ for all $\alpha, \beta \in \text{Con}(A)$. Moreover, since the meet in Con(A) is the intersection of congruences, the surjectivity of the map $p_{\equiv_A} : \text{Con}(A) \to \text{Con}(A)/\equiv_A$ ensures us that:

Proposition 7.2. If A is a member of a congruence-modular variety and $\equiv_A = \{(\theta, \zeta) \in \operatorname{Con}(A)^2 \mid \rho_A(\theta) = \rho_A(\zeta)\}$, then \equiv_A is a complete congruence of the complete lattice $\operatorname{Con}(A)$, so all its classes are intervals.

Again by Proposition 5.9, the radical congruences of A are the maxima of the classes of \equiv_A and, for each radical congruence θ of A, $\min(\theta/\equiv_A) = \min\{\alpha \in$ $\operatorname{Con}(A) \mid \rho_A(\alpha) = \theta\}$; also, for all $\beta \in \theta/\equiv_A$, we have $[\beta, \min(\theta/\equiv_A)]_A =$ $\min(\theta/\equiv_A)$. By Proposition 5.17, $\operatorname{Con}(A)/\equiv_A$ is a frame, so Proposition 5.15, (iii), gives us:

Corollary 7.3. If A is a member of a congruence-modular variety, then all congruences of A are radical iff the commutator of A equals the intersection of congruences.

Note that, if A is semiprime, then, for $\theta \equiv \equiv_A$, the annihilators in Con(A) satisfy the properties from Lemmas 4.2 and 4.3. Also, if A is semiprime, then the properties from Lemmas 5.18 and 4.11 and Proposition 5.19 hold, in particular:

Corollary 7.4. If A is a semiprime member of a congruence-modular variety, in particular if A belongs to a congruence-distributive variety, then:

- $\mathcal{A}nn(\operatorname{Con}(A)) = \operatorname{PAnn}(\operatorname{Con}(A)) \subseteq \operatorname{PId}(\operatorname{Con}(A));$
- $\operatorname{Con}(A)$ is Stone iff $\operatorname{Con}(A)$ is strongly Stone.

So, if A is semiprime, then all annihilators in Con(A) have maxima, so that Con(A) is Stone iff these maxima are complemented; see also Proposition 6.9, (ii).

By Lemma 6.6 and Proposition 6.7, if \mathcal{V} is semi-degenerate or congruence -distributive, then $[\alpha, \theta]_A = \alpha \cap \theta$ for all $\alpha \in \operatorname{Con}(A)$ and all $\theta \in \mathcal{B}(\operatorname{Con}(A))$ and $\mathcal{B}(\operatorname{Con}(A))$ is a Boolean sublattice of $\operatorname{Con}(A)$ in which the complementation is defined by $\neg \theta = \max(\operatorname{Ann}_{\operatorname{Con}(A)}(\theta))$ for all $\theta \in \mathcal{B}(\operatorname{Con}(A))$. By Proposition 6.11, if \mathcal{V} is semi-degenerate, then all complemented congruences of A are compact.

Remark 7.5. By Proposition 6.10 and Proposition 6.12, (iii), if A is semiprime and $\nabla_A / \equiv_A = {\nabla_A}$, in particular if A is semiprime and \mathcal{V} is semi-degenerate, then the complemented elements of Id(Con(A)) are the principal ideals of Con(A) generated by complemented congruences of A, $\mathcal{B}(\text{Con}(A)) / \equiv_A = \mathcal{B}(\text{Con}(A) / \equiv_A)$, that is \equiv_A has the BLP, and, for all $\theta \in \text{Con}(A)$, we have: $\theta \in \mathcal{B}(\text{Con}(A))$ iff $\theta / \equiv_A \in \mathcal{B}(\text{Con}(A) / \equiv_A)$.

By Proposition 6.12, if \mathcal{V} is semi-degenerate or congruence-distributive, then $p_{\equiv_A} \mid_{\mathcal{B}(\operatorname{Con}(A))} : \mathcal{B}(\operatorname{Con}(A)) \to \mathcal{B}(\operatorname{Con}(A)/\equiv_A)$ is an injective Boolean morphism, which is an isomorphism if A is semiprime and $\nabla_A / \equiv_A = \{\nabla_A\}$, in particular if \mathcal{V} is congruence-distributive or A is semiprime and \mathcal{V} is semi-degenerate.

If A is semiprime, $\nabla_A / \equiv_A = \{\nabla_A\}$ and $[\alpha, \nabla_A]_A = \alpha$ for all $\alpha \in \text{Con}(A)$, in particular if A is semiprime and \mathcal{V} is semi-degenerate or, for instance, \mathcal{V} is congruence-distributive, then Con(A) satisfies the properties from Proposition 6.14.

Corollary 7.6. Let A be a member of a congruence-modular variety \mathcal{V} , $\equiv_A = \{(\theta, \zeta) \in \operatorname{Con}(A)^2 \mid \rho_A(\theta) = \rho_A(\zeta)\}$ and κ be an arbitrary cardinality.

Then $(5)_{\kappa,\operatorname{Con}(A)}$ implies $(5)_{\kappa,\operatorname{Con}(A)/\equiv_A}$.

If A is semiprime, in particular if the commutator of A equals the intersection, in particular if \mathcal{V} is congruence-distributive, then:

- if Con(A) is Stone (equivalently, strongly Stone), then Con(A)/≡_A is Stone (equivalently, strongly Stone); in particular, if κ is nonzero, then, for any cardinality λ, (1)_{κ,Con(A)} implies (1)_{λ,Con(A)/≡_A};
- (3)_{κ ,Con(A)} is equivalent to (3)_{κ ,Con(A)/ \equiv_A ;}
- $(iv)_{\operatorname{Con}(A)}$, $(4)_{\kappa,\operatorname{Con}(A)}$ and $(4)_{\kappa,\operatorname{Con}(A)/\equiv_A}$ are equivalent.

If A is semiprime and $\{\theta \in \operatorname{Con}(A) \mid \rho_A(\theta) = \nabla_A\} = \{\nabla_A\}$, in particular if \mathcal{V} is congruence-distributive or A is semiprime, $\nabla_A \in \operatorname{Cp}(\operatorname{Con}(A))$ and $[\nabla_A, \nabla_A]_A = \nabla_A$, in particular if A is semiprime and \mathcal{V} is semi-degenerate, then:

- $(2)_{\kappa,\operatorname{Con}(A)}$ implies $(2)_{\kappa,\operatorname{Con}(A)/\equiv_A}$;
- $(5)_{\kappa,\operatorname{Con}(A)}$ is equivalent to $(5)_{\kappa,\operatorname{Con}(A)/\equiv_A}$.

If A is semiprime, $\{\theta \in \text{Con}(A) \mid \rho_A(\theta) = \nabla_A\} = \{\nabla_A\}$ and $[\alpha, \nabla_A]_A = \alpha$ for all $\alpha \in \text{Con}(A)$, in particular if \mathcal{V} is congruence-distributive or A is semiprime and \mathcal{V} is semi-degenerate, then:

- Con(A) is Stone iff Con(A) is strongly Stone iff Con(A)/≡_A is Stone iff Con(A)/≡_A is strongly Stone;
- for each $i \in \overline{1,5}$, $(i)_{\kappa,\operatorname{Con}(A)}$ is equivalent to $(i)_{\kappa,\operatorname{Con}(A)/\equiv_A}$.

Proof. By Propositions 6.15, 6.17, 6.18, 6.19 and 6.20 and Corollary 6.16.

Corollary 7.7. Let A be a member of a congruence-modular variety \mathcal{V} . If A is semiprime, $\{\theta \in \operatorname{Con}(A) \mid \rho_A(\theta) = \nabla_A\} = \{\nabla_A\}$ and $[\alpha, \nabla_A]_A = \alpha$ for all $\alpha \in \operatorname{Con}(A)$, in particular if \mathcal{V} is congruence-distributive or A is semiprime and \mathcal{V} is semi-degenerate, then, for any $i, j \in \overline{1,5}$ and any nonzero cardinalities κ and μ , conditions $(iv)_{\operatorname{Con}(A)}$, $(i)_{\kappa,\operatorname{Con}(A)}$ and $(j)_{\mu,\operatorname{Con}(A)}$ are equivalent, and thus each of them is equivalent to $(i)_{<\infty,\operatorname{Con}(A)}$ and to $(i)_{\operatorname{Con}(A)}$ and $\operatorname{Con}(A)$ satisfies the equivalences from Corollary 4.21, (ii).

Corollary 7.8. Let A be a member of a congruence-modular variety \mathcal{V} .

- If A is semiprime, in particular if the commutator of A equals the intersection, in particular if \mathcal{V} is congruence-distributive, then, for any nonzero cardinalities κ and μ , conditions $(3)_{\kappa,\operatorname{Con}(A)}$, $(4)_{\mu,\operatorname{Con}(A)}$ and $(iv)_{\operatorname{Con}(A)}$ are equivalent.
- If $\{\theta \in \operatorname{Con}(A) \mid \rho_A(\theta) = \nabla_A\} = \{\nabla_A\}$, in particular if $\nabla_A \in \operatorname{Cp}(\operatorname{Con}(A))$ and $[\nabla_A, \nabla_A]_A = \nabla_A$ or the commutator of A equals the intersection, in particular if \mathcal{V} is congruence-distributive, then, for any nonzero cardinalities κ, λ and μ , conditions $(3)_{\kappa,\operatorname{Con}(A)}, (4)_{\lambda,\operatorname{Con}(A)}, (iv)_{\operatorname{Con}(A)}$ and $(5)_{\mu,\operatorname{Con}(A)}$ are equivalent.

Proof. By Corollary 6.22.

Remark 7.9. Note that, for any nonempty family $(L_i)_{i\in I}$ of bounded lattices, $\mathcal{B}(\prod_{i\in I} L_i) = \prod_{i\in I} \mathcal{B}(L_i)$ and, for all $(a_i)_{i\in I} \in \prod_{i\in I} L_i$, $\operatorname{Ann}_{\prod_{i\in I} L_i}((a_i)_{i\in I}) = \prod_{i\in I} \operatorname{Ann}_{L_i}(a_i)$ and $((a_i)_{i\in I}]_{\prod_{i\in I} L_i} = \prod_{i\in I}(a_i]_{L_i}$. Moreover, if $pr_j : \prod_{i\in I} L_i \to L_j$ is the canonical projection for each $j \in I$, then, for all $U \subseteq \prod_{i\in I} L_i$, $\operatorname{Ann}_{\prod_{i\in I} L_i}(U) = \prod_{i\in I} \operatorname{Ann}_{L_i}(pr_i(U))$ and $(U]_{\prod_{i\in I} L_i} = \prod_{i\in I}(pr_i(U)]_{L_i}$. Hence, for any cardinality κ and each $h \in \overline{1, 5}$, $(h)_{\kappa, \prod_{i\in I} L_i}$ is satisfied iff $(h)_{\kappa, L_i}$ is satisfied for all $i \in I$.

Corollary 7.10. Let A and B be members of a congruence-modular variety \mathcal{V} and κ be a cardinality.

• If the direct product $A \times B$ has no skew congruences, in particular if \mathcal{V} is semi-degenerate, then: $A \times B$ is semiprime iff A and B are semiprime.

- If the direct product $A \times B$ has no skew congruences and A and B are semiprime, then: $Con(A \times B)$ is Stone iff Con(A) and Con(B) are Stone.
- If the direct product A×B has no skew congruences, A and B are semiprime and B(Con(A)) and B(Con(B)) are closed w.r.t. the intersection, in particular if they are (Boolean) sublattices of Con(A) and Con(B), respectively, then: Con(A×B) is κ-Stone iff Con(A) and Con(B) are κ-Stone.
- If \mathcal{V} is semi-degenerate and A and B are semiprime, then: $\operatorname{Con}(A \times B)$ is κ -Stone iff $\operatorname{Con}(A)$ and $\operatorname{Con}(B)$ are κ -Stone.
- If \mathcal{V} is congruence-distributive, then: $\operatorname{Con}(A \times B)$ is κ -Stone iff $\operatorname{Con}(A)$ and $\operatorname{Con}(B)$ are κ -Stone.

Proof. According to [16, Theorem 5.17, p. 48], for all $\alpha, \theta \in \text{Con}(A)$ and all $\beta, \zeta \in \text{Con}(B)$, we have $[\alpha \times \beta, \theta \times \zeta]_{A \times B} = [\alpha, \theta]_A \times [\beta, \zeta]_B$, hence, if the direct product $A \times B$ has no skew congruences, in particular if \mathcal{V} is congruence-distributive or semi-degenerate ([7, Theorem 8.5, p. 85],[1, Lemma 5.2]), then $\text{Spec}(A \times B) = \{\phi \times \nabla_B, \nabla_A \times \psi \mid \phi \in \text{Spec}(A), \psi \in \text{Spec}(B)\}$, so that $R(\text{Con}(A \times B)) = R(\text{Con}(A)) \times R(\text{Con}(B))$ and hence $A \times B$ is semiprime iff A and B are semiprime (see also [10]).

Since, for any bounded lattice L, if $\mathcal{B}(L)$ is closed w.r.t. the meet, in particular if $\mathcal{B}(L)$ is a (Boolean) sublattice of L, then $1_{1,L}$ is equivalent to $1_{<\infty,L}$, and, if κ is an infinite cardinality, then, for any $U \subseteq A$ and any $V \subseteq B$, we have: $|U| \leq \kappa$ and $|V| \leq \kappa$ iff $|U \times V| \leq \kappa$, by Remark 7.9 we get the statements in the enunciation.

Remark 7.11. Let M be a bounded sublattice of a bounded lattice L and $U, V \subseteq L$. Then it is straightforward that $\operatorname{Ann}_L(U) \cap M \subseteq \operatorname{Ann}_M(U \cap M)$, $(U \cap M]_L \cap M = (U \cap M]_M$ and, if L is distributive, then $(U]_L \cap (V]_L = (U \cap V]_L$.

Lemma 7.12. Let L be a bounded distributive lattice and M a bounded sublattice of L.

- (i) If $U \subseteq M$ is such that $\operatorname{Ann}_L(U) \vee \operatorname{Ann}_L(\operatorname{Ann}_L(U)) = L$, then $\operatorname{Ann}_M(U) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(U)) = M$.
- (ii) For any cardinality κ , $(5)_{\kappa,L}$ implies $(5)_{\kappa,M}$.

Proof. (i) If U is as in the hypothesis, then $\operatorname{Ann}_M(U) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(U)) \supseteq$ $(\operatorname{Ann}_L(U) \cap M) \vee (\operatorname{Ann}_M(\operatorname{Ann}_L(U) \cap M) \supseteq (\operatorname{Ann}_L(U) \cap M) \vee (\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \cap M) \supseteq (\operatorname{Ann}_L(U) \cap M) \cup (\operatorname{Ann}_L(\operatorname{Ann}_L(U)) \cap M) = (\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L(\operatorname{Ann}_L(U)) \cap M)$ M, thus $\operatorname{Ann}_M(U) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(U)) \supseteq (\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L(\operatorname{Ann}_L(U))]_M \cap (M]_M$ $= (\operatorname{Ann}_L(U) \cup \operatorname{Ann}_L\operatorname{Ann}_L(U))]_L \cap M \cap M = (\operatorname{Ann}_L(U) \vee \operatorname{Ann}_L(\operatorname{Ann}_L(U))) \cap M =$ $L \cap M = M$. (ii) Assume that $(5)_{\kappa,L}$ is fulfilled, and let $U \subseteq M \subseteq L$ with $|U| \leq \kappa$, so that $\operatorname{Ann}_L(U) \vee \operatorname{Ann}_L(\operatorname{Ann}_L(U)) = L$, hence $\operatorname{Ann}_M(U) \vee \operatorname{Ann}_M(\operatorname{Ann}_M(U)) = M$ by (i).

Let us assume that the set $\operatorname{Cp}(\operatorname{Con}(A))$ of the compact congruences of A contains ∇_A and is closed w.r.t. the commutator of A. In [10], under these hypotheses we have constructed the *reticulation* $\mathcal{L}(A)$ of A, which, by definition, is a bounded distributive lattice whose prime spectrum of ideals (or filters, but our construction in [10] fulfills this property for ideals) is homeomorphic to the prime spectrum of congruences of A, w.r.t. the Stone topologies. $\mathcal{L}(A)$ is unique modulo a lattice isomorphism and, by our construction from [10]: $\mathcal{L}(A) = \operatorname{Cp}(\operatorname{Con}(A))/\equiv_A$, which is a bounded sublattice of $\operatorname{Con}(A)/\equiv_A$.

Proposition 7.13. Let A be a member of a congruence-modular variety \mathcal{V} such that $\nabla_A \in \operatorname{Cp}(\operatorname{Con}(A))$ and $\operatorname{Cp}(\operatorname{Con}(A))$ is closed w.r.t. the commutator of A. Then, for any cardinality κ , $(5)_{\kappa,\operatorname{Con}(A)}$ implies $(5)_{\kappa,\mathcal{L}(A)}$.

Proof. By Corollary 7.6 and Lemma 7.12, (ii).

Corollary 7.14. Let A be a member of a congruence-modular variety \mathcal{V} such that $\nabla_A \in \operatorname{Cp}(\operatorname{Con}(A))$ and $\operatorname{Cp}(\operatorname{Con}(A))$ is closed w.r.t. the commutator of A. If A is semiprime and $\operatorname{Con}(A)$ is Stone, then $\mathcal{L}(A)$ is strongly Stone.

Proof. By Proposition 7.13, Corollary 7.6, the distributivity of $\mathcal{L}(A)$ and Theorem 3.3, (ii).

8. TRANSFERRING DAVEY'S THEOREM TO COMMUTATIVE UNITARY RINGS

Let us see how we can to obtain versions of Davey's Theorem for the elements of semiprime algebras from congruence-modular varieties by transferring results such as Corollary 7.7 from their congruence lattices. We exemplify here for semiprime commutative unitary rings.

Let $(T, \lor, \land, \odot, \rightarrow, 0, 1)$ be a residuated lattice (otherwise called a commutative integral bounded lattice-ordered monoid), which means that $(T, \lor, \land, 0, 1)$ is a bounded lattice, $(T, \odot, 1)$ is a commutative monoid and \rightarrow is a binary operation on T which fulfills the *law of residuation*: for all $a, b, c \in T$, $a \leq b \rightarrow c$ iff $a \odot b \leq c$. See more about residuated lattices in [8], [12], [17]. Residuated lattices form a semi-degenerate congruence-distributive variety, hence they are semiprime and thus their congruence lattices satisfy Theorem 3.3, (i), and even the equivalences from Corollary 7.7. But they also fulfill a theorem of this form for elements, which can be expressed in the following way, since we notice that the bounded lattice of the filters of T is a bounded sublattice of that of the filters of the underlying bounded lattice of T.

Theorem 8.1 [14, Theorem 5.2.6], [15, Theorem 3.13]. If S is the dual of the underlying bounded lattice of a residuated lattice, then conditions $(1)_{m,S}$, $(2)_{m,S}$, $(3)_{m,S}$, $(4)_{m,S}$ and $(5)_{m,S}$ are equivalent.

In [14, 15], we have proven Theorem 8.1 by transferring the dual of Theorem 3.3 from bounded distributive lattices to residuated lattices through the reticulation functor for residuated lattices.

Remark 8.2. Note from Lemma 6.5 that, if $(L, \lor, \land, [\cdot, \cdot], 0, 1)$ is a commutator lattice in which the operation $[\cdot, \cdot]$ is associative and satisfies [a, 1] = a for all $a \in L$, then L is a complete residuated lattice with the residuation \rightarrow defined above Remark 6.4.

For instance, rings form a semi-degenerate congruence-modular variety with associative commutators, so that, for any commutative unitary ring R, $(\operatorname{Con}(R), \vee, \cap, [\cdot, \cdot]_R, \to, \Delta_R, \nabla_R)$ is a complete residuated lattice.

Throughout the rest of this section, unless mentioned otherwise, $(R, +, \cdot, 0, 1)$ will be a commutative unitary ring.

We denote by $(\mathrm{Id}(R), \vee = +, \cap, [\cdot, \cdot] = \cdot, \{0\}, R)$ the commutator lattice of the ideals of R and by $\iota \gamma_R : \mathrm{Id}(R) \to \mathrm{Con}(R)$ the canonical lattice isomorphism: for all $I \in \mathrm{Id}(R), \, \iota \gamma_R(I) = \{(x, y) \in I^2 \mid x - y \in I\}$. We denote by $\mathrm{Spec}_{\mathrm{Id}}(R) =$ $\mathrm{Spec}_{\mathrm{Id}(R)}$ the set of the prime ideals of R (w.r.t. the commutator operation given by the multiplication of ideals). Recall that $\iota \gamma_R$ preserves the commutator operation, that is $[\iota \gamma_R(I), \iota \gamma_R(J)]_R = \iota \gamma_R(I \cdot J)$ for all $I, J \in \mathrm{Id}(R)$, from which it is easy to deduce that $\iota \gamma_R(\mathrm{Spec}_{\mathrm{Id}}(R)) = \mathrm{Spec}(R)$ and thus $\iota \gamma_R(R(\mathrm{Id}(R))) =$ $R(\mathrm{Con}(R))$. If we denote, for each $I \in \mathrm{Id}(R)$, by $\sqrt{I} = \bigcap\{P \in \mathrm{Spec}_{\mathrm{Id}}(R) \mid I \subseteq P\}$ the radical of I, then note that R is semiprime iff $\{0\} \in R(\mathrm{Id}(R))$ iff $\sqrt{\{0\}} = \{0\}$.

For every $U \subseteq R$, $\langle U \rangle_R$ shall be the ideal of R generated by U, so, for each $x \in R$, $\langle \{x\} \rangle_R = xR$. Let $\operatorname{PId}(R)$ be the set of the principal ideals of R and note that $\operatorname{Cp}(\operatorname{Id}(R))$ is the set of the finitely generated ideals of R. It is straightforward that, for all $x, a, b \in R$, $\iota \gamma_R(xR) = Cg_R(x, 0)$ and $Cg_R(a, b) = Cg_R(a - b, 0)$, hence $\iota \gamma_R(\operatorname{PId}(R)) = \operatorname{PCon}(R)$ and thus $\iota \gamma_R(\operatorname{Cp}(\operatorname{Id}(R))) = \operatorname{Cp}(\operatorname{Con}(R))$. Notice that, for any $k, n \in \mathbb{N}^*$ and any $x_1, \ldots, x_k, y_1, \ldots, y_n \in R$, $\langle \{x_1, \ldots, x_k\} \rangle_R \cdot \langle \{y_1, \ldots, y_k\} \rangle_R = (x_1R + \ldots + x_kR) \cdot (y_1R + \ldots + y_nR) = \langle \{x_iy_j \mid i \in \overline{1,k,j} \in \overline{1,n} \} \rangle_R$, so $\operatorname{Cp}(\operatorname{Id}(R))$ is closed w.r.t. \cdot , thus $\operatorname{Cp}(\operatorname{Con}(R))$ is closed w.r.t. $[\cdot, \cdot]_R$. Let R^* be the reticulation of R, as constructed in [3, 4] (see also [13, 19]): $R^* = \operatorname{Id}(R)/\sim_R$, where \sim_R is the complete lattice congruence of $\operatorname{Id}(R)$ defined by: $\sim_R = \{(I,J) \in (\operatorname{Id}(R))^2 \mid \sqrt{I} = \sqrt{J}\}$ (see also Proposition 7.2); by the above, R^* is isomorphic to $\operatorname{Con}(R)/\equiv_R$. Regarding the results from [3] we are using, note that, since R is commutative, it follows that R is quasicommutative, thus, by $[3, \operatorname{Theorem 3}]$, R fulfills condition (*) from [3].

Remark 8.3. By [3, Lemma, p. 1861], for all $I \in Id(R)$, there exists a $K \in Cp(Id(R))$ such that $K \subseteq I$ and $\sqrt{K} = \sqrt{I}$, hence $R^* = Id(R)/\sim_R = Cp(Id(R))/\sim_R$, therefore $Con(R)/\equiv_R = Cp(Con(R))/\equiv_R = \mathcal{L}(R)$, thus, as expected by the uniqueness of the reticulation, R^* is isomorphic to $\mathcal{L}(R)$.

The fact that the variety of commutative unitary rings is semi-degenerate and congruence-modular and Corollary 7.7, along with the fact that the lattices $\operatorname{Con}(R)$ and $\operatorname{Id}(R)$ are isomorphic, give us:

Corollary 8.4. If R is a semiprime commutative unitary ring, then, for any $i, j \in \overline{1,5}$ and any nonzero cardinalities κ and μ , conditions $(iv)_{\mathrm{Id}(R)}$, $(i)_{\kappa,\mathrm{Id}(R)}$ and $(j)_{\mu,\mathrm{Id}(R)}$ are equivalent, in particular $\mathrm{Id}(R)$ is a Stone lattice iff it is a strongly Stone lattice.

Let us see that, similarly to what happens in residuated lattices, commutative unitary rings also fulfill an analogue of Davey's Theorem for elements instead of congruences.

Let κ be an arbitrary cardinality.

For any $a \in R$ and any $U \subseteq R$, $\operatorname{Ann}_R(a)$ and $\operatorname{Ann}_R(U)$ will denote the *annihilator* of a and that of U, respectively: $\operatorname{Ann}_R(a) = \{x \in R \mid xa = 0\}$ and $\operatorname{Ann}_R(U) = \bigcap_{u \in U} \operatorname{Ann}_R(u)$. As in the case of bounded lattices, let us denote by $\operatorname{Ann}(R) = \{\operatorname{Ann}_R(U) \mid U \subseteq R\}$, $\operatorname{Ann}_{<\infty}(R) = \{\operatorname{Ann}_R(U) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(U) \mid U \subseteq R, |U| \leq \kappa\}$, $\operatorname{PAnn}(R) = \{\operatorname{Ann}_R(a) \mid a \in R\} = \operatorname{Ann}_1(R)$, $\operatorname{2Ann}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \aleph_0\}$, $\operatorname{2Ann}_{\kappa}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(U)) \mid U \subseteq R, |U| < \kappa\}$ and $\operatorname{P2Ann}(R) = \{\operatorname{Ann}_R(\operatorname{Ann}_R(a)) \mid a \in R\} = \operatorname{2Ann}_1(R)$. It is well known and straightforward that $\operatorname{Ann}(R) \subseteq \operatorname{Id}(R)$.

E(R) will denote the set of the idempotent elements of R. Recall that $(E(R), \lor, \land = \cdot, \neg, 0, 1)$ is a Boolean algebra, where, for every $e, f \in E(R)$, $\neg e = 1 - e$ and $e \lor f = \neg (\neg e \land \neg f) = 1 - (1 - e) \cdot (1 - f)$.

R is called a *Baer ring* iff, for any $a \in R$, there exists an $e \in E(R)$ such that $\operatorname{Ann}_R(a) = eR$. By analogy to the case of bounded lattices, we shall call R a *strongly Baer ring*, respectively a κ -*Baer ring* iff, for any $U \subseteq R$, respectively any $U \subseteq R$ with $|U| \leq \kappa$, there exists an $e \in E(R)$ such that $\operatorname{Ann}_R(U) = eR$.

Let us consider the following conditions on R, where κ is an arbitrary cardinality:

- $(1^{\circ})_{\kappa,R}$ R is a κ -Baer ring;
- $(1^{\circ})_{<\infty,R}$ $\mathcal{A}nn_{<\infty}(R) \subseteq \{eR \mid e \in E(R)\};$
- $(1^{\circ})_R$ R is a strongly Baer ring;
- $(2^{\circ})_{\kappa,R}$ R is a Baer ring and E(R) is a κ -complete Boolean algebra;
- $(2^{\circ})_{<\infty,R}$ R is a Baer ring and E(R) is a Boolean algebra;
- $(2^{\circ})_R$ R is a Baer ring and E(R) is a complete Boolean algebra;

| $(3^{\circ})_{\kappa,R}$ | $2\operatorname{Ann}(R)$ is a κ -complete Boolean sublattice of $\operatorname{Id}(R)$ such that |
|---------------------------|---|
| | $I \mapsto \operatorname{Ann}_R(\operatorname{Ann}_R(I))$ is a lattice morphism from $\operatorname{Id}(R)$ to $2\operatorname{Ann}(R)$; |
| $(3^\circ)_{<\infty,R}$ | 2Ann(R) is a Boolean sublattice of $Id(R)$ such that |
| | $I \mapsto \operatorname{Ann}_R(\operatorname{Ann}_R(I))$ is a lattice morphism from $\operatorname{Id}(R)$ to $2\operatorname{Ann}(R)$; |
| $(3^{\circ})_R$ | 2Ann(R) is a complete Boolean sublattice of $Id(R)$ such that |
| | $I \mapsto \operatorname{Ann}_R(\operatorname{Ann}_R(I))$ is a lattice morphism from $\operatorname{Id}(R)$ to $2\operatorname{Ann}(R)$; |
| $(4^{\circ})_{\kappa,R}$ | for all $I, J \in Id(R)$, $Ann_R(I \cap J) = Ann_R(I) \vee Ann_R(J)$, and |
| | $2\operatorname{Ann}_{\kappa}(R) \subseteq \mathcal{A}nn_{<\infty}(R);$ |
| $(4^{\circ})_{<\infty,R}$ | for all $I, J \in Id(R)$, $Ann_R(I \cap J) = Ann_R(I) \vee Ann_R(J)$, and |
| | $2\operatorname{Ann}_{<\infty}(R) \subseteq \mathcal{A}nn_{<\infty}(R);$ |
| $(4^{\circ})_R$ | for all $I, J \in Id(R)$, $Ann_R(I \cap J) = Ann_R(I) \vee Ann_R(J)$, and |
| | $2\operatorname{Ann}(R) \subseteq \mathcal{A}nn_{<\infty}(R);$ |
| $(iv^{\circ})_R$ | for all $I, J \in Id(R)$, $Ann_R(I \cap J) = Ann_R(I) \vee Ann_R(J)$; |
| $(5^{\circ})_{\kappa,R}$ | for each $U \subseteq R$ with $ U \le \kappa$, $\operatorname{Ann}_R(U) \lor \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$; |
| $(5^\circ)_{<\infty,R}$ | for each finite $U \subseteq R$, $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$; |
| $(5^{\circ})_R$ | for each $U \subseteq R$, $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$. |

Obviously, conditions $(iv^{\circ})_R$, $(h^{\circ})_{\kappa,R}$, $(i^{\circ})_{<\infty,R}$ and $(j^{\circ})_R$ satisfy the properties stated after Remark 3.1 for conditions $(iv)_L$, $(h)_{\kappa,L}$, $(i)_{<\infty,L}$ and $(j)_L$, where $h, i, j \in \overline{1, 5}$.

Remark 8.5. If $n \in \mathbb{N}^*$, $u_1, \ldots, u_n \in R$ and, for each $i \in \overline{1, n}$, $\operatorname{Ann}_R(u_i) = e_i R$ for some $e_i \in E(R)$, then $\operatorname{Ann}_R(\{u_1, \ldots, u_n\}) = \bigcap_{i=1}^n \operatorname{Ann}_R(u_i) = \bigcap_{i=1}^n e_i R = e_1 R \cdots e_n R = e_1 \cdots e_n R$, with $e_1 \cdots e_n = e_1 \wedge \cdots \wedge e_n \in E(R)$, hence $(1^\circ)_{1,R}$ implies $(1^\circ)_{<\infty,R}$. Therefore $(1^\circ)_{1,R}$ is equivalent to $(1^\circ)_{<\infty,R}$, that is R is a Baer ring iff R satisfies $(1^\circ)_{<\infty,R}$.

Hence, if $\mathcal{A}nn(R) = \mathcal{A}nn_{<\infty}(R)$, so that $(1^{\circ})_{<\infty,R}$ is equivalent to $(1^{\circ})_R$, then $(1^{\circ})_{1,R}$ is equivalent to $(1^{\circ})_R$, that is R is a Baer ring iff R is a strongly Baer ring.

Remark 8.6. Proposition 6.7 and the fact that the lattices Con(R) and Id(R) are isomorphic ensure us that $\mathcal{B}(Id(R))$ is a Boolean sublattice of Id(R).

If R is semiprime, then, by [3, Lemma, p. 1863], the map $e \mapsto eR/\sim_R$ from E(R) to $\mathcal{B}(R^*)$ is a Boolean isomorphism, so, by Proposition 6.12, (iii), it follows that the map $e \mapsto eR$ from E(R) to $\mathcal{B}(\mathrm{Id}(R))$ is a Boolean isomorphism.

Lemma 8.7. If R is a semiprime commutative unitary ring, then:

- (i) $\mathcal{B}(\mathrm{Id}(R)) = \{eR \mid e \in E(R)\}$ and the map $e \mapsto eR$ from E(R) to $\mathcal{B}(\mathrm{Id}(R))$ is a Boolean isomorphism;
- (ii) if $U \subseteq R$, then $U \cap \operatorname{Ann}_R(U) \subseteq \{0\}$; if $I \in \operatorname{Id}(R)$, then $I \cap \operatorname{Ann}_R(I) = \{0\}$;
- (iii) if $U \subseteq R$ such that $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$, then $\operatorname{Ann}_R(U) = eR$ for some $e \in E(R)$.

Proof. (i) By Remark 8.6.

(ii) If $U \subseteq R$ and $x \in U \cap \operatorname{Ann}_R(U)$, then $x \cdot x = 0$, so that x = 0 since a semiprime commutative unitary ring has no nonzero nilpotents [11, p.125,126]. Now, if $I \in \operatorname{Id}(R)$, then $0 \in I \cap \operatorname{Ann}_R(I)$.

(iii) By (ii), it follows that $\operatorname{Ann}_R(U) \in \mathcal{B}(\operatorname{Id}(R))$, having $\operatorname{Ann}_R(\operatorname{Ann}_R(U))$ as a complement, so that $\operatorname{Ann}_R(U) = eR$ for some $e \in E(R)$ by (i).

Lemma 8.8. If R is a commutative unitary ring, then:

- for any $U \subseteq R$, $\operatorname{Ann}_R(U) = \operatorname{Ann}_R(\langle U \rangle_R)$;
- if all ideals of R are finitely generated, then $2\operatorname{Ann}(R) = 2\operatorname{Ann}_{<\infty}(R) \subseteq \operatorname{Ann}(R) = \operatorname{Ann}_{<\infty}(R);$
- for any $I \in Id(R)$, $Ann_{Id(R)}(I) = (Ann_R(I)]_{Id(R)}$ and $Ann_{Id(R)}(Ann_{Id(R)}(I))$ = $(Ann_R(Ann_R(I))]_{Id(R)}$.

Proof. Let $U \subseteq R$, arbitrary. Since $U \subseteq \langle U \rangle_R$, we have $\operatorname{Ann}_R(\langle U \rangle_R) \subseteq \operatorname{Ann}_R(U)$. The converse inclusion holds, as well, since, given any $a \in \langle U \rangle_R$ and any $x \in \operatorname{Ann}_R(U)$, we have $a = a_1 \cdot u_1 + \cdots + a_n \cdot u_n$ for some $n \in \mathbb{N}^*$, $a_1, \ldots, a_n \in R$ and $u_1, \ldots, u_n \in U$, so that $x \cdot u_1 = \cdots = x \cdot u_n = 0$, therefore $x \cdot a = 0$, so $x \in \operatorname{Ann}_R(\langle U \rangle_R)$.

Thus, in the particular case when all ideals of R are finitely generated, so that there exists a finite $F \subseteq R$ such that $\langle U \rangle_R = \langle F \rangle_R$, then $\operatorname{Ann}_R(U) = \operatorname{Ann}_R(\langle U \rangle_R) = \operatorname{Ann}_R(\langle F \rangle_R) = \operatorname{Ann}_R(F)$, hence $\operatorname{Ann}(R) = \operatorname{Ann}_{<\infty}(R)$.

Let $J \in \mathrm{Id}(R)$. Then: $J \in (\mathrm{Ann}_R(I)]_{\mathrm{Id}(R)}$ iff $J \subseteq \mathrm{Ann}_R(I)$ iff $x \in \mathrm{Ann}_R(I)$ for all $x \in J$ iff $x \cdot y = 0$ for all $x \in J$ and all $y \in I$ iff $J \cdot I = \{0\}$ iff $J \in \mathrm{Ann}_{\mathrm{Id}(R)}(I)$. Hence $\mathrm{Ann}_{\mathrm{Id}(R)}(I) = (\mathrm{Ann}_R(I)]_{\mathrm{Id}(R)}$, therefore $\mathrm{Ann}_{\mathrm{Id}(R)}(\mathrm{Ann}_{\mathrm{Id}(R)}(I)) = \mathrm{Ann}_{\mathrm{Id}(R)}((\mathrm{Ann}_R(I))) = (\mathrm{Ann}_R(I))_{\mathrm{Id}(R)}$.

Lemma 8.9. If R is a semiprime commutative unitary ring, then, for any $U \subseteq R$, there exists a finite subset $S \subseteq \langle U \rangle_R$ such that $\operatorname{Ann}_R(U) = \operatorname{Ann}_R(S)$, so $2\operatorname{Ann}(R) = 2\operatorname{Ann}_{<\infty}(R) \subseteq \operatorname{Ann}(R) = \operatorname{Ann}_{<\infty}(R)$.

Proof. By Remark 8.3 and Lemmas 8.8 and 4.2, for an appropriate finite subset $S \subseteq \langle U \rangle_R$, we have $\langle U \rangle_R / \sim_R = \langle S \rangle_R / \sim_R$, thus $\operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R) / \sim_R = \operatorname{Ann}_{R^*}(\langle U \rangle_R / \sim_R) = \operatorname{Ann}_{R^*}(\langle S \rangle_R / \sim_R) = \operatorname{Ann}_{\operatorname{Id}(R)}(\langle S \rangle_R) / \sim_R$, hence $(\operatorname{Ann}_R(U)]_{\operatorname{Id}(R)} = (\operatorname{Ann}_R(\langle U \rangle_R)]_{\operatorname{Id}(R)} = \operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R) = \operatorname{Ann}_{\operatorname{Id}(R)}(\langle S \rangle_R) = (\operatorname{Ann}_R(\langle S \rangle_R)]_{\operatorname{Id}(R)} = (\operatorname{Ann}_R(S)]_{\operatorname{Id}(R)}$, thus $\operatorname{Ann}_R(U) = \operatorname{Ann}_R(S)$.

Proposition 8.10. Let R be a commutative unitary ring.

- (i) If all ideals of R are finitely generated, then R is a Baer ring iff R is a strongly Baer ring.
- (ii) If R is semiprime, then: R is a Baer ring iff R is a strongly Baer ring iff Id(R) is a Stone lattice iff Id(R) is a strongly Stone lattice.

Proof. (i) By Remark 8.5 and Lemma 8.8.

(ii) By Remark 8.5 and Lemma 8.9, R is Baer iff R is strongly Baer.

For any $U \subseteq R$, we have $(\operatorname{Ann}_R(U)]_{\operatorname{Id}(R)} = (\operatorname{Ann}_R(\langle U \rangle_R)]_{\operatorname{Id}(R)} = \operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R)$ by Lemma 8.8, so that, for any $e \in R$, $\operatorname{Ann}_R(U) = eR$ iff $\operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R) = (eR]_{\operatorname{Id}(R)}$. According to Lemma 8.7, (i), $e \in E(R)$ iff $eR \in \mathcal{B}(\operatorname{Id}(R))$. Hence $(1)_{1,\operatorname{Id}(R)}$ is equivalent to $(1^\circ)_R$, that is $\operatorname{Id}(R)$ is a Stone lattice iff R is a strongly Baer ring.

Finally, by Corollary 8.4, Id(R) is a Stone lattice iff Id(R) is a strongly Stone lattice.

See also [3, Theorem 8] and [19, Theorem 2.6], according to which, if R is semiprime, then R is a Baer ring iff R^* is a Stone lattice, which, by Corollary 7.6 and the fact that the lattices Con(R) and Id(R) are isomorphic, is equivalent to Id(R) being a Stone lattice.

Proposition 8.11. For any semiprime commutative unitary ring R and any cardinality κ , conditions $(2^{\circ})_{\kappa,R}$ and $(2)_{\kappa,\mathrm{Id}(R)}$ are equivalent.

Proof. By Lemma 8.7, (i), and Proposition 8.10, (ii).

For the next lemma, recall that $2\operatorname{Ann}(R) \subseteq \operatorname{Ann}(R) \subseteq \operatorname{Id}(R)$, $\operatorname{PAnn}(\operatorname{Id}(R)) \subseteq \operatorname{Id}(\operatorname{Id}(R))$ and $\operatorname{P2Ann}(\operatorname{Id}(R)) \subseteq \operatorname{Id}(\operatorname{Id}(R))$, and we will be referring to these sets of annihilators as subposets of $\operatorname{Id}(R)$, respectively $\operatorname{Id}(\operatorname{Id}(R))$.

Lemma 8.12. For any commutative unitary ring R, the map $x \mapsto (x]_{\mathrm{Id}(R)}$ from $\mathcal{A}nn(R)$ to $\mathrm{PAnn}(\mathrm{Id}(R))$, as well as from $2\mathrm{Ann}(R)$ to $\mathrm{P2Ann}(\mathrm{Id}(R))$, is an order isomorphism.

Proof. By Lemma 8.8, $Ann(R) = \{Ann_R(I) \mid I \in Id(R)\}$ and $2Ann(R) = \{Ann_R(Ann_R(I)) \mid I \in Id(R)\}$, hence these maps are completely defined. By the same lemma, these maps are well defined and surjective. Clearly, they are injective, thus bijective, and both these maps and their inverses are order-preserving.

Proposition 8.13. For any commutative unitary ring R and any cardinality κ , the properties $(3^{\circ})_{\kappa,R}$ and $(3)_{\kappa,\mathrm{Id}(R)}$ are equivalent.

Proof. By Lemma 8.12 and the fact that, by Lemma 8.8, the map $x \mapsto (x]_{\mathrm{Id}(R)}$ from $2\mathrm{Ann}(R) \subseteq \mathrm{Id}(R)$ to $\mathrm{P2Ann}(\mathrm{Id}(R))$ composed with the map $I \mapsto \mathrm{Ann}_R$ $(\mathrm{Ann}_R(I))$ from $\mathrm{Id}(R)$ to $2\mathrm{Ann}(R)$ equals the map $I \mapsto \mathrm{Ann}_{\mathrm{Id}(R)}(\mathrm{Ann}_{\mathrm{Id}(R)}(I))$ from $\mathrm{Id}(R)$ to $\mathrm{P2Ann}(\mathrm{Id}(R)) \subseteq \mathrm{Id}(\mathrm{Id}(R))$.

Proposition 8.14. Let R be a commutative unitary ring. Then:

(i) $(iv^{\circ})_R$ is equivalent to $(iv)_{\mathrm{Id}(R)}$;

- (ii) if R has all ideals finitely generated, then conditions (iv)_{Id(R)}, (iv°)_R and (4°)_R are equivalent;
- (iii) if R is semiprime, then conditions $(iv)_{Id(R)}$, $(4)_{Id(R)}$, $(iv^{\circ})_R$ and $(4^{\circ})_R$ are equivalent.

Proof. (i) Let $I, J \in Id(R)$. By Lemma 8.8, $\operatorname{Ann}_{Id(R)}(I \cap J) = (\operatorname{Ann}_R(I \cap J)]_{Id(R)}$ and $\operatorname{Ann}_{Id(R)}(I) \vee \operatorname{Ann}_{Id(R)}(J) = (\operatorname{Ann}_R(I)]_{Id(R)} \vee (\operatorname{Ann}_R(J)]_{Id(R)} = (\operatorname{Ann}_R(I) \vee \operatorname{Ann}_R(J)]_{Id(R)}$, hence: $\operatorname{Ann}_R(I \cap J) = \operatorname{Ann}_R(I) \vee \operatorname{Ann}_R(J)$ iff $\operatorname{Ann}_{Id(R)}(I \cap J) = \operatorname{Ann}_{Id(R)}(I) \vee \operatorname{Ann}_{Id(R)}(J)$.

(ii), (iii) By Lemmas 8.8 and 8.9, if R has all ideals principal or it is semiprime, then the second part of condition $(4^{\circ})_R$ is trivially satisfied, so that $(4^{\circ})_R$ is equivalent to $(iv^{\circ})_R$.

By (i), $(iv^{\circ})_R$ is equivalent to $(iv)_{\mathrm{Id}(R)}$.

Finally, by Corollary 8.4, if R is semiprime, then $(iv)_{\mathrm{Id}(R)}$ is equivalent to $(4)_{\mathrm{Id}(R)}$.

Proposition 8.15. Let R be a commutative unitary ring. Then:

- $(5)_{1,\mathrm{Id}(R)}$ is equivalent to $(5^{\circ})_R$;
- if all ideals of R are finitely generated, then $(5)_{1,\mathrm{Id}(R)}$, $(5^{\circ})_{R}$ and $(5^{\circ})_{<\infty,R}$ are equivalent;
- if R is semiprime, then $(5)_{1,\mathrm{Id}(R)}$, $(5)_{\mathrm{Id}(R)}$, $(5^{\circ})_R$ and $(5^{\circ})_{<\infty,R}$ are equivalent.

Proof. By Lemma 8.8, for any $U \subseteq R$, we have $\operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R) \vee \operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R)) = (\operatorname{Ann}_R(\langle U \rangle_R)]_{\operatorname{Id}(R)} \vee (\operatorname{Ann}_R(\operatorname{Ann}_R(\langle U \rangle_R))]_{\operatorname{Id}(R)} = (\operatorname{Ann}_R(U))_{\operatorname{Id}(R)} \vee (\operatorname{Ann}_R(\operatorname{Ann}_R(U))]_{\operatorname{Id}(R)} = (\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U))]_{\operatorname{Id}(R)} \in \operatorname{PId}(\operatorname{Id}(R))$ since $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U) \in \operatorname{Id}(R),$ hence $\operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R) \vee \operatorname{Ann}_{\operatorname{Id}(R)}(\langle U \rangle_R)) = \operatorname{Id}(R) = (R]_{\operatorname{Id}(R)}$ iff $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$. Therefore $(5)_{1,\operatorname{Id}(R)}$ is equivalent to $(5^\circ)_R$.

Clearly, if $\mathcal{A}nn(R) = \mathcal{A}nn_{<\infty}(R)$, in particular if R has all ideals principal or R is semiprime, then $(5^{\circ})_R$ is equivalent to $(5^{\circ})_{<\infty,R}$.

By Corollary 8.4, if R is semiprime, then $(5)_{1,\mathrm{Id}(R)}$ is equivalent to $(5)_{\mathrm{Id}(R)}$.

Theorem 8.16. If R is a semiprime commutative unitary ring, then, for any nonzero cardinality κ and any $h, i, j \in \overline{1,5}$, $(iv^{\circ})_R$, $(h^{\circ})_{\kappa,R}$, $(i^{\circ})_{<\infty,R}$ and $(j^{\circ})_R$ are equivalent.

Proof. By Corollary 8.4 and Propositions 8.10, 8.11, 8.13, 8.14 and 8.15.

Remark 8.17. Let S be a commutative unitary ring. Since the variety of commutative unitary rings is semi-degenerate and thus it has no skew congruences, it follows that $Id(R \times S) = Id(R) \times Id(S)$, hence, if R and S are semiprime, then, for any cardinality κ , the ring $R \times S$ is κ -Baer iff R and S are κ -Baer, according to Corollary 7.10.

If we eliminate from Theorem 8.16 the trivial implications, along with those that immediately follow from Lemma 8.9, then we obtain the following.

Corollary 8.18. If R is a semiprime commutative unitary ring, then the following are equivalent:

- R is a Baer ring;
- R is a strongly Baer ring and E(R) is a complete Boolean algebra;
- $2\operatorname{Ann}(R)$ is a Boolean sublattice of $\operatorname{Id}(R)$ such that $I \mapsto \operatorname{Ann}_R(\operatorname{Ann}_R(I))$ is a lattice morphism from $\operatorname{Id}(R)$ to $2\operatorname{Ann}(R)$;
- $2\operatorname{Ann}(R)$ is a complete Boolean sublattice of $\operatorname{Id}(R)$ such that $I \mapsto \operatorname{Ann}_R(I)$ is a lattice morphism from $\operatorname{Id}(R)$ to $2\operatorname{Ann}(R)$;
- for all $I, J \in \mathrm{Id}(R)$, $\mathrm{Ann}_R(I \cap J) = \mathrm{Ann}_R(I) \vee \mathrm{Ann}_R(J)$;
- for any $U \subseteq R$, $\operatorname{Ann}_R(U) \vee \operatorname{Ann}_R(\operatorname{Ann}_R(U)) = R$.

Propositions 8.10, 8.11, 8.13, 8.14 and 8.15, along with Theorem 3.3, (iii), also give us:

Corollary 8.19. Let R be a commutative unitary ring and m be a nonzero cardinality such that the intersection in Id(R) is distributive w.r.t. the joins of families of cardinality at most m. Then, for any nonzero cardinalities $\kappa \leq m$, $\lambda \leq m$ and $\mu \leq m$ and any infinite cardinality $\iota \leq m$:

- conditions $(2^{\circ})_{\kappa,R}$, $(3^{\circ})_{\lambda,R}$ and $(5^{\circ})_R$ are equivalent;
- if all ideals of R are finitely generated, then conditions $(2^{\circ})_{\kappa,R}$, $(3^{\circ})_{\lambda,R}$, $(iv^{\circ})_{R}$, $(4^{\circ})_{\mu,R}$, $(4^{\circ})_{R}$, $(5^{\circ})_{<\infty,R}$, $(5^{\circ})_{\iota,R}$ and $(5^{\circ})_{R}$ are equivalent.

9. Conclusions

Determining what kinds of complete algebraic modular lattices are congruence lattices of semiprime algebras from semi-degenerate congruence-modular varieties may be of interest, since it will follow that the equivalences in Corollary 7.7 hold for all those kinds of lattices.

Another theme for future research is studying further extensions of Davey's Theorem to different kinds of lattices, as well as finding more classes of algebras in which, given an appropriate setting (regarding definitions for annihilators and a Boolean center), Davey's Theorem holds not only for congruences, but also for elements, as in the case of bounded distributive lattices, commutator lattices, residuated lattices and commutative unitary rings.

94

Acknowledgements

This work was supported by the research grant "Clonoids: a Unifying Approach to Equational Logic and Clones" of the Austrian Science Fund FWF (P29931).

I thank Erhard Aichinger for very helpful discussions on the problems tackled in this paper. INTO: I thank Erhard Aichinger for proposing some of the previously open problems tackled in this work and George Georgescu for suggesting the study on rings in Section 8 of this paper.

References

- P. Agliano, Prime spectra in modular varieties, Algebra Univ. 30 (1993) 581–597. https://doi.org/10.1007/BF01195383
- [2] E. Aichinger, Congruence lattices forcing nilpotency, J. Algebra and Its Appl. 17 (2) (2018). https://doi.org/10.1142/S0219498818500330
- [3] L.P. Belluce, Spectral spaces and non-commutative rings, Commun. in Algebra 19 (7) (1991) 1855–1865. https://doi.org/10.1080/00927879108824234
- [4] L.P. Belluce, Spectral closure for non-commutative rings, Commun. in Algebra 25 (5) (1997) 1513–1536. https://doi.org/10.1080/00927879708825933
- J. Czelakowski, Additivity of the commutator and residuation, Rep. Math. Logic 43 (2008) 109–132. https://doi.org/10.1007/978-3-319-21200-5
- [6] B.A. Davey, *m-stone lattices*, Canad. J. Math. 24 (6) (1972) 1027–1032. https://doi.org/10.4153/CJM-1972-104-x
- [7] R. Freese and R. McKenzie, Commutator Theory for Congruence-modular Varieties, London Mathematical Society Lecture Note Series 125 (Cambridge University Press, 1987). https://doi.org/10.1112/blms/20.4.362
- [8] N. Galatos, P. Jipsen, T. Kowalski and H. Ono, Residuated Lattices: An Algebraic Glimpse at Substructural Logics, Studies in Logic and The Foundations of Mathematics 151 (Elsevier, Amsterdam/ Boston /Heidelberg /London /New York /Oxford /Paris /San Diego/ San Francisco /Singapore /Sydney /Tokyo, 2007). https://doi.org/10.1016/s0049-237x(07)x8002-9
- G. Georgescu and C. Mureşan, Congruence Boolean lifting property, J. Multiple-Valued Logic and Soft Comp. 29 (3–4) (2017) 225–274.
- [10] G. Georgescu and C. Mureşan, The reticulation of a universal algebra, Sci. Ann. Comp. Sci. XXVIII (2018) 67–113. https://doi.org/10.7561/SACS.2018.1.67

- J.E. Kist, Two characterizations of commutative Baer rings, Pacific J. Math. 50 (1974). https://doi.org/10.2140/pjm.1974.50.125
- [12] A. Iorgulescu, Algebras of Logic as BCK Algebras (Editura ASE, Bucharest, 2008).
- [13] A. Joyal, Le théorème de Chevalley-Tarski et Remarques sur l'algèbre constructive, Cahiers Topol. Géom. Différ. 16 (1975) 256–258.
- [14] C. Mureşan, Algebras of Many-valued Logic. Contributions to the Theory of Residuated Lattices, Ph.D. Thesis, 2009.
- C. Mureşan, Co-stone residuated lattices, Annals of the University of Craiova, Math. Comp. Sci. Ser. 40 (2013) 52–75. https://doi.org/10.7561/SACS.2018.1.67
- [16] P. Ouwehand, Commutator Theory and Abelian Algebras. arXiv:1309.0662 [math.RA].
- [17] D. Piciu, Algebras of Fuzzy Logic (Editura Universitaria Craiova, Craiova, 2007).
- [18] D. Schweigert, Tolerances and commutators on lattices, Bull. Austral. Math. Soc. 37 (2) (1988) 213–219. https://doi.org/10.1017/S0004972700026745
- [19] H. Simmons, *Reticulated rings*, J. Algebra 66 (1980) 169–192. https://doi.org/10.1016/0021-8693(80)90118-0

Received 8 April 2019 Revised 29 March 2020 Accepted 30 November 2020