

RADICALS OF GENERALIZED PRIME IDEALS IN TERNARY SEMIGROUPS

NARAKON SATVONG¹, THAWHAT CHANGPHAS²

AND

PANUWAT LUANGCHAISRI³

^{1,2,3}*Department of Mathematics*
Khon Kaen University, Khon Kaen 40002, Thailand

²*Centre of Excellence in Mathematics*
CHE, Si Ayuttaya Rd., Bangkok 10400, Thailand

e-mail: satvong1994@gmail.com
thacha@kku.ac.th
desparadoskku@hotmail.com

Abstract

In this paper, the concepts of f -prime ideals and f -semiprime ideals on a ternary semigroup are considered as a generalization of pseudo prime ideals and pseudo semiprime ideals, respectively. Then such ideals introduced are used to describe left (respectively, right) f -primary ideals on a ternary semigroup.

Keywords: ternary semigroup, pseudo prime ideal, good mapping, f -prime ideal, left f -primary decomposition.

2010 Mathematics Subject Classification: 20N99, 20M12, 03E75.

1. INTRODUCTION

A *ternary semigroup* is a nonempty set T together with a ternary operation $[] : T \times T \times T \rightarrow T$ written as $(a, b, c) \rightarrow [abc]$ satisfying the associative law of the first kind:

$$[[abc]uv] = [a[bcu]v] = [ab[cuv]]$$

for all $a, b, c, u, v \in T$. Yet prior, the structures was contemplated by Kasner [9] who give the possibility of n -ary algebras. In [6], Hewitt and Zuckerman studied

the concept of ternary semigroups and showed that every semigroup can be considered as a ternary semigroup but there is an example of a ternary semigroup which does not reduce to a semigroup. Recently, ternary semigroups have been widely studied (see, [1, 2, 3, 4, 11, 13]).

Let $(T, [\])$ be a ternary semigroup. For nonempty subsets A, B, C of T , the set product $[ABC]$ is defined by

$$[ABC] = \{[abc] \mid a \in A, b \in B, c \in C\}.$$

If $A = \{a\}$, we write $[ABC]$ as $[aBC]$. The cases $[A\{b\}C]$ and $[AB\{c\}]$ can be written similarly.

For convenience, we shall write T for a ternary semigroup $(T, [\])$, and write ABC for the product $[ABC]$.

Let T be a ternary semigroup. A nonempty subset A of T is said to be

- (1) a *left ideal* of T if $TTA \subseteq A$;
- (2) a *right ideal* of T if $ATT \subseteq A$;
- (3) a *lateral ideal* (or *middle ideal*) of T if $TAT \subseteq A$;
- (4) a *two-sided ideal* of T if it is both a left and a right ideal of T ;
- (5) an *ideal* of T if it is a left, a right and a lateral ideal of T .

For any $a \in T$, it is clear that the intersection of all ideals of T containing a is an ideal of T . It is called the *principal ideal* of T generated by a and denoted by (a) .

There are many authors considered m -systems and n -systems on semigroups being related to pseudo prime ideals and pseudo semiprime ideals of the semigroups. By amplifying these concepts, Rao and Sarala [10] introduced an m -system and an n -system on ternary semigroups and then presented properties of pseudo prime ideals and pseudo semiprime ideals on ternary semigroups by these systems. The following definitions we refer to [10] (see also [7]).

Definition. Let T be a ternary semigroup and P a proper ideal of T . Then P is said to be a *pseudo prime ideal* of T if for any ideals A, B, C of T ,

$$ABC \subseteq P \text{ implies } A \subseteq P, B \subseteq P \text{ or } C \subseteq P.$$

Lemma 1 [7]. *Let P be a proper ideal of a ternary semigroup T . Then P is pseudo prime if and only if for any $a, b, c \in T$, $aTbTc \subseteq P$, $aTTbTTc \subseteq P$, $TaTbTTc \subseteq P$ and $aTTbTcT \subseteq P$ imply $a \in P, b \in P$ or $c \in P$.*

Definition. Let T be a ternary semigroup and P a proper ideal of T . Then P is said to be a *pseudo semiprime ideal* of T if for any ideal A of T ,

$$AAA \subseteq P \text{ implies } A \subseteq P.$$

Remark 2. Let P be a proper ideal of a ternary semigroup T . Then P is pseudo semiprime if and only if for any $a \in T$, $aTaTa \subseteq P$, $aTTaTTa \subseteq P$, $TaTaTTa \subseteq P$ and $aTTaTaT \subseteq P$ imply $a \in P$.

Definition. Let T be a ternary semigroup. Then a subset A of T is called an m -system of T if for any $a, b, c \in A$,

$$(aTbTc \cup aTTbTTc \cup TaTbTTc \cup aTTbTcT) \cap A \neq \emptyset.$$

Definition. Let T be a ternary semigroup. Then a subset A of T is called an n -system of T if for any $a \in A$,

$$(aTaTa \cup aTTaTTa \cup TaTaTTa \cup aTTaTaT) \cap A \neq \emptyset.$$

2. f -PRIME IDEALS AND f -PRIME RADICALS

The notion of f -prime ideals on a ring was introduced to be a generalization of prime ideals by Murata, Kurata and Marubayashi [8]. Then such concept was extended to semirings by Sardar and Goswami [12]. After that, it appear on ordered algebraic structures having one operation, called ordered semigroups, by Gu [5]. In this section, we extend this concept to ternary semigroups.

Definition. Let $A(T)$ denote the set of all ideals of a ternary semigroup T . A mapping $f : T \rightarrow A(T)$ is called a *good mapping* on T if

- (i) $\forall a \in T, a \in f(a)$;
- (ii) $\forall a \in T \forall A \in A(T), x \in f(a) \cup A \Rightarrow f(x) \subseteq f(a) \cup A$.

Definition. Let f be a good mapping on a ternary semigroup T . A subset F of T is called an f -system of T if F contains an m -system F^* of T such that

$$\forall t \in F, f(t) \cap F^* \neq \emptyset.$$

Here, F^* is called the *kernel* of F .

Remark 3. Every m -system of a ternary semigroup T is an f -system of T with kernel itself.

Definition. Let T be a ternary semigroup and f a good mapping on T . An ideal A of T is called an f -prime ideal of T if its complement $C(A)$ is an f -system.

Remark 4. Let T be a ternary semigroup. Then the following statements hold:

- (1) if P is a pseudo prime ideal of T , then $C(P)$ is an m -system of T ;
- (2) if P is a pseudo prime ideal of T , then P is an f -prime ideal of T .

Proof. (1) Assume that P is a pseudo prime ideal of T and let $a, b, c \in C(P)$. Since P is pseudo prime,

$$aTbTc \not\subseteq P, aTTbTTc \not\subseteq P, TaTbTTc \not\subseteq P \text{ or } aTTbTcT \not\subseteq P.$$

Thus

$$(aTbTc \cup aTTbTTc \cup TaTbTTc \cup aTTbTcT) \cap C(P) \neq \emptyset.$$

Therefore $C(P)$ is an m -system of T .

(2) It is obtained directly from (1) and Remark 3. \blacksquare

Lemma 5. Let P be an f -prime ideal of a ternary semigroup T . For any $a, b, c \in T$, if

$$f(a)Tf(b)Tf(c) \subseteq P, Tf(a)Tf(b)TTf(c) \subseteq P \text{ and } f(a)TTf(b)Tf(c)T \subseteq P,$$

then $a \in P, b \in P$ or $c \in P$.

Proof. Let $a, b, c \in T$. Suppose contrary that $a, b, c \in C(P)$. Since $C(P)$ is an f -system of T ,

$$f(a) \cap (C(P))^* \neq \emptyset, f(b) \cap (C(P))^* \neq \emptyset \text{ and } f(c) \cap (C(P))^* \neq \emptyset.$$

Let $x_1 \in f(a) \cap (C(P))^*, x_2 \in f(b) \cap (C(P))^*$ and $x_3 \in f(c) \cap (C(P))^*$. Since $(C(P))^*$ is an m -system of T ,

$$(x_1Tx_2Tx_3 \cup x_1TTx_2TTx_3 \cup Tx_1Tx_2TTx_3 \cup x_1TTx_2Tx_3T) \cap (C(P))^* \neq \emptyset.$$

Consequently,

$$x_1Tx_2Tx_3 \not\subseteq P, x_1TTx_2TTx_3 \not\subseteq P, Tx_1Tx_2TTx_3 \not\subseteq P, \text{ or } x_1TTx_2Tx_3T \not\subseteq P. \blacksquare$$

Definition. Let T be a ternary semigroup and

$$\begin{aligned} fS(T) &= \text{the set of all } f\text{-systems of } T; \\ fPI(T) &= \text{the set of all } f\text{-prime ideals of } T. \end{aligned}$$

For any ideal A of T , the f -prime radicals of A is defined by

$$r_f(A) = \{a \in T \mid \forall F \in fS(T), a \in F \Rightarrow F \cap A \neq \emptyset\}.$$

Theorem 6. Let T be a ternary semigroup and A be an ideal of T . Then

$$r_f(A) = \bigcap \{P \mid A \subseteq P \in fPI(T)\}.$$

Proof. Suppose that $x \notin \bigcap \{P \mid A \subseteq P \in fPI(T)\}$. Then $x \notin P$ for some $P \in fPI(T)$ with $A \subseteq P$. Since $C(P)$ is an f -system of T and $C(P) \cap A = \emptyset$,

then $x \notin r_f(A)$. Hence $r_f(A) \subseteq \bigcap \{P \mid A \subseteq P \in fPI(T)\}$. On the other hand, suppose that $x \notin r_f(A)$. Then there exists $F \in fS(T)$ such that $x \in F$ and $F \cap A = \emptyset$. Thus $C(F)$ is an f -prime ideal of T containing A . Then $x \notin C(F)$ implies $x \notin \bigcap \{P \mid A \subseteq P \in fPI(T)\}$. Therefore

$$\bigcap \{P \mid A \subseteq P \in fPI(T)\} \subseteq r_f(A). \quad \blacksquare$$

3. f -SEMIPRIME IDEALS

Definition. Let T be a ternary semigroup. A subset E of T is said to be an f *n*-system of T if

$$E = \bigcup_{i \in \Gamma} F_i$$

where $\{F_i \mid i \in \Gamma\} \subseteq fS(T)$.

Definition. An ideal P of a ternary semigroup T is said to be an f -semiprime ideal of T if its complement $C(P)$ in T is an f *n*-system of T .

Remark 7. Let T be a ternary semigroup. Then the following statements hold:

- (1) every f -system of T is an f *n*-system of T ;
- (2) every f -prime ideal of T is an f -semiprime ideal of T .

The following example shows that f -prime need not be pseudo prime and f -semiprime need not be pseudo semiprime.

Example 8. Consider the ternary semigroup $T = \{a, b, c\}$ defined by:

a	a	b	c	b	a	b	c	c	a	b	c
a	a	a	a	a	a	a	a	a	a	a	a
b	a	a	a	b	a	a	a	b	a	a	b
c	a	a	a	c	a	a	b	c	a	b	c

The ideals of T are $\{a\}$, $\{a, b\}$ and T .

Define $f(a) = \{a\}$, $f(b) = f(c) = T$; then f is a good mapping. Let $A = \{a\}$. Then $C(A) = \{b, c\}$ is an f -system with kernel $F^* = \{c\}$. Hence, A is f -prime and f -semiprime. However, A is not pseudo prime and not pseudo semiprime. Indeed $\{a, b\}\{a, b\}\{a, b\} \subseteq A$, but $\{a, b\} \not\subseteq A$.

Lemma 9. Let T be a ternary semigroup. Then the following statements hold:

- (1) if P is a pseudo semiprime ideal of T , then $C(P)$ is an n -system of T ;
- (2) if N is an n -system of T , then N is the union of some m -systems of T .

Proof. (1) Assume that P is a pseudo semiprime ideal of T . Let $a \in C(P)$. Then $aTaTa \not\subseteq P$, $aTTaTTa \not\subseteq P$, $TaTaTTa \not\subseteq P$ or $aTTaTaT \not\subseteq P$. Thus,

$$(aTaTa \cup aTTaTTa \cup TaTaTTa \cup aTTaTaT) \cap C(P) \neq \emptyset.$$

(2) Assume that N is an n -system of T . Let $a \in N$. Then

$$(aTaTa \cup aTTaTTa \cup TaTaTTa \cup aTTaTaT) \cap N \neq \emptyset.$$

Given

$$a_1 \in (aTaTa \cup aTTaTTa \cup TaTaTTa \cup aTTaTaT) \cap N.$$

We have that

$$(a_1Ta_1Ta_1 \cup a_1TTa_1TTa_1 \cup Ta_1Ta_1TTa_1 \cup a_1TTa_1Ta_1T) \cap N \neq \emptyset.$$

Then there exists

$$a_2 \in (a_1Ta_1Ta_1 \cup a_1TTa_1TTa_1 \cup Ta_1Ta_1TTa_1 \cup a_1TTa_1Ta_1T) \cap N.$$

Continue in the same manner, we obtain the set

$$M_a = \{a_0 = a, a_1, a_2, \dots\}.$$

Next, we will show that M_a is an m -system of T . Let $a_i, a_j, a_k \in M_a$.

If $i = \max\{i, j, k\}$, we have

$$\begin{aligned} a_{i+1} &\in a_iTa_iTa_i \cup a_iTTa_iTTa_i \cup Ta_iTa_iTTa_i \cup a_iTTa_iTa_iT \\ &\subseteq a_iTa_{i-1}Ta_{i-1} \cup a_iTTa_{i-1}TTa_{i-1} \cup Ta_iTa_{i-1}TTa_{i-1} \cup a_iTTa_{i-1}Ta_{i-1}T \\ &\vdots \\ &\subseteq a_iTa_jTa_k \cup a_iTTa_jTTa_k \cup Ta_iTa_jTTa_k \cup a_iTTa_jTa_kT. \end{aligned}$$

If $j = \max\{i, j, k\}$ or $k = \max\{i, j, k\}$, we obtain that

$$a_{j+1} \in a_iTa_jTa_k \cup a_iTTa_jTTa_k \cup Ta_iTa_jTTa_k \cup a_iTTa_jTa_kT$$

and

$$a_{k+1} \in a_iTa_jTa_k \cup a_iTTa_jTTa_k \cup Ta_iTa_jTTa_k \cup a_iTTa_jTa_kT,$$

respectively. Thus

$$(a_iTa_jTa_k \cup a_iTTa_jTTa_k \cup Ta_iTa_jTTa_k \cup a_iTTa_jTa_kT) \cap M_a \neq \emptyset.$$

Therefore M_a is an m -system of T . Hence

$$N = \bigcup_{a \in N} M_a$$

is the union of m -systems of T . ■

Lemma 10. *Let T be a ternary semigroup. If A is a pseudo semiprime ideal of T , then A is an f -semiprime ideal of T .*

Proof. Assume that A is a pseudo semiprime ideal of T . By Lemma 9(1) and (2), $C(A)$ is the union of some m -systems of T . By Remark 3, $C(A)$ is the union of some f -systems of T . Hence $C(A)$ is an fn -system of T . Therefore A is an f -semiprime ideal of T . ■

Lemma 11. *Let P be an f -semiprime ideal of a ternary semigroup T and $a \in T$. If $f(a)Tf(a)Tf(a) \subseteq P$, $Tf(a)Tf(a)TTf(a) \subseteq P$ and $f(a)TTf(a)Tf(a)T \subseteq P$, then $a \in P$.*

Proof. Given

$$C(P) = \bigcup_{i \in \Gamma} F_i,$$

where $\{F_i \mid i \in \Gamma\} \subseteq fS(T)$. Let $a \in C(P)$. Then $a \in F_i$ for some $i \in \Gamma$. Since F_i is an f -system of T , it follows that $f(a) \cap F_i^* \neq \emptyset$. Let $x \in f(a) \cap F_i^*$. Since F_i^* is an m -system of T , we obtain

$$\emptyset \neq (xTxTx \cup xTTxTTx \cup TxTxTTx \cup xTTxTxT) \cap F_i^* \subseteq F_i^* \subseteq F_i \subseteq C(P).$$

These imply that $xTxTx \not\subseteq P, xTTxTTx \not\subseteq P, TxTxTTx \not\subseteq P$ or $xTTxTxT \not\subseteq P$. This proof is complete. ■

Lemma 12. *Let P be an f -semiprime ideal of a ternary semigroup T . Then $r_f(P) = P$.*

Proof. It is clear that $P \subseteq r_f(P)$. We will show that $r_f(P) \subseteq P$, by proving that $C(P) \subseteq C(r_f(P))$. Let $x \in C(P)$. Since $C(P)$ is an fn -system of T ,

$$C(P) = \bigcup_{i \in \Gamma} F_i,$$

where $\{F_i \mid i \in \Gamma\} \subseteq fS(T)$. This implies $x \in F_i$ for some $i \in \Gamma$. Since $x \notin P$ and F_i is an f -system of T such that $F_i \cap P = \emptyset$, then $x \notin r_f(P)$. Thus $x \in C(r_f(P))$. The proof is complete. ■

Theorem 13. *Let A be an ideal of a ternary semigroup T . Then $r_f(A)$ is the smallest f -semiprime ideal of T containing A .*

Proof. It is clear that $A \subseteq r_f(A)$. By Theorem 6,

$$r_f(A) = \bigcap \{P \mid A \subseteq P \in fPI(T)\}.$$

Then

$$C(r_f(A)) = \bigcup \{C(P) \mid A \subseteq P \in fPI(T)\}.$$

Thus $C(r_f(A))$ is an fn -system of T . Therefore $r_f(A)$ is an f -semiprime ideal of T . Let P be an f -semiprime ideal of T containing A . Then

$$r_f(A) \subseteq r_f(P) = P.$$

Thus $r_f(A)$ is the least f -semiprime ideal of T containing A . ■

Definition. Let f be a good mapping of a ternary semigroup T . For $a \in T$ and $A \in A(T)$, define the set $A : a$ to be

$$A : a := \{x \in T \mid f(a)Tf(a)Tf(x) \subseteq A, Tf(a)Tf(a)TTf(x) \text{ and} \\ f(a)TTf(a)Tf(x)T \subseteq A\}.$$

This set is called the *left f -quotient* of A by a . Moreover, for any $B \in A(T)$, the *left f -quotient* of A by B is defined by

$$A : B = \bigcap_{b \in B} (A : b).$$

We note that $A : a$ may be empty. See the following example.

Example 14. We consider the ternary semigroup T of Example 8. If we define $f(a) = f(b) = f(c) = T$, then the mapping f is a good mapping. Let $A = \{a\}$. Then $A : a, A : b$ and $A : c$ are all empty.

Lemma 15. Let f be a good mapping on a ternary semigroup T . Let $A, A_1, A_2, B, B_1, B_2 \in A(T)$ and $a \in T$.

- (1) $A_1 \subseteq A_2 \Rightarrow A_1 : a \subseteq A_2 : a$.
- (2) $A_1 \subseteq A_2 \Rightarrow A_1 : B \subseteq A_2 : B$.
- (3) $B_1 \subseteq B_2 \Rightarrow A : B_1 \supseteq A : B_2$.
- (4) $(A_1 \cap A_2) : a = (A_1 : a) \cap (A_2 : a)$.
- (5) $(A_1 \cap A_2) : B = (A_1 : B) \cap (A_2 : B)$.

Lemma 16. Let f be a good mapping on a ternary semigroup T . For any $A \in A(T)$ and $a \in T$, $A : a$ is either empty or an ideal of T containing A .

Proof. Let A be an ideal of T and $a \in T$. Suppose that $A : a \neq \emptyset$. We will show that $A : a$ is an ideal of T . Let $x \in A : a$ and $r_1, r_2 \in T$. Then the following inclusions hold:

$$\begin{aligned} f(a)Tf(a)Tf(xr_1r_2) &\subseteq f(a)Tf(a)Tf(x) \subseteq A; \\ Tf(a)Tf(a)TTf(xr_1r_2) &\subseteq Tf(a)Tf(a)TTf(x) \subseteq A; \\ f(a)TTf(a)Tf(xr_1r_2)T &\subseteq f(a)TTf(a)Tf(x)T \subseteq A. \end{aligned}$$

Thus $xr_1r_2 \in A : a$. We can show on the same way that $r_1xr_2 \in A : a$ and $r_1r_2x \in A : a$. Let $y \in A : a$ and $a' \in A$. Then $f(a)Tf(a)Tf(a') \subseteq f(a)Tf(a)T(f(y) \cup A) = f(a)Tf(a)Tf(y) \cup f(a)Tf(a)TA \subseteq A$.

Similarly,

$$Tf(a)Tf(a)TTf(a') \subseteq A$$

and

$$f(a)TTf(a)Tf(a')T \subseteq A.$$

Thus $a' \in A : a$. Therefore $A : a$ is an ideal of T containing A . ■

Let f be a good mapping on a ternary semigroup T . Denote the following condition by (α) :

$$\forall F \in fS(T) \forall A \in A(T), F \cap A \neq \emptyset \Rightarrow F^* \cap A \neq \emptyset.$$

Remark 17. If $f(a) = (a)$ for every $a \in T$, then T satisfies the condition (α) .

Proof. Assume $f(a) = (a)$ for every $a \in T$. Let $F \in fS(T)$ and $A \in A(T)$ such that $F \cap A \neq \emptyset$. Let $a \in F \cap A$. Since F is an f -system, $F^* \cap f(a) \neq \emptyset$. By assumption,

$$\emptyset \neq F^* \cap f(a) = F^* \cap (a) \subseteq F^* \cap A. \quad \blacksquare$$

Lemma 18. Let f be a good mapping on a ternary semigroup T and $A, B \in A(T)$. Then the following statements hold:

- (1) $A \subseteq B \Rightarrow r_f(A) \subseteq r_f(B)$;
- (2) $r_f(r_f(A)) = r_f(A)$;
- (3) if T satisfies the condition (α) , then $r_f(A \cap B) = r_f(A) \cap r_f(B)$.

Proof. (1) Assume that $A \subseteq B$. Then

$$r_f(A) = \bigcap \{P \mid A \subseteq P \in fPI(T)\} \subseteq \bigcap \{P \mid B \subseteq P \in fPI(T)\} = r_f(B).$$

(2) It is clear that $r_f(A) \subseteq r_f(r_f(A))$. Let $a \in r_f(r_f(A))$ and F be an f -system of T containing a . Then

$$F \cap r_f(A) \neq \emptyset.$$

This implies

$$F \cap A \neq \emptyset.$$

Thus $a \in r_f(A)$.

(3) Assume that T satisfies the condition (α) . It is clear by (1) that $r_f(A \cap B) \subseteq r_f(A) \cap r_f(B)$. To prove the composite inclusion, let $x \in r_f(A) \cap r_f(B)$. For any f -system F of T containing x , we have that $F \cap A \neq \emptyset$ and $F \cap B \neq \emptyset$. By the condition (α) ,

$$F^* \cap A \neq \emptyset \text{ and } F^* \cap B \neq \emptyset.$$

Let $a \in F^* \cap A$ and $b \in F^* \cap B$. Since F^* is an m -system of T ,

$$\begin{aligned} \emptyset \neq (aTF^*Tb \cup aTTF^*TTb \cup TaTF^*TTb \cup aTTF^*TbT) \cap F^* \\ \subseteq (A \cap B) \cap F. \end{aligned}$$

Thus $x \in r_f(A \cap B)$. ■

4. f -LEFT PRIMARY DECOMPOSITIONS

Definition. Let f be a good mapping on a ternary semigroup T . An ideal P of T is called *left f -primary* if, for $a, b, c \in T$,

$$f(a)Tf(b)Tf(c) \subseteq P, Tf(a)Tf(b)TTf(c) \subseteq P \text{ and } f(a)TTf(b)Tf(c)T \subseteq P$$

imply

$$a \in r_f(P), b \in r_f(P) \text{ or } c \in P.$$

Remark 19. Every f -prime ideal of a ternary semigroup T is a left f -primary ideal of T .

Theorem 20. Let T be a ternary semigroup satisfying the condition (α) . If P_1 and P_2 are left f -primary ideals of T such that $r_f(P_1) = r_f(P_2)$, then $P = P_1 \cap P_2$ is also a left f -primary ideal of T such that $r_f(P) = r_f(P_1) = r_f(P_2)$.

Proof. Assume that P_1 and P_2 are left f -primary ideals of T such that $r_f(P_1) = r_f(P_2)$. Let $P = P_1 \cap P_2$. Then $\emptyset \neq P_1 P_2 \subseteq P_1 \cap P_2$ and

$$r_f(P) = r_f(P_1 \cap P_2) = r_f(P_1) \cap r_f(P_2) = r_f(P_1) \cap r_f(P_1) = r_f(P_1).$$

Let $a, b, c \in T$ be such that $f(a)Tf(b)Tf(c) \subseteq P$, $Tf(a)Tf(b)TTf(c) \subseteq P$ and $f(a)TTf(b)Tf(c) \subseteq P$. Since P_1 and P_2 are left f -primary, $a \in r_f(P)$, $b \in r_f(P)$ or $c \in P_1 \cap P_2 = P$. Thus P is a left f -primary ideal of T . ■

Let T be a ternary semigroup and f a good mapping on T . Denote the following condition by (β) :

$$\forall A, B \in A(T), B \not\subseteq r_f(A) \Rightarrow A : B \neq \emptyset.$$

Theorem 21. Let T be a ternary semigroup satisfying the condition (β) . If an ideal A of T is left f -primary, then $A : B = A$ for every ideal $B \not\subseteq r_f(A)$.

Proof. Assume that A is a left f -primary ideal of T . Let B be an ideal of T not contained in $r_f(A)$. Since $A : B \neq \emptyset$,

$$A : b \neq \emptyset$$

for all $b \in B$. Therefore $A \subseteq A : b$ for all $b \in B$. Hence $A \subseteq A : B$. To show the opposite inclusion, let $a \in A : B$ and $c \in B \setminus r_f(A)$. Then $A : c \neq \emptyset$ and

$$f(c)Tf(c)Tf(a) \subseteq A, Tf(c)Tf(c)TTf(a) \subseteq A \text{ and } f(c)TTf(c)Tf(a)T \subseteq A.$$

Since A is left f -primary and $c \notin r_f(A)$, then $a \in A$. Thus $A : B \subseteq A$. ■

Definition. If an ideal P of a ternary semigroup T can be written as

$$P = P_1 \cap P_2 \cap \dots \cap P_n$$

where each P_i is a left f -primary ideal, then this is called a *left f -primary decomposition* of T and each P_i is called the *left f -primary component* of the decomposition.

Definition. Let $P = \bigcap_{i \in \mathcal{I}} P_i$ be a left f -primary decomposition of a ternary semigroup T . Then P is called *irredundant* if

$$\bigcap_{i \in \mathcal{I} \setminus \{j\}} P_i \not\subseteq P_j$$

for all $j \in \mathcal{I}$. Moreover, an irredundant left f -primary decomposition is called a *normal decomposition* if

$$r_f(P_i) \neq r_f(P_j)$$

for all $i, j \in \mathcal{I}$ such that $i \neq j$.

Let f be a good mapping on a ternary semigroup T . Denote the following condition by (γ) :

$$\text{for any left } f\text{-primary ideal } P \text{ of } T, \text{ we have } P : P = T.$$

Remark 22. Let T be a ternary semigroup. If $f(a) = (a)$ for all $a \in T$, then T satisfies the condition (γ) .

Proof. Let P be a left f -primary ideal of T . For any $x \in T$ and $a \in P$,

$$\begin{aligned} f(a)Tf(a)Tf(x) &= (a)T(a)T(x) \subseteq P; \\ Tf(a)Tf(a)TTf(x) &= T(a)T(a)TT(x) \subseteq P; \\ f(a)TTf(a)Tf(x)T &= (a)TT(a)T(x)T \subseteq P. \end{aligned}$$

Hence $T \subseteq P : P$. ■

Theorem 23. *Let T be a ternary semigroup satisfying the conditions (α) , (β) and (γ) . If an ideal K of T has two normal left f -primary decompositions*

$$K = \bigcap_{i=1}^n P_i = \bigcap_{i=1}^m Q_i,$$

then $n = m$ and $r_f(P_i) = r_f(Q_i)$ for $1 \leq i \leq n = m$ by a suitable ordering.

Proof. This proof is a modification of the proof of Theorem 4.7 in [5]. It is easy to see that the result holds in the case $K = T$. Next we assume that $K \neq T$, where all left f -primary components $P_1, \dots, P_n, Q_1, \dots, Q_m$ are proper ideals of T . We may assume that $r_f(P_1)$ is maximal in the set

$$\{r_f(P_1), \dots, r_f(P_n), r_f(Q_1), \dots, r_f(Q_m)\}.$$

Now we prove that $r_f(P_1) = r_f(Q_i)$ for some $1 \leq i \leq m$. It is enough to show that $P_1 \subseteq r_f(Q_i)$. Suppose that $P_1 \not\subseteq r_f(Q_i)$ for all $1 \leq i \leq m$. Then, by Theorem 21, we have

$$Q_i : P_1 = Q_i$$

for all $1 \leq i \leq m$. Then $K : P_1 = (Q_1 \cap Q_2 \cap \dots \cap Q_m) : P_1 = (Q_1 : P_1) \cap (Q_2 : P_1) \cap \dots \cap (Q_m : P_1) = Q_1 \cap Q_2 \cap \dots \cap Q_m = K$.

Case 1. $n = 1$. By the condition (γ) , we obtain

$$T = P_1 : P_1 = K : P_1 = K,$$

which is a contradiction.

Case 2. $n > 1$. By the condition (γ) and the fact that $P_1 \not\subseteq r_f(P_i)$ for all $2 \leq i \leq n$, we have

$$K = K : P_1 = (P_1 \cap P_2 \cap \dots \cap P_n) : P_1 = (P_1 : P_1) \cap (P_2 : P_1) \cap \dots \cap (P_n : P_1) = T \cap (P_2 : P_1) \cap \dots \cap (P_n : P_1) = (P_2 : P_1) \cap \dots \cap (P_n : P_1) = P_2 \cap P_3 \cap \dots \cap P_n.$$

This is also a contradiction. Thus, $r_f(P_1) \subseteq r_f(Q_i)$ for some $1 \leq i \leq m$. By a suitable ordering, we assume $r_f(P_1) = r_f(Q_1)$.

We use an induction on the number n of left f -primary components. For $n = 1$, we have

$$K = P_1 = \bigcap_{j=1}^m Q_j.$$

Suppose that $m > 1$. Then $P_1 \not\subseteq r_f(Q_j)$ for all $2 \leq j \leq m$. It follows that

$$T = P_1 : P_1 = (Q_1 : P_1) \cap (Q_2 : P_1) \cap \dots \cap (Q_m : P_1) \subseteq Q_m : P_1 = Q_m.$$

This is a contradiction. Thus $m = 1 = n$. Now let us suppose that the conclusion hold for the ideals which are represented by fewer than n of f -primary components. Let $P = P_1 \cap Q_1$. Then P is a left f -primary ideal such that

$$r_f(P) = r_f(P_1) = r_f(Q_1).$$

By the condition (γ) ,

$$T = P_1 : P_1 \subseteq P_1 : P$$

and thus $P_1 : P = T$. From the fact that $P \not\subseteq r_f(P_i)$ for all $2 \leq i \leq n$, we obtain $P_i : P = P_i$ for all $2 \leq i \leq n$. Hence $K : P = \bigcap_{i=1}^n (P_i : P) = (P_1 : P) \cap (P_2 : P) \cap \dots \cap (P_n : P) = T \cap P_2 \cap P_3 \cap \dots \cap P_n = \bigcap_{i=2}^n P_i$.

Similarly, we can show that

$$K : P = \bigcap_{j=2}^m Q_j.$$

Since both decompositions are normal, $n - 1 = m - 1$ implies $n = m$. Moreover, by a suitable ordering, we have $r_f(P_i) = r_f(Q_i)$ for all $2 \leq i \leq n = m$. ■

REFERENCES

[1] S. Bashir and M. Shabir, *Pure ideals in ternary semigroups*, Quasigroups and Related Systems **17** (2009) 149–160.

[2] P. Choo-suwan and R. Chinram, *A study on quasi-ideals in ternary semigroups*, Int. J. Pure and Appl. Math. **77** (2012) 639–647.

[3] V.N. Dixit and S. Dewan, *A note on quasi and bi-ideals in ternary semigroups*, Int. J. Math. and Math. Sci. **18** (1995) 501–508.
<https://doi.org/10.1155/S0161171295000640>

[4] T.K. Dutta, S. Kar and B.K. Maity, *On ideals in regular ternary semigroups*, Discuss. Math. Gen. Alg. and Appl. **28** (2008) 147–159.
<https://doi.org/10.7151/dmgaa.1140>

[5] Z. Gu, *On f -prime radical in ordered semigroups*, Open Mathematics **16** (2018) 574–580.
<https://doi.org/10.1515/math-2018-0053>

[6] E. Hewitt and H.S. Zuckerman, *Ternary operations and semigroups*, in: 1969 Semigroups (Proc. Sympos., Wayne State Univ., Detroit, Mich. 1968), (Academic Press, New York, 1968) 55–83.

[7] S. Kar and B.K. Maity, *Some ideals of ternary semigroups*, Analele Științifice ale Universității Al I Cuza din Iași - Matematică **LVII** (2011) 247–258.
<https://doi.org/10.2478/v10157-011-0024-1>

- [8] K. Murata, Y. Kurata and H. Marubayashi, *A generalization of prime ideals in rings*, Osaka J. Math. **6** (1969) 291–301.
- [9] E. Kasner, *An extension of the group concept*, Bull. Amer. Math. Soc. **10** (1904) 290–291.
- [10] D.M. Rao and Y. Sarala, *A study on d -system, m -system and n -system in ternary semigroups*, Int. J. Development Res. **4** (2014) 195–199.
- [11] M.L. Santiago and S. Sri Bala, *Ternary semigroups*, Semigroup Forum **81** (2010) 380–388.
<https://doi.org/10.1007/s00233-010-9254-x>
- [12] S.K. Sardar and S. Goswami, *f -prime radical of semirings*, South. Asian Bull. Math. **35** (2011) 319–328.
- [13] M. Shabir and M. Bano, *Prime bi-ideals in ternary semigroups*, Quasigroups and Related Systems **16** (2008) 239–256.

Received 28 October 2020

Revised 7 December 2020

Accepted 23 June 2022