Discussiones Mathematicae General Algebra and Applications 43 (2023) 63–74 https://doi.org/10.7151/dmgaa.1410

USING THE SWING LEMMA AND C_1 -DIAGRAMS FOR CONGRUENCES OF PLANAR SEMIMODULAR LATTICES

George Grätzer

Department of Mathematics University of Manitoba Winnipeg, MB R3T 2N2, Canada

e-mail: gratzer@mac.com

Abstract

A planar semimodular lattice K is slim if M_3 is not a sublattice of K. In a recent paper, G. Czédli found four new properties of congruence lattices of slim, planar, semimodular lattices, including the No Child Property: Let \mathcal{P} be the ordered set of join-irreducible congruences of K. Let $x, y, z \in \mathcal{P}$ and let z be a maximal element of \mathcal{P} . If $x \neq y$ and $x, y \prec z$ in \mathcal{P} , then there is no element u of \mathcal{P} such that $u \prec x, y$ in \mathcal{P} .

The Swing Lemma and a standardized diagram type are used to give direct proofs of Czédli's four properties.

 $\label{eq:constraint} \textbf{Keywords:} \ \text{rectangular lattice, slim planar semimodular lattice, congruence lattice.}$

2020 Mathematics Subject Classification: 06C10.

1. INTRODUCTION

Let K be a planar semimodular lattice. We call the lattice K slim if M_3 is not a sublattice of K. In the paper [19, Theorem 1.5], I found a property of congruence lattices of slim, planar, semimodular lattices. In the same paper (see also Problem 24.1 in Grätzer [18]), I proposed the following.

Problem. Characterize the congruence lattices of slim, planar, semimodular lattices.

Czédli [4, Corollaries 3.4, 3.5, Theorem 4.3] found four new properties of congruence lattices of slim, planar, semimodular lattices.

Theorem. Let K be a slim, planar, semimodular lattice with at least three elements and let \mathcal{P} be the ordered set of join-irreducible congruences of K.

- (i) Partition Property: The set of maximal elements of P can be divided into a disjoint union of two nonempty subsets such that no two distinct elements in the same subset have a common lower cover.
- (ii) Maximal Cover Property: If $v \in \mathcal{P}$ is covered by a maximal element u of \mathcal{P} , then u is not the only cover of v.
- (iii) No Child Property: Let x ≠ y ∈ P and let u be a maximal element of P. If x, y ≺ u in P, then there is no element z ∈ P such that z ≺ x, y in P.
- (iv) Four-Cown Two-pendant Property: There is no cover-preserving embedding of the ordered set \mathcal{R} in Figure 1 into \mathcal{P} satisfying the property: any maximal element of \mathcal{R} is a maximal element of \mathcal{P} .

In this note, we will provide a short and direct proof of this theorem using only the Swing Lemma and C_1 -diagrams, see Section 3.

Czédli [4] uses different names for three of the four properties of the Theorem.

Czédli [4]:

Our terminology:

Forbidden Marriage Property Dioecious Maximal Elements Property Bipartite Maximal Elements Property No Child Property Maximal Cover Property Partition Property



Figure 1. The Four-crown Two-pendant ordered set ${\mathfrak R}$ with notation; the covering ${\sf S}_7$ sublattice with edge and element notation.

Outline

Section 2 provides the motivation for Czédli's Theorem. Section 3 provides the tools we need: the Swing Lemma, C_1 -diagrams, and forks.

Section 4 proves the Partition Property, Section 5 does the Maximal Cover Property, while Section 6 verifies the No Child Property.

Finally, The Four-Crown Two-pendant Property is proved in Section 7.

2. MOTIVATION

In my paper [29] with Lakser and Schmidt, we proved that every finite distributive lattice D can be represented as the congruence lattice of a semimodular lattice L. To our surprise, the semimodular lattice K we constructed was *planar*.

Grätzer and Knapp [24]–[28] started the study of planar semimodular lattices. I continued it with my "Notes on planar semimodular lattices" series (started with Knapp): [30] (with Wares), [6, 14] (with Czédli), [21, 23]. See also Czédli and Schmidt [10] and Czédli [1]–[5].

A major subchapter of the theory of planar semimodular lattices started with the observation that in the construction of the lattice K, as in the first paragraph of this section, M_3 sublattices play a crucial role. It was natural to raise the question what can be said about congruence lattices of slim, planar, semimodular (SPS) lattices (see [CFL2, Problem 24.1], originally raised in Grätzer [19]). In [19], I found the first necessary condition and Czédli [2] proved that this condition is not sufficient (see also my related papers [17] and [21]).

A number of papers developed tools to tackle this problem: the Swing Lemma (Grätzer [16]), trajectory coloring (Czédli [1]), special diagrams (Czédli [3]), lamps (Czédli [4]). Some of these results require long proofs. The proof of the trajectory coloring theorem is just shy of 20 pages, while the basic theory of lamps and its application to Theorem 1 is 23 pages.

There are a number of surveys of this field, see the book chapters Czédli and Grätzer [7] and Grätzer [13] in Grätzer and Wehrung, eds. [31]. My presentation [22] provides a gentle review for the background of this topic.

There is an approach to congruences of SPS lattices that is very different from what we are doing in this note. This started in McKenzie [33], Nation [34], Jónsson and Nation [33] and Day [11], viewing SPS lattices as upper bounded homomorphic images of free lattices. Thus the meet-prime elements of a rectangular SPS lattice are along the upper boundaries, so the edges on the upper boundary of a rectangular SPS lattice correspond to maximal join-irreducible congruences, providing a background for the Swing Lemma and its corollaries in Section 3.1. This topic is covered in detail in the book Freese, Ježek and Nation [12, Chapter II], see also the lecture notes of Nation [35, Chapter 10]. The referee informs me that the Maximal Cover Property holds for all finite upper bounded lattices.

3. The tools we need

Most basic concepts and notation not defined in this paper are available in Part I of the book [18], see

https://www.researchgate.net/publication/299594715 arXiv:2104.06539 It is available to the reader and will be referenced as [CFL2].

In particular, we use the notation $C \sim D$, $C \stackrel{\text{up}}{\sim} D$, and $C \stackrel{\text{dn}}{\sim} D$ for perspectivity, up-perspectivity, and down-perspectivity, respectively. As usual, for planar lattices, a prime interval (or covering interval) is called an *edge*. For a finite lattice K and a finite ordered set S, a *cover-preserving* embedding $\varepsilon : S \to K$ is an embedding ε mapping edges of S to edges of K. We define a *cover-preserving* sublattice similarly. For the lattice S_7 of Figure 1, we need a variant: an S_7 sublattice S (a sublattice isomorphic to S_7) is a *peak sublattice* if the three top edges (L, M, and R in Figure 1) are edges in K.

By Grätzer and Knapp [27], every slim, planar, semimodular lattice K has a congruence-preserving extension (see [CFL2, page 43]) \hat{K} to a slim rectangular lattice. Any of the properties (i)–(iv) holds for K iff it holds for \hat{K} . Therefore, in the rest of this paper, we can assume that K is a slim rectangular lattice, simplifying the discussion.

3.1. Swing Lemma

For an edge E of an SPS lattice K, let $E = [0_E, 1_E]$ and define col(E), the *color* of E, as con(E), the (join-irreducible) congruence generated by collapsing E (see [CFL2, Section 3.2]). We write \mathcal{P} for Ji Con K, the ordered set of join-irreducible congruences of K.

As in my paper [16], for the edges U, V of an SPS lattice K, we define a binary relation: U swings to V, in formula, $U \subseteq V$, if $1_U = 1_V$, the element $1_U = 1_V$ of K covers at least three elements, and 0_V is neither the left-most nor the right-most element covered by $1_U = 1_V$; if also 0_U is such, then the swing is *interior*, otherwise, it is *exterior*, denoted by $U \subseteq V$ and $U \subseteq V$, respectively; see Figure 2.



Figure 2. Two swings, $U \subseteq V$; the first $U \stackrel{\text{ex}}{\subseteq} V$, the second $U \stackrel{\text{in}}{\subseteq} V$.

Swing Lemma (Grätzer [16]). Let K be an SPS lattice and let U and V be edges in K. Then $col(V) \leq col(U)$ iff there exists an edge R such that U is

up-perspective to R and there exists a sequence of edges and a sequence of binary relations

$$R = R_0 \ \varrho_1 \ R_1 \ \varrho_2 \cdots \varrho_n \ R_n = V_1$$

where each relation ϱ_i is $\stackrel{\text{dn}}{\sim}$ (down-perspective) or \smile (swing). In addition, this sequence also satisfies

$$1_{R_0} \ge 1_{R_1} \ge \cdots \ge 1_{R_n}.$$

The following statements are immediate consequences of the Swing Lemma, see my papers [16] and [20].

Corollary 1. We use the assumptions of the Swing Lemma.

(i) The equality col(U) = col(V) holds in \mathcal{P} iff there exist edges S and T in K, such that

$$U \stackrel{\text{up}}{\sim} S, S \stackrel{\text{in}}{\smile} T, T \stackrel{\text{dn}}{\sim} V.$$

- (ii) Let us further assume that the element 0_U is meet-irreducible. Then the equality $\operatorname{col}(U) = \operatorname{col}(V)$ holds in \mathcal{P} iff there exists an edge T such that $U \stackrel{\text{in}}{\hookrightarrow} T \stackrel{\text{dn}}{\simeq} V.$
- (iii) If the lattice K is rectangular and U is on the upper boundary of K, then the equality $\operatorname{col}(U) = \operatorname{col}(V)$ holds in \mathfrak{P} iff $U \stackrel{\mathrm{dn}}{\sim} V$.

Note that in (i) the edges S, T, U, V need not be distinct, so we have as special cases $U = V, U \sim V, S = T$, and others.

Corollary 2. We use the assumptions of the Swing Lemma.

(i) The covering $\operatorname{col}(V) \prec \operatorname{col}(U)$ holds in \mathcal{P} iff there exist edges R_1, \ldots, R_4 in K, such that

 $U \stackrel{\mathrm{up}}{\sim} R_1, \ R_1 \stackrel{\mathrm{in}}{\smile} R_2, \ R_2 \stackrel{\mathrm{dn}}{\sim} R_3, \ R_3 \stackrel{\mathrm{ex}}{\smile} R_4, \ R_4 \stackrel{\mathrm{dn}}{\sim} V.$

(ii) If the element 0_U is meet-irreducible, then the covering $\operatorname{col}(V) \prec \operatorname{col}(U)$ holds in \mathfrak{P} iff there exist edges S, T in K, so that $U \stackrel{\mathrm{dn}}{\sim} S \stackrel{\mathrm{ex}}{\hookrightarrow} T \stackrel{\mathrm{dn}}{\sim} V.$

Corollary 3. Let K be a slim rectangular lattice, let U and V be edges in K, and let U be in the upper-left boundary of K.

(i) The covering $\operatorname{col}(V) \prec \operatorname{col}(U)$ holds in \mathcal{P} iff there exist edges S, T in K, such that

(1)
$$U \stackrel{\text{dn}}{\sim} S \stackrel{\text{ex}}{\smile} T \stackrel{\text{dn}}{\sim} V.$$

(ii) Define the element $t = 1_S = 1_T \in K$ and let $S = E_1, E_2, \dots, E_n = W$ enumerate, from left to right, all the edges E of K with $1_E = t$. Then

(2)
$$\operatorname{col}(S) \neq \operatorname{col}(W),$$

- $\operatorname{col}(S) \neq \operatorname{col}(W),$ $\operatorname{col}(E_2) = \dots = \operatorname{col}(E_{n-1}) = \operatorname{col}(T),$ (3)
- $\operatorname{col}(T) \prec \operatorname{col}(S), \operatorname{col}(W).$ (4)

Corollary 4. Let the edge U be on the upper edge of the rectangular lattice K. Then $\operatorname{col}(U)$ is a maximal element of \mathcal{P} .

The converse of this statement is stated in Corollary 8.

3.2. \mathcal{C}_1 -diagrams

In the diagram of a planar lattice K, a normal edge (line) has a slope of 45° or 135° . If it is the first, we call it a normal-up edge (line), otherwise, a normal-down edge (line). Any edge of slope strictly between 45° and 135° is steep (Czédli [3] calls these edges *precipitous*). In Figure 3, for instance, the edges A and D are normal, while the edge S is steep.



Figure 3. Illustrating the proof of The Four-Crown Two-pendant Property.

Definition 5. A diagram of a rectangular lattice K is a C_1 -diagram if the middle edge of any covering S_7 is steep and all other edges are normal.

This concept was introduced in G. Czédli [3, Definition 5.3], see also Czédli [4, Definition 2.1] and Czédli and Grätzer [8, Definition 3.1]. The following is the existence theorem of \mathcal{C}_1 -diagrams in Czédli [3, Theorem 5.5].

Theorem 6. Every rectangular lattice lattice K has a C_1 -diagram.

See Figure 3 for a C_1 -diagram of a rectangular lattice. For a short and direct proof for the existence of \mathcal{C}_1 -diagrams, see my paper [23].

In this paper, K denotes a slim rectangular lattice with a fixed \mathcal{C}_1 -diagram and \mathcal{P} is the ordered set of join-irreducible congruences of K.

Let C and D be maximal chains in an interval [a, b] of K such that $C \cap D = \{a, b\}$. If there is no element of K between C and D, then we call $C \cup D$ a cell. A four-element cell is a 4-cell. Opposite edges of a 4-cell are called *consecutive*. Planar semimodular lattices are 4-cell lattices, that is, all of its cells are 4-cells, see Grätzer and Knapp [24, Lemmas 4, 5] and [CFL2, Section 4.1] for more detail.

The following statement illustrates the use of \mathcal{C}_1 -diagrams.

Lemma 7. Let K be a slim rectangular lattice K with a fixed C_1 -diagram and let X be a normal-up edge of K. Then X is up-perspective either to an edge in the upper-left boundary of K or to a steep edge.

Proof. If X is not steep nor it is in the upper-left boundary of K, then there is a 4-cell C whose lower-right edge is X. If the upper-left edge is steep or it is in the upper-left boundary, then we are done. Otherwise, we proceed the same way until we reach a steep edge or an edge the upper-left boundary.

Corollary 8. Let the edge U be on the upper boundary of K. Then col(U) is a maximal element of \mathcal{P} . Conversely, if u is a maximal element of \mathcal{P} , then there is an edge U on the upper boundary of K so that col(U) = u.

3.3. Trajectories

Czédli and Schmidt [9] introduced a *trajectory* in K as a maximal sequence of consecutive edges, see also [CFL2, Section 4.1]. The *top edge* T of a trajectory is either in the upper boundary of K or it is steep by Lemma 7. For such an edge T, we denote by traj(T) the trajectory with top edge T.

By Grätzer and Knapp [24, Lemma 8], an element a in an SPS lattice K has at most two covers. Therefore, a trajectory has at most one top edge and at most one steep edge. So we conclude the following statement.

Lemma 9. Let K be a slim rectangular lattice K with a fixed C_1 -diagram. Let X and Y be distinct steep edges of K. Then $\operatorname{traj}(X)$ and $\operatorname{traj}(Y)$ are disjoint.

4. The Partition Property

First, we verify the Partition Property for the slim rectangular lattice K with a fixed C_1 -diagram. Let us start with a lemma.

Lemma 10. Let X and Y be distinct edges on the upper-left boundary of K. Then there is no edge Z of K such that $col(Z) \prec col(X), col(Y)$.

Proof. By way of contradiction, let Z be an edge such that $col(Z) \prec col(X), col(Y)$. Since X and Y are on the upper-left boundary, Corollary 3(i) applies. Therefore, there exist normal-up edges S_X, S_Y and steep edges T_X, T_Y such that

$$X \stackrel{\mathrm{dn}}{\sim} S_X \stackrel{\mathrm{ex}}{\smile} T_X, \quad Y \stackrel{\mathrm{dn}}{\sim} S_Y \stackrel{\mathrm{ex}}{\smile} T_Y, \quad Z \in \operatorname{traj}(T_X) \cap \operatorname{traj}(T_Y).$$

By Lemma 9, the third formula implies that $T_X = T_Y$. Since S_X, S_Y are normalup, it follows that X = Y, contradicting the assumption that $X \neq Y$.

By Corollary 8, the set of maximal elements of \mathcal{P} is the same as the set of colors of edges in the upper boundaries, which we can partition into the set of edges \mathcal{L} in the upper-left boundary and the set of edges \mathcal{R} in the upper-right boundary. Let X and Y be distinct edges in \mathcal{L} . By Lemma 10, there is no edge Z of K such that $\operatorname{col}(Z) \prec \operatorname{col}(X), \operatorname{col}(Y)$. By symmetry, this verifies the Partition Property.

5. The Maximal Cover Property

Next, we verify the Maximal Cover Property for the slim rectangular lattice K and with a fixed \mathcal{C}_1 -diagram.

Let $x \in \mathcal{P}$ be covered by a maximal element u of \mathcal{P} in K. By Corollary 8, we can choose an edge U of color u on the upper boundary of K, by symmetry, on the upper-left boundary of K. By Corollary 3(ii), we can choose the edges S, T in K so that $U \stackrel{\text{dn}}{\sim} S \stackrel{\text{ex}}{\smile} T$, $\operatorname{col}(S) = u$, and $\operatorname{col}(T) = x$. By Corollary 3(ii), specifically, by equations (2) and (4), we have $x \prec u, \operatorname{col}(W)$ and $u \neq \operatorname{col}(W)$, verifying the Maximal Cover Property.

6. The No Child Property

In this section, we verify the No Child Property for the slim rectangular lattice K and with a fixed \mathcal{C}_1 -diagram.

Let $x, y, z, u \in \mathcal{P}$ with $x \neq y \in \mathcal{P}$, let u be a maximal element of \mathcal{P} , and let $x, y \prec u$ in \mathcal{P} . By way of contradiction, let us assume that there is an element $z \in \mathcal{P}$ such that $z \prec x, y$ in \mathcal{P} .

By Corollary 8, the element u colors an edge U on the upper boundary of K, say, in the upper-left boundary. By Corollary 2(i), for $z \prec x \in \mathcal{P}$, we get a peak sublattice S_7 in which the middle edge Z is colored by z and upper-left edge X is colored by x, or symmetrically. The upper-right edge Y must have color y.

Now we apply Corollary 3(ii) to the edge U and middle edge Z of the peak sublattice S_7 , obtaining that $U \stackrel{\text{dn}}{\sim} Y \subseteq Z$, in particular, $U \stackrel{\text{dn}}{\sim} Y$. This is a contradiction, since U is normal-up and Y is normal-down.

7. The Four-Crown Two-pendant Property

Finally, we verify the Four-Crown Two-pendant Property for the slim rectangular lattice K and with a fixed C_1 -diagram.

By way of contradiction, assume that the ordered set \mathcal{R} of Figure 1 is a cover-preserving ordered subset of \mathcal{P} , where a, b, c, d are maximal elements of \mathcal{P} . By Corollary 8, there are edges A, B, C, D on the upper boundary of K, so that $\operatorname{col}(A) = a, \operatorname{col}(B) = b, \operatorname{col}(C) = c, \operatorname{col}(D) = d$. By left-right symmetry, we can assume that the edge A is on the upper-left boundary of K. Since $p \prec a, b$ in \mathcal{P} , it follows from Lemma 10 that the edge B is on the upper-right boundary of K, and so is D. Similarly, C is on the upper-left boundary of K.

Because of the automorphisms of the ordered set \mathcal{R} , it is sufficient to deal with one case only: C is below A and B is below D.

By Corollary 2(ii), there is a peak sublattice S_7 with middle edge P (as in the first diagram of Figure 3 so that A and B are down-perspective to the upper-left edge and the upper-right edge of this peak sublattice, respectively. We define, similarly, the edge Q for C and B, the edge S for A and D, the edge R for C and D, and the edge U for R and P. Finally, $v \prec q, s$ in \mathcal{R} , therefore, there is a peak sublattice S_7 with middle edge V with upper-left edge V_l and the upper-right edge V_r so that $S \stackrel{\text{dn}}{\sim} V_l$ and $Q \stackrel{\text{dn}}{\sim} V_r$, or symmetrically. Then V is in both traj(S) and traj(Q), contradicting Lemma 9. This concludes the proof of the Four-Crown Two-pendant Property and of Czédli's Theorem.

Of course, the diagram in Figure 3 is only an illustration. The grid could be much larger, the edges A, C and B, D may not be adjacent, and there maybe lots of other elements in K.

References

- G. Czédli, Patch extensions and trajectory colorings of slim rectangular lattices, Algebra Universalis 72 (2014) 125–154. https://doi.org/10.1007/s00012-014-0294-z
- G. Czédli, A note on congruence lattices of slim semimodular lattices, Algebra Universalis 72 (2014) 225–230. https://doi.org/10.1007/s00012-014-0286-z
- G. Czédli, Diagrams and rectangular extensions of planar semimodular lattices, Algebra Universalis 77 (2017) 443–498. https://doi.org/10.1007/s00012-017-0437-0
- [4] G. Czédli, Lamps in slim rectangular planar semimodular lattices, Acta Sci. Math. (Szeged) 87 (2021) 381–413. https://doi.org/10.14232/actasm-021-865-y0

- [5] G. Czédli, Non-finite axiomatizability of some finite structures. arXiv:2102.00526
- G. Czédli and G. Grätzer, Notes on planar semimodular lattices VII. Resections of planar semimodular lattices, Order 30 (2013) 847–858. https://doi.org/10.1007/s11083-012-9281-1
- [7] G. Czédli and G. Grätzer, Planar Semimodular Lattices: Structure and Diagrams, Chapter 3 in [31]. https://doi.org/10.1007/978-3-319-06413-0_3
- [8] G. Czédli and G. Grätzer, A new property of congruence lattices of slim, planar, semimodular lattices. arXiv:2103.04458
- G. Czédli and E.T. Schmidt, The Jordan-Hölder theorem with uniqueness for groups and semimodular lattices, Algebra Universalis 66 (1-2) (2011) 69-79. https://doi.org/10.1007/s00012-011-0144-1
- [10] G. Czédli and E.T. Schmidt, Slim semimodular lattices I. A visual approach, OR-DER 29 (2012) 481–497. https://doi.org/10.1007/s11083-011-9215-3
- [11] A. Day, Characterizations of finite lattices that are bounded-homomorphic images or sublattices of free lattices, Canad. J. Math. 31 (1979) 69–78.
- [12] R. Freese, J. Ježek and J.B. Nation, Free Lattices, Mathematical Surveys and Monographs 42 (American Mathematical Society, Providence, RI, 1995). https://doi.org/10.1090/surv/042
- G. Grätzer, Planar Semimodular Lattices: Congruences, Chapter 4 in [31]. https://doi.org/10.1007/978-3-319-06413-0_4
- [14] G. Grätzer, Notes on planar semimodular lattices VI. On the structure theorem of planar semimodular lattices, Algebra Universalis 69 (2013) 301–304. https://doi.org/10.1007/s00012-013-0233-4
- G. Grätzer, Two Topics Related to Congruence Lattices of Lattices, Chapter 10 in [31]. https://doi.org/10.1007/978-3-319-06413-0_10
- [16] G. Grätzer, Congruences in slim, planar, semimodular lattices: The Swing Lemma, Acta Sci. Math. (Szeged) 81 (2015) 381–397. https://doi.org/10.1007/978-3-319-38798-7_25
- [17] G. Grätzer, On a result of Gábor Czédli concerning congruence lattices of planar semimodular lattices, Acta Sci. Math. (Szeged) 81 (2015) 25–32. https://doi.org/10.14232/actasm-014-024-1
- [18] G. Grätzer, The Congruences of a Finite Lattice, A Proof-by-Picture Approach, Second Edition (Birkhäuser, 2016). Part I is accessible at arXiv:2104.06539 https://doi.org/10.1007/978-3-319-38798-7

- [19] G. Grätzer, Congruences of fork extensions of lattices, Algebra Universalis 76 (2016) 139–154.
 arXiv:1307.0778
- [20] G. Grätzer, Congruences and trajectories in planar semimodular lattices, Discuss. Math. GAA 38 (2018) 131–142. https://doi.org/10.7151/dmgaa.1280
- [21] G. Grätzer, Notes on planar semimodular lattices VIII. Congruence lattices of SPS lattices, Algebra Universalis 81 (2020). https://doi.org/10.1007/s00012-020-0641-1
- [22] G. Grätzer, A gentle introduction to congruences of planar semimodular lattices, Presentation at the meeting AAA 101, Novi Sad, 2021.
- [23] G. Grätzer, Notes on planar semimodular lattices IX. On Czédli diagrams. arXiv:1307.0778
- [24] G. Grätzer and E. Knapp, Notes on planar semimodular lattices I. Construction, Acta Sci. Math. (Szeged) 73 (2007) 445–462. arXiv:0705.3366
- G. Grätzer and E. Knapp, A note on planar semimodular lattices, Algebra Universalis 58 (2008) 497–499. https://doi.org/10.1007/s00012-008-2089-6
- [26] G. Grätzer and E. Knapp, Notes on planar semimodular lattices II. Congruences, Acta Sci. Math. (Szeged) 74 (2008) 37–47.
- [27] G. Grätzer and E. Knapp, Notes on planar semimodular lattices III. Rectangular lattices, Acta Sci. Math. (Szeged) 75 (2009) 29–48.
- [28] G. Grätzer and E. Knapp, Notes on planar semimodular lattices IV. The size of a minimal congruence lattice representation with rectangular lattices, Acta Sci. Math. (Szeged) 76 (2010) 3–26.
- [29] G. Grätzer, H. Lakser and E.T. Schmidt, Congruence lattices of finite semimodular lattices, Canad. Math. Bull. 41 (1998) 290–297. https://doi.org/10.4153/cmb-1998-041-7
- [30] G. Grätzer and T. Wares, Notes on planar semimodular lattices V. Cover-preserving embeddings of finite semimodular lattices into simple semimodular lattices, Acta Sci. Math. (Szeged) 76 (2010) 27–33.
- [31] G. Grätzer and F. Wehrung eds., Lattice Theory: Special Topics and Applications 1 (Birkhäuser Verlag, Basel, 2014). https://doi.org/10.1007/978-3-319-06413-0
- [32] B. Jónsson and J.B. Nation, Representation of 2-distributive modular lattices of finite length, Acta Sci. Math. (Szeged) 51 (1987) 123–128.
- [33] R.N. McKenzie, Equational bases and nonmodular lattice varieties, Trans. Amer. Math. Soc. 174 (1972) 1–43.

- [34] J.B. Nation, Bounded finite lattices. Universal algebra (Esztergom, 1977), 531–533, Colloq. Math. Soc. János Bolyai 29 (North-Holland, Amsterdam-New York, 1982).
- [35] J.B. Nation, Revised Notes on Lattice Theory. http://www.math.hawaii.edu/jb/books.html

Received 6 June 2021 Revised 18 July 2021 Accepted 19 July 2021