

4  $\pi$ -INVERSE ORDERED SEMIGROUPS

5 AMLAN JAMADAR

6 *Department of Mathematics, Rampurhat College*  
7 *Rampurhat-731224*  
8 *West Bengal, India*

9 **e-mail:** amlanjamadar@gmail.com

10 **Abstract**

11 This article deals with the generalization of  $\pi$ -inverse semigroups with-  
12 out order to ordered semigroups. Here we characterize  $\pi$ -inverse ordered  
13 semigroups by their ordered idempotents and bi-ideals.

14 **Keywords:** bi-ideals, ordered idempotent,  $\pi$ -regular,  $\pi$ -inverse, inverse.

15 **2010 Mathematics Subject Classification:** 06F05, 20M10.

16 1. INTRODUCTION

17 A semigroup  $(S, \cdot)$  with an order relation  $\leq$  is called an ordered semigroup([2],[7])  
18 if for all  $a, b, x \in S$ ,  $a \leq b$  implies  $xa \leq xb$  and  $ax \leq bx$ . It is denoted by  $(S, \cdot, \leq)$ .  
19 Let  $(S, \cdot, \leq)$  be an ordered semigroup. For a subset  $A$  of  $S$ , let  $[A] = \{x \in S :$   
20  $x \leq a, \text{ for some } a \in A\}$ .

21 An element  $a$  of  $S$  is said to be regular (completely regular) [9] if there exists  
22  $x \in S$  such that  $a \leq axa$  ( $a \leq a^2xa^2$ ).  $S$  is called a regular (completely regular)  
23 ordered semigroup if every element of  $S$  is regular (completely regular). Note  
24 that  $S$  is regular (completely regular) if and only if  $a \in (aSa]$  ( $a \in (a^2Sa^2]$ ) for  
25 all  $a \in S$ .

26 An element  $b \in S$  is called an inverse [5] of  $a$  if  $a \leq aba$  and  $b \leq bab$ . The set  
27 of all inverses of an element  $a \in S$  is denoted by  $V_{\leq}(a)$ .  $a', a''$  are the inverse of  
28  $a$  unless otherwise stated.

29 An element  $e \in S$  is said to be an ordered idempotent if  $e \leq e^2$ . The set of  
30 all ordered idempotents of  $S$  is denoted by  $E_{\leq}(S)$ .

31 Bhuniya and Hansda [1] studied the ordered semigroups in which any two  
32 inverses of an element are  $\mathcal{H}$ -related. Class of these ordered semigroups are natu-  
33 ral generalization of the class of all inverse semigroups. Hansda and Jamadar [5]

named these ordered semigroups as inverse ordered semigroups and studied their different aspects. In this paper, we further extend inverse ordered semigroups to  $\pi$ -inverse ordered semigroups.

A nonempty subset  $A$  of  $S$  is called a left (right) ideal [8] of  $S$ , if  $SA \subseteq A$  ( $AS \subseteq A$ ) and  $(A] = A$ . A nonempty subset  $A$  is called a (two-sided) ideal of  $S$  if it is both a left and a right ideal of  $S$ . Following Kehayopulu [9], a nonempty subset  $B$  of an ordered semigroup  $S$  is called a bi-ideal of  $S$  if  $BSB \subseteq B$  and  $(B] = B$ . Hansda [4] studied algebraic properties of bi-ideals in completely regular and Clifford ordered semigroups.

The principal [8] left ideal, right ideal, ideal and bi-ideal [9] generated by  $a \in S$  are denoted by  $L(a)$ ,  $R(a)$ ,  $I(a)$  and  $B(a)$  respectively. It is easy to show that

$$L(a) = (a \cup Sa], R(a) = (a \cup aS], I(a) = (a \cup Sa \cup aS \cup SaS] \text{ and } B(a) = (a \cup aSa].$$

Kehayopulu [8] defined Green's relations  $\mathcal{L}$ ,  $\mathcal{R}$ ,  $\mathcal{J}$  and  $\mathcal{H}$  on an ordered semigroup  $S$  as follows:

$$a\mathcal{L}b \text{ if } L(a) = L(b), a\mathcal{R}b \text{ if } R(a) = R(b), a\mathcal{J}b \text{ if } I(a) = I(b) \text{ and } \mathcal{H} = \mathcal{L} \cap \mathcal{R}.$$

These four relations are equivalence relations on  $S$ .

An ordered semigroup  $S$  is called  $\pi$ -regular (resp. completely  $\pi$ -regular) [3] if for every  $a \in S$  there is  $m \in \mathbb{N}$  such that  $a^m \in (a^m Sa^m]$  (resp.  $a^m \in (a^{2m} Sa^{2m}]$ ). The set of all regular, completely regular, inverse and  $\pi$ -regular elements in an ordered semigroup  $S$  is denoted by  $Reg_{\leq}(S)$ ,  $Gr_{\leq}(S)$ ,  $V_{\leq}(S)$  and  $\pi Reg_{\leq}(S)$  respectively.

Let  $S$  be an ordered semigroup and  $\rho$  be an equivalence relation on  $S$ . Following Hansda and Jamadar [5], an element  $a \in S$  of type  $\tau$  is said to be a  $\rho$ -unique element in  $S$  if for every other element  $b \in S$  of type  $\tau$  we have  $a\rho b$ .

**Theorem 1** [5]. *The following conditions are equivalent on an ordered semigroup  $S$ .*

1.  $S$  is an inverse ordered semigroup;
2.  $S$  is regular and its idempotents are  $\mathcal{H}$ -commutative;
3. For every  $e, f \in E_{\leq}(S)$ ,  $e\mathcal{L}f(e\mathcal{R}f)$  implies  $e\mathcal{H}f$ .

## 2. $\pi$ -INVERSE ORDERED SEMIGROUP

This section deals with the characterization of the class of  $\pi$ -inverse ordered semigroups.

Let  $S$  be a  $\pi$ -regular ordered semigroup. Then for every  $a \in S$  there is  $m \in \mathbb{N}$  such that  $a^m \leq a^m x a^m \leq a^m (x a^m x) a^m$  and  $x a^m x \leq x a^m x (a^m) x a^m x$ . Thus for every  $a \in S$  there is  $m \in \mathbb{N}$  such that  $V_{\leq}(a^m) \neq \phi$ .

**Definition.** A  $\pi$ -regular ordered semigroup  $S$  is called  $\pi$ -inverse if for every  $a \in S$ , there is  $m \in \mathbb{N}$  such that any two inverses of  $a^m$  are  $\mathcal{H}$ -related.

For  $a \in S$ , there is  $m \in \mathbb{N}$  such that every principal left ideal and every principal right ideal generated by  $a^m$  in a  $\pi$ -inverse ordered semigroup have  $\mathcal{H}$ -unique ordered idempotent generator. This has been shown in the following theorem.

**Theorem 2.** A  $\pi$ -regular ordered semigroup  $S$  is  $\pi$ -inverse if and only if for every  $a \in S$  there is  $m \in \mathbb{N}$  such that  $(Sa^m]$  and  $(a^m S]$  are generated by an  $\mathcal{H}$ -unique ordered idempotent.

**Proof.** Suppose that  $S$  is  $\pi$ -inverse. Let  $a \in S$ . Since  $S$  is  $\pi$ -regular, there is  $m \in \mathbb{N}$  such that  $a^m \leq a^m z a^m$  for some  $z \in S$ . Let  $I = (Sa^m]$ . Then clearly  $I = (Sa^m z a^m] = (Se]$ , where  $e = z a^m \in E_{\leq}(S)$ . If possible let  $I = (Sf]$  for some  $f \in E_{\leq}(S)$ . Then  $e \mathcal{L} f$  and so  $e \leq x f$  and  $f \leq y e$  for some  $x, y \in S$ . Now  $e \leq e e \leq e e e \leq e x f e$ . Therefore  $e x f \leq e x f e x f$  so that  $e x f \in E_{\leq}(S)$ . Also  $e x f \leq e x f e x f \leq e x f (f e) e x f$  and  $f e \leq f e e e \leq f e x f e \leq f e (e x f) f e$ . Therefore  $f e \in V_{\leq}(e x f)$ . Also  $e x f \in V_{\leq}(e x f)$ . Since  $S$  is  $\pi$ -inverse for  $f e, e x f \in V_{\leq}(e x f)$  we have  $f e \mathcal{H} e x f$ . Then  $e \leq e e \leq e e e \leq e x f e \leq e x f f e \leq f e t t_1 e x f$  for some  $t, t_1 \in S$  and so  $e \leq f z_1$ , where  $z_1 = e t t_1 e x f$ . Similarly  $f \leq e z_2$  for some  $z_2 \in S$ . So  $e \mathcal{R} f$ . Hence  $e \mathcal{H} f$ . Likewise  $(a^m S]$  is generated by an  $\mathcal{H}$ -unique ordered idempotent.

Conversely assume that given condition holds in  $S$ . Then  $S$  is  $\pi$ -regular. Let  $a \in S$  and  $a', a'' \in V_{\leq}(a^m)$  for some  $m \in \mathbb{N}$ . Clearly  $(Sa^m] = (Sa' a^m] = (Sa'' a^m]$ . Since  $a' a^m, a'' a^m \in E_{\leq}(S)$  we have that  $a' a^m \mathcal{H} a'' a^m$ , by given condition. Then there are  $s, v \in S$  such that  $a' \leq a' a^m a' \leq a'' a^m s a'$  and  $a'' \leq a' a^m v a''$ . Thus  $a' \mathcal{R} a''$ . Likewise  $a' \mathcal{L} a''$ , that is  $a' \mathcal{H} a''$ . Hence  $S$  is a  $\pi$ -inverse ordered semigroup.

■

The following theorem shows some equivalent conditions for an ordered semigroup  $S$  to be  $\pi$ -inverse.

**Theorem 3.** The following conditions are equivalent on an ordered semigroup  $S$ .

1.  $S$  is a  $\pi$ -inverse ordered semigroup;
2.  $S$  is  $\pi$ -regular and for every  $e, f \in E_{\leq}(S)$ , there is  $m \in \mathbb{N}$  such that  $(e f)^m \in (f S e]$ ;

96 3.  $S$  is  $\pi$ -regular and for every  $e, f \in E_{\leq}(S)$ ,  $e\mathcal{L}f(e\mathcal{R}f)$  implies  $e\mathcal{H}f$ .

97 **Proof.** (1)  $\Rightarrow$  (2): First suppose  $S$  is  $\pi$ -inverse. Then  $S$  is  $\pi$ -regular. Let  $e, f \in$   
 98  $E_{\leq}(S)$ . Since  $S$  is  $\pi$ -regular, for  $ef \in S$  there is  $x \in S$  such that  $x \in V_{\leq}(ef)^m$   
 99 for some  $m \in \mathbb{N}$ . We consider the following cases.

100 **Case 1:** If  $m = 1$  then  $ef \in (fSe]$  holds, by Theorem 1.

101 **Case 2:** If  $m > 1$  then  $x \leq x(ef)^m x$  implies that  $fxe \leq fxe(ef)^m fxe$ . Also  
 102  $(ef)^m \leq (ef)^m x(ef)^m$  implies that  $(ef)^m \leq (ef)^m (fxe)(ef)^m$ . Thus  $(ef)^m \in$   
 103  $V_{\leq}(fxe)$ . Now  $x \leq x(ef)^m x = xe(fe)^{m-1} fx$  so that  $fxe \leq fxe(fe)^{m-1} fxe \leq$   
 104  $fxe(fe)^{m-1} fxe(fe)^{m-1} fxe$  and  $(fe)^{m-1} fxe(fe)^{m-1} \leq (fe)^{m-1} fxe(fe)^{m-1} fxe$   
 105  $(fe)^{m-1} \leq (fe)^{m-1} fxe(fe)^{m-1} fxe(fe)^{m-1} fxe$   
 106  $(fe)^{m-1}$ . This gives  $(fe)^{m-1} fxe(fe)^{m-1} \in V_{\leq}(fxe)$ . Thus  $(ef)^m, (fe)^{m-1} fxe(fe)^{m-1} \in$   
 107  $V_{\leq}(fxe)$ . Since  $S$  is  $\pi$ -inverse, we have that  $(fe)^{m-1} fxe(fe)^{m-1} \mathcal{H}(ef)^m$ . Then  
 108 there are  $s_1, s_2 \in S$  such that  $(ef)^m \leq (fe)^{m-1} fxe(fe)^{m-1} s_1$  and  $(ef)^m \leq$   
 109  $s_2 (fe)^{m-1} fxe(fe)^{m-1}$ . Thus from the inequality  $(ef)^m \leq (ef)^m x(ef)^m$  we have  
 110 that  $(ef)^m \leq (fe)^{m-1} fxe(fe)^{m-1}$   
 111  $s_1 x s_2 (fe)^{m-1} fxe(fe)^{m-1} \leq f(fe)^{m-1} fxe(fe)^{m-1} s_1 x s_2 (fe)^{m-1} fxe(fe)^{m-1} e$ . There-  
 112 fore  $(ef)^m \leq fye$ , where  $y = (fe)^{m-1} fxe(fe)^{m-1} s_1 x s_2 (fe)^{m-1} fxe(fe)^{m-1} \in S$ .  
 113 Hence  $(ef)^m \in (fSe]$ .

114 (2)  $\Rightarrow$  (3): Let  $e, f \in E_{\leq}(S)$  be such that  $e\mathcal{L}f$ . Then  $e \leq xf$  and  $f \leq ye$  for  
 115 some  $x, y \in S$ . Now  $e \leq xf$  implies  $e \leq exf$  and so  $e \leq ee \leq exfe$ , which implies  
 116 that  $exf \leq exfexf$ . So  $exf \in E_{\leq}(S)$ . Similarly  $f \leq fye$  and  $fye \in E_{\leq}(S)$ . Now

$$(1) \quad e \leq exf \leq exff \leq (exf)(fye).$$

117 Since  $exf, fye \in E_{\leq}(S)$ , there exists  $m \in \mathbb{N}$  such that  $(exffye)^m \in ((fye)S(exf))$ ,  
 118 by condition (2). Then there exists  $z \in S$  such that  $(exffye)^m \leq (fye)z(exf)$ .  
 119 Thus  $e \leq e^m$  together with (1) implies that  $e \leq (exffye)^m$  and therefore  
 120  $e \in ((fye)S(exf)) \subseteq (fS]$ . Likewise  $f \in (eS]$ , that is,  $e\mathcal{R}f$ . Hence  $e\mathcal{H}f$ .

121 For  $e\mathcal{R}f$ ,  $e\mathcal{H}f$  follows dually.

122 (3)  $\Rightarrow$  (1): Let  $a \in S$  and  $a', a'' \in V_{\leq}(a^m)$  for some  $m \in \mathbb{N}$ . Now  $a^m a' \leq$   
 123  $a^m a'' a^m a'$  and  $a^m a'' \leq a^m a' a^m a''$  which gives  $a^m a' \mathcal{R} a^m a''$  so that  $a^m a' \mathcal{H} a^m a''$ ,  
 124 by the condition (3). Likewise  $a' a^m \mathcal{H} a'' a^m$ . Then  $a' \leq a' a^m a'$  gives that  $a' \leq$   
 125  $a'' a^m x a^m$  for some  $x \in S$ . Therefore  $a' \leq a'' t$  where  $t = a^m x a^m$ . In a similar  
 126 manner it is possible to get  $u, v, w \in S$  such that  $a' \leq ua''$ ,  $a'' \leq a'v$  and  $a'' \leq wa'$ .  
 127 So  $a' \mathcal{H} a''$ . Hence  $S$  is a  $\pi$ -inverse ordered semigroup. ■

128 Let  $S$  be a  $\pi$ -regular ordered semigroup. Then for every  $a \in S$  there is  $m \in \mathbb{N}$   
 129 such that  $a^m \leq a^m x a^m$  for some  $x \in S$  which gives that  $a^m \leq a^m x (a^m) x a^m$ . Here  
 130  $a^m x, x a^m \in E_{\leq}(S)$  so that  $a^m \in (eSf]$ , for  $e = a^m x$  and  $f = x a^m$ .

131 Following this idea we find a condition for a  $\pi$ -regular ordered semigroup to  
 132 be  $\pi$ -inverse.

**Theorem 4.** *A  $\pi$ -regular ordered semigroup  $S$  is  $\pi$ -inverse if and only if for every  $e, f \in E_{\leq}(S)$  and  $x \in S$  whenever  $x^m \in (eSf]$  for some  $m \in \mathbb{N}$ , then  $x' \in (fSe]$  for every  $x' \in V_{\leq}(x^m)$ .*

**Proof.** First suppose that  $S$  is a  $\pi$ -inverse ordered semigroup. Then there is  $m \in \mathbb{N}$  such that  $V_{\leq}(x^m) \neq \emptyset$ . Let  $x' \in V_{\leq}(x^m)$ . Suppose  $x^m \in (eSf]$  for  $e, f \in E_{\leq}(S)$ . Then  $x^m \leq es_1f$  for some  $s_1 \in S$ . Now  $x' \leq x'x^mx' \leq x'es_1fx'$  and so  $es_1fx' \leq es_1fx'es_1fx'$ , that is  $es_1fx' \in E_{\leq}(S)$ . Similarly  $x'es_1f \in E_{\leq}(S)$ . Therefore  $x' \leq x'(es_1fx')^r$  and  $x' \leq (x'es_1f)^rx'$  for all  $r \in \mathbb{N}$ . Now since  $S$  is  $\pi$ -inverse, for  $f, x'es_1f \in E_{\leq}(S)$  there are  $s_2 \in S$  and  $n \in \mathbb{N}$  such that  $(x'es_1ff)^n \leq fs_2x'es_1f$ , by Theorem 3(2). Similarly for  $e, es_1fx' \in E_{\leq}(S)$  we have  $(ees_1fx')^k \leq es_1fx's_3e$ , for some  $s_3 \in S$  and  $k \in \mathbb{N}$ . Then  $x' \leq x'x^mx'$  implies that  $x' \leq (x'es_1ff)^nx'(ees_1fx')^k \leq fs_2x'es_1x'fes_1fx's_3e$ . Hence  $x' \in (fSe]$ .

Conversely, assume that the given condition holds in  $S$ . Let  $e, f \in E_{\leq}(S)$  be such that  $e\mathcal{L}f$ , this yields that  $e \leq ee \leq ezf$  for some  $z \in S$ . Therefore  $e^m \in (eSf]$ . Since  $e \in V_{\leq}(e^m)$  we have  $e \in (fSe]$ , by given condition. Likewise  $f \in (eSf]$ . This implies that  $e\mathcal{R}f$  and so  $e\mathcal{H}f$ . Thus by Theorem 3,  $S$  is a  $\pi$ -inverse ordered semigroup. ■

**Corollary 5.** *The following conditions are equivalent on a  $\pi$ -regular ordered semigroup  $S$ .*

1.  $S$  is a  $\pi$ -inverse ordered semigroup;
2. Let  $a \in S$ . Then there are  $m, n \in \mathbb{N}$  such that  $(a^ma'a^ma^m)^n \in (a'Sa']$ , for every  $a' \in V_{\leq}(a^m)$ ;
3. Any two inverses of an ordered idempotent in  $S$  are  $\mathcal{H}$ -related;
4. All inverses of  $e$  are  $\mathcal{H}$ -commutative, for every  $e \in E_{\leq}(S)$ ;
5. For any  $e \in E_{\leq}(S)$  and  $e' \in V_{\leq}(e)$ ,  $ee'e'e \in (e'Se']$ .

**Proof.** (1)  $\Rightarrow$  (2), (2)  $\Rightarrow$  (3), (3)  $\Rightarrow$  (4): These are obvious.

(4)  $\Rightarrow$  (5): Let  $e \in E_{\leq}(S)$  and  $e' \in V_{\leq}(e)$ . Then  $ee'e'e \leq e's_1ees_2e'$  for some  $s_1, s_2 \in S$ . Hence  $ee'e'e \in (e'Se']$ .

(5)  $\Rightarrow$  (1): Let  $a \in S$  and  $a', a'' \in V_{\leq}(a^m)$  for some  $m \in \mathbb{N}$ . Then  $a' \leq a'a^ma' \leq a'a^ma''a^ma' \leq a''a^ms_4a'a^ma'$ , for some  $s_4 \in S$ . Therefore  $a' \leq a''t_1$  where  $t_1 = a^ms_4a'a^ma'$ . Similarly there exists  $t_2 \in S$  such that  $a' \leq t_2a''$ . Also there are  $t_3, t_4 \in S$  such that  $a'' \leq t_3a'$  and  $a'' \leq a't_4$ . Thus  $a'\mathcal{H}a''$ . Hence  $S$  is a  $\pi$ -inverse ordered semigroup. ■

**Corollary 6.** *Let  $S$  be a  $\pi$ -inverse ordered semigroup and  $a, b \in S$ . If  $m, n \in \mathbb{N}$  are such that  $V_{\leq}(a^m), V_{\leq}(b^n) \neq \emptyset$ , then the following statements hold in  $S$ .*

- 169 1.  $a^m \mathcal{L}b^n$  if and only if  $a'a^m \mathcal{H}b'b^n$  for every  $a' \in V_{\leq}(a^m)$  and  $b' \in V_{\leq}(b^n)$ ;  
 170 2.  $a^m \mathcal{R}b^n$  if and only if  $a^m a' \mathcal{H}b'b^n$  for every  $a' \in V_{\leq}(a^m)$  and  $b' \in V_{\leq}(b^n)$ ;  
 171 3.  $a^m \mathcal{H}b^n$  if and only if  $a'a^m \mathcal{H}b'b^n$  and  $a^m a' \mathcal{H}b'b^n$  for every  $a' \in V_{\leq}(a^m)$  and  
 172  $b' \in V_{\leq}(b^n)$ .

173 **Proof.** (1): Let  $a, b \in S$ . Since  $S$  is  $\pi$ -inverse, there are  $m, n \in \mathbb{N}$  such that  
 174  $V_{\leq}(a^m), V_{\leq}(b^n) \neq \phi$ . Let  $a' \in V_{\leq}(a^m), b' \in V_{\leq}(b^n)$ . Let  $a^m \mathcal{L}b^n$ . Since  $a^m \leq$   
 175  $a^m a' a^m$  and  $a' a^m \leq a' a^m a' a^m$ , we have  $a^m \mathcal{L}a' a^m$ , which implies that  $b^n \mathcal{L}a' a^m$ .  
 176 Also  $b^n \mathcal{L}b'b^n$ . Hence  $a' a^m \mathcal{L}b'b^n$ . Since  $a' a^m, b'b^n \in E_{\leq}(S)$  and  $S$  is  $\pi$ -inverse we  
 177 have  $a' a^m \mathcal{H}b'b^n$ , by Theorem 3(3).

178 Conversely suppose that given condition holds in  $S$ . Let  $a, b \in S$  with  $a' \in$   
 179  $V_{\leq}(a^m)$  and  $b' \in V_{\leq}(b^n)$  for some  $m, n \in \mathbb{N}$ . Then by given condition  $a' a^m \mathcal{H}b'b^n$ .  
 180 Also we have  $a^m \mathcal{L}a' a^m$  and  $b^n \mathcal{L}b'b^n$  so that  $a^m \mathcal{L}b^n$ .

181 (2) and (3): These follow dually. ■

### 182 3. BI-IDEALS IN $\pi$ -INVERSE ORDERED SEMIGROUPS

183 In this section we characterize a  $\pi$ -inverse ordered semigroup  $S$  by the principal  
 184 bi-ideals of  $S$ .

185 **Theorem 7.** *Let  $S$  be a  $\pi$ -regular ordered semigroup. Then the following condi-*  
 186 *tions are equivalent.*

- 187 1.  $S$  is a  $\pi$ -inverse ordered semigroup;  
 188 2. For any  $a \in S$ , there is  $m \in \mathbb{N}$  such that  $B(a') = B(a'')$  for every  $a', a'' \in$   
 189  $V_{\leq}(a^m)$ ;  
 190 3. For any  $e, f \in E_{\leq}(S)$ ,  $B((ef)^m) \subseteq B(e) \cap B(f)$  for some  $m \in \mathbb{N}$ ;  
 191 4. For any  $e \in E_{\leq}(S)$  and  $x \in V_{\leq}(e)$ ,  $B(ex) = B(xe)$ .

192 **Proof.** (1)  $\Rightarrow$  (2): First suppose that  $S$  is a  $\pi$ -inverse ordered semigroup. Let  
 193  $a \in S$ . Then there is  $m \in \mathbb{N}$  such that  $a', a'' \in V_{\leq}(a^m)$ . Suppose  $x \in B(a')$ .  
 194 Therefore  $x \leq a'$  or  $x \leq a'ya'$  for some  $y \in S$ . Since  $S$  is  $\pi$ -inverse,  $a' \mathcal{H}a''$ . If  
 195  $x \leq a'$  then  $x \leq a' a^m a' \leq a'' s_1 a^m s_2 a''$  for some  $s_1, s_2 \in S$ . Therefore  $x \leq a'' s a''$   
 196 where  $s = s_1 a^m s_2$ . If  $x \leq a'ya'$  then there is  $s_3 \in S$  such that  $x \leq a'' s_3 a''$ . Thus  
 197 in either case  $x \in B(a'')$ . Also  $a' \in B(a'')$  implies that  $B(a') \subseteq B(a'')$ . Similarly  
 198  $B(a'') \subseteq B(a')$ . Hence  $B(a') = B(a'')$ .

199 (2)  $\Rightarrow$  (3): Let  $e, f \in E_{\leq}(S)$  and  $x \in V_{\leq}(ef)^m$  for some  $m \in \mathbb{N}$ . Clearly  
 200  $(ef)^m, (fe)^{m-1} fxe(fe)^{m-1} \in V_{\leq}(fxe)$  and so by the condition (2) it follows that

201  $B((ef)^m) = B((fe)^{m-1}fxe(fe)^{m-1})$ . Now  $(ef)^m \in B((fe)^{m-1}fxe(fe)^{m-1})$  im-  
 202 plies  $(ef)^m \leq (fe)^{m-1}fxe(fe)^{m-1}$  or  $(ef)^m \leq (fe)^{m-1}fxe(fe)^{m-1}h(fe)^{m-1}fxe(fe)^{m-1}$   
 203 for some  $h \in S$ . So in either case  $(ef)^m \leq h_1(fe)^{m-1}fxe(fe)^{m-1}$  and  $(ef)^m \leq$   
 204  $(fe)^{m-1}fxe(fe)^{m-1}h_2$  for some  $h_1, h_2 \in S$ . Likewise there are  $h_3, h_4 \in S$  such  
 205 that  $(fe)^{m-1}fxe(fe)^{m-1} \leq h_3(ef)^m$  and  $(fe)^{m-1}fxe(fe)^{m-1} \leq (ef)^mh_4$ . Hence  
 206  $(ef)^m\mathcal{H}(fe)^{m-1}$   
 207  $fxe(fe)^{m-1}$ .

208 Let  $w \in B(ef)^m$ . Then either  $w \leq (ef)^m$  or  $w \leq (ef)^ms_1(ef)^m$  for some  $s_1 \in$   
 209  $S$ . If  $w \leq (ef)^m$  then  $w \leq (ef)^m \leq (ef)^mx(ef)^m \leq (ef)^mxs_2(fe)^{m-1}fxe(fe)^{m-1}$   
 210 for some  $s_2 \in S$ .

211 Also  $w \leq (ef)^ms_1(ef)^m$  gives  $w \leq efs_1s_3(fe)^{m-1}fxe(fe)^{m-1}$  for some  $s_3 \in$   
 212  $S$ . So in either case  $w \in B(e)$ . Likewise  $w \in B(f)$ . Therefore  $w \in B(e) \cap B(f)$   
 213 and hence  $B(ef)^m \subseteq B(e) \cap B(f)$ .

214 (3)  $\Rightarrow$  (4): Let  $e \in E_{\leq}(S)$  and  $x \in V_{\leq}(e)$ . Then  $e, xe, ex \in E_{\leq}(S)$ . Now  
 215 by condition (3)  $B((exe)^m) \subseteq B(e) \cap B(xe)$  for some  $m \in \mathbb{N}$ . Let  $y \in B(e)$ .  
 216 Then either  $y \leq e$  or  $y \leq es_3e$  for some  $s_3 \in S$ . If  $y \leq e$  then  $y \leq exe \leq$   
 217  $eeexe \leq exeeexe \leq \dots \leq (exe)^m$ . So  $y \in B((exe)^m)$ . Likewise  $y \in B((exe)^m)$   
 218 for the case  $y \leq es_3e$ . Therefore  $B(e) = B((exe)^m)$  and so  $B(e) \subseteq B(xe)$ .  
 219 Also  $B((xee)^n) \subseteq B(e) \cap B(xe)$  for some  $n \in \mathbb{N}$ , then by a similar argument  
 220  $B(xe) \subseteq B(e)$ . Therefore  $B(e) = B(xe)$ . Likewise  $B(e) = B(ex)$ . Therefore  
 221  $B(xe) = B(ex)$ .

222 (4)  $\Rightarrow$  (1): By condition (4) we have  $ex\mathcal{H}xe$ . Also  $ex \in B(e)$  and  $ex \in B(x)$ .  
 223 Then  $ex \leq e$  or  $ex \leq eb_1e$  and  $ex \leq x$  or  $ex \leq xb_2x$  for some  $b_1, b_2 \in S$ . Here  
 224 following cases arise.

225 **Case(1):** If  $ex \leq e$  and  $ex \leq x$  then  $ex \leq exex \leq xe \leq xexe = xae$  where  
 226  $a = ex$ .

227 **Case(2):** If  $ex \leq e$  and  $ex \leq xb_2x$  then  $ex \leq exex \leq xb_2xe = xbe$  where  
 228  $b = b_2x$ .

229 **Case(3):** If  $ex \leq eb_1e$  and  $ex \leq x$  then  $ex \leq exex \leq xeb_1e = xce$  where  
 230  $c = eb_1$ .

231 **Case(4):** If  $ex \leq eb_1e$  and  $ex \leq xb_2x$  then  $ex \leq exex \leq xb_2xeb_1e = xde$   
 232 where  $d = b_2xeb_1$ . Therefore in either case  $ex \leq xse$  for some  $s \in S$ . Similarly  
 233  $xe \leq etx$  for some  $t \in S$ . Thus  $e, x$  are  $\mathcal{H}$ -commutative. Hence by Corollary 5,  $S$   
 234 is a  $\pi$ -inverse ordered semigroup. ■

235 **Corollary 8.** A  $\pi$ -regular ordered semigroup  $S$  is  $\pi$ -inverse if and only if for any  
 236  $e \in E_{\leq}(S)$  and  $x \in V_{\leq}(e)$ ,  $B(ex) = B(e) \cap B(x) = B(xe) = B(e) = B(x)$ .

237 **Proof.** This follows from Theorem 7. ■

238 **Corollary 9.** A  $\pi$ -regular ordered semigroup  $S$  is  $\pi$ -inverse if and only if for any  
 239  $e, f \in E_{\leq}(S)$ ,  $e\mathcal{L}f(e\mathcal{R}f)$  implies  $B(e) = B(f)$ .

**Proof.** Let  $S$  be a  $\pi$ -inverse ordered semigroup. Since  $S$  is  $\pi$ -inverse  $e\mathcal{L}f(e\mathcal{R}f)$  implies  $e\mathcal{H}f$  by Theorem 3. So it is easy to check that  $B(e) = B(f)$ .

Conversely suppose that the condition holds in  $S$ . Now  $B(e) = B(f)$  gives that  $e \in B(f)$  and  $f \in B(e)$ . Therefore  $e \leq f$  or  $e \leq fxf$  and  $f \leq e$  or  $f \leq eye$  for some  $x, y \in S$ . In either case  $e\mathcal{R}f$ . So  $e\mathcal{L}f$  implies  $e\mathcal{H}f$ . Hence  $S$  is a  $\pi$ -inverse ordered semigroup, by Theorem 3. ■

**Corollary 10.** Let  $S$  be a  $\pi$ -inverse ordered semigroup and  $a, b \in S$ . If  $a' \in V_{\leq}(a^m)$ ,  $b' \in V_{\leq}(b^n)$ , for some  $m, n \in \mathbb{N}$ , then the following conditions hold on  $S$ .

1.  $a^m\mathcal{L}b^n$  if and only if  $B(a'a^m) = B(b'b^n)$ .

2.  $a^m\mathcal{R}b^n$  if and only if  $B(a^ma') = B(b^nb')$ .

**Proof.** (1): Let  $S$  be a  $\pi$ -inverse ordered semigroup and  $a, b \in S$ . Also let  $a' \in V_{\leq}(a^m)$ ,  $b' \in V_{\leq}(b^n)$  for some  $m, n \in \mathbb{N}$ , such that  $a^m\mathcal{L}b^n$ . So by Corollary 6  $a'a^m\mathcal{H}b'b^n$ . Let  $x \in B(a'a^m)$ . Therefore  $x \leq a'a^m$  or  $x \leq a'a^ms_1a'a^m$  for some  $s_1 \in S$ . So it is easy to verify that  $x \in B(b'b^n)$ . Also  $a'a^m \in B(b'b^n)$ . So  $B(a'a^m) \subseteq B(b'b^n)$ . Similarly  $B(b'b^n) \subseteq B(a'a^m)$ . So  $B(a'a^m) = B(b'b^n)$ .

Converse follows easily.

(2): This is similar to (1). ■

### Acknowledgements

I express my deepest gratitude to the editor of the journal Professor Joanna Skowronek-Kaziow for communicating the paper and to the referee of the paper for their important valuable comments and suggestions to enrich the quality of the paper both in value and content.

### REFERENCES

- [1] A.K. Bhuniya and K. Hansda, *On completely regular and Clifford ordered semigroups*, Afrika Mat. **31** (2020) 1029–1045.  
<https://doi.org/10.1007/s13370-020-00778-1>
- [2] G. Birkhoff, *Lattice Theory* (Providence, 1969).
- [3] Y. Cao and X. Xinzhai, *Nil-extensions of simple po-semigroups*, Comm. Algebra **28**(5) (2000) 2477–2496.
- [4] K. Hansda, *Bi-ideals in Clifford ordered semigroup*, Discuss. Math. General Alg. Appl. **33** (2013) 73–84.
- [5] K. Hansda and A. Jamadar, *Characterization of inverse ordered semigroups by their ordered idempotents and bi-ideals*, Quasigroups and Related Systems **28** (2020) 77–88.



- 274 [6] N. Kehayopulu, *Remark in ordered semigroups*, Math. Japonica **35** (1990) 1061–  
275 1063.
- 276 [7] N. Kehayopulu and M. Tsingelis, *On left regular ordered semigroups*, South. Asian  
277 Bull. Math. **25** (2002) 609–615.
- 278 [8] N. Kehayopulu, *Ideals and Green's relations in ordered semigroups*, Int. J. Math.  
279 and Math. Sci. **2006** (1–8) Article ID 61286.
- 280 [9] N. Kehayopulu, *On completely regular poe-semigroups*, Math. Japonica **37** (1992)  
281 123–130.

282  
283  
284

Received  
Revised  
Accepted