

4 **MEDIAN FILTERS OF PSEUDO-COMPLEMENTED**
5 **DISTRIBUTIVE LATTICES**

6 M. SAMBASIVA RAO
7 *Department of Mathematics*
8 *MVGR College of Engineering, Vizianagaram*
9 *Andhra Pradesh, India-535005*
10 **e-mail:** mssraomaths35@rediffmail.com

11 **Abstract**

12 Coherent filters, strongly coherent filters, and τ -closed filters are intro-
13 duced in pseudo-complemented distributive lattices and their characteriza-
14 tion theorems are derived. A set of equivalent conditions is derived for every
15 filter of a pseudo-complemented distributive lattice to become a coherent
16 filter. The notion of median filters is introduced and some equivalent con-
17 ditions are derived for every maximal filter of a pseudo-complemented dis-
18 tributive lattice to become a median filter which leads to a characterization
19 of Stone lattices.

20 **Keywords:** Coherent filter; strongly coherent filter; median filter; minimal
21 prime filter; maximal filter; Stone lattice.

22 **2010 Mathematics Subject Classification:** 06D99.

23 **1. INTRODUCTION**

24 The theory of pseudo-complements in lattices, and particularly in distributive
25 lattices was developed by M.H. Stone [10], O. Frink [5], and George Gratzner [6].
26 Later many authors like R. Balbes [1], T.P. Speed [9], and O. Frink [5]etc., ex-
27 tended the study of pseudo-complements to characterize Stone lattices. In [3],
28 I. Chajda, R. Halaš and J. Kühr extensively studied the structure of pseudo-
29 complemented semilattices. In [7], the concept of δ -ideals was introduced in
30 pseudo-complemented distributive lattices and then Stone lattices were charac-
31 terized in terms of δ -ideals. In [8], the authors investigated the properties of
32 D -filters and prime D -filters of distributive lattices and characterized the mini-
33 mal prime D -filters of distributive lattices.

In this note, the concepts of coherent filters and strongly coherent filters are introduced in pseudo-complemented distributive lattices. A set of equivalent conditions is derived for every filter of a pseudo-complemented distributive lattice to become a coherent filter which characterizes a Boolean algebra. It is showed that every strongly coherent filter of a pseudo-complemented distributive lattice is coherent. The concepts of τ -closed filters and semi Stone lattices are introduced within pseudo-complemented distributive lattices and the class of all semi Stone lattices is characterized using the τ -closed filters. It is observed that the classes of maximal filters and prime D -filters coincide in a pseudo-complemented lattice. This observation precisely motivates to investigate the properties of certain class of filters under the name median filters as a special subclass of maximal filters of pseudo-complemented distributive lattices. Median filters are characterized and it is shown that every median filter of a pseudo-complemented distributive lattice is a coherent filter. A set of equivalent conditions is derived for every maximal filter of a pseudo-complemented distributive lattice to become a strongly coherent filter. Some equivalent conditions are stated for maximal filters of a pseudo-complemented distributive lattice to become median filters which leads to a characterization of Stone lattices.

2. PRELIMINARIES

The reader is referred to [2], [3] and [8] for the elementary notions and notations of pseudo-complemented distributive lattices. However some of the preliminary definitions and results are presented for the ready reference of the reader.

A non-empty subset A of a lattice L is called an *ideal (filter)* [2] of L if $a \vee b \in A$ ($a \wedge b \in A$) and $a \wedge x \in A$ ($a \vee x \in A$) whenever $a, b \in A$ and $x \in L$. The set $[a] = \{x \in L \mid x \leq a\}$ (resp. $[a] = \{x \in L \mid a \leq x\}$) is called a *principal ideal* (resp. *principal filter*) generated by a . The set $\mathcal{I}(L)$ of all ideals of a distributive lattice L with 0 forms a complete distributive lattice. The set $\mathcal{F}(L)$ of all filters of a distributive lattice L with 1 forms a complete distributive lattice in which $F \vee G = \{i \wedge j \mid i \in F \text{ and } j \in G\}$ for any two filters F and G . A proper filter P of a lattice L is said to be *prime* if for any $x, y \in L$, $x \vee y \in P$ implies $x \in P$ or $y \in P$. A proper filter P of a lattice L is called *maximal* if there exists no proper filter Q of L such that $P \subset Q$. A proper filter P of a distributive lattice is *minimal* if there exists no prime filter Q of L such that $Q \subset P$.

The *pseudo-complement* b^* of an element b is the element satisfying

$$a \wedge b = 0 \Leftrightarrow a \wedge b^* = a \Leftrightarrow a \leq b^*$$

where \leq is the induced order of L .

A distributive lattice L in which every element has a pseudo-complement is

71 called a *pseudo-complemented distributive lattice*. For any two elements a, b of a
 72 pseudo-complemented semilattice [3], we have the following.

- 73 (1) $a \leq b$ implies $b^* \leq a^*$,
- 74 (2) $a \leq a^{**}$,
- 75 (3) $a^{***} = a^*$,
- 76 (4) $(a \vee b)^* = a^* \wedge b^*$,
- 77 (5) $(a \wedge b)^{**} = a^{**} \wedge b^{**}$.

78 An element a of a pseudo-complemented distributive lattice L is called a *dense*
 79 *element* if $a^* = 0$ and the set D of all dense elements of L forms a filter in L .

80 **Definition** [2]. A pseudo-complemented distributive lattice L is called a *Stone*
 81 *lattice* if $x^* \vee x^{**} = 1$ for all $x \in L$

82 **Theorem 1** [2]. *The following conditions are equivalent in a pseudo-complemented*
 83 *distributive lattice L :*

- 84 (1) L is a Stone lattice;
- 85 (2) for $x, y \in L$, $(x \wedge y)^* = x^* \vee y^*$;
- 86 (3) for $x, y \in L$, $(x \vee y)^{**} = x^{**} \vee y^{**}$.

87 A filter F of a distributive lattice L is called a *D-filter* [8] if $D \subseteq F$. For any
 88 non-empty subset A of a distributive lattice L , the set $A^\circ = \{x \in L \mid x \vee a \in$
 89 $D \text{ for all } a \in A\}$ is a *D-filter* of L . In case of $A = \{a\}$, we simply represent $\{a\}^\circ$
 90 by $(a)^\circ$. A prime *D-filter* of a distributive lattice is minimal if it is the minimal
 91 element in the poset of all prime *D-filters*. A prime *D-filter* of a distributive
 92 lattice is minimal [8] if and only if to each $x \in P$, there exists $y \notin P$ such that
 93 $x \vee y \in D$. Throughout this note, all lattices are considered to be bounded
 94 pseudo-complemented distributive lattices unless otherwise mentioned.

95 3. COHERENT FILTERS

96 In this section, the concepts of coherent filters and strongly coherent filter are
 97 introduced. Stone lattices are characterized with help of coherent filters. A set of
 98 equivalent conditions is derived for every filter of a lattice to become a coherent
 99 filter which leads to a characterization of a Boolean algebra.

100 **Definition.** For any non-empty subset A of a lattice L , define

$$101 \quad A^\tau = \{x \in L \mid a^{**} \vee x^{**} = 1 \text{ for all } a \in A\}$$

102 Clearly $D^\tau = D$ and $L^\tau = D$. For any $a \in L$, we denote $(\{a\})^\tau$ simply by
 103 $(a)^\tau$. It is obvious that $(0)^\tau = D$ and $(1)^\tau = L$. For any $\emptyset \neq A \subseteq L$, $A \cap A^\tau \subseteq D$.

104 **Proposition 2.** *For any non-empty subset A of L , A^τ is a D -filter in L .*

105 **Proof.** Clearly $D \subseteq A^\tau$. Let $x, y \in A^\tau$. For any $a \in A$, we get $(x \wedge y)^{**} \vee a^{**} =$
 106 $(x^{**} \wedge y^{**}) \vee a^{**} = (x^{**} \vee a^{**}) \wedge (y^{**} \vee a^{**}) = 1 \wedge 1 = 1$. Hence $x \wedge y \in A^\tau$. Again,
 107 let $x \in A^\tau$ and $x \leq y$. Then $x^{**} \vee a^{**} = 1$ for any $a \in A$. Since $x \leq y$, we get
 108 $x^{**} \leq y^{**}$. For any $c \in A$, we get $1 = x^{**} \vee c^{**} \leq y^{**} \vee c^{**}$. Thus $y^{**} \vee c^{**} = 1$.
 109 Hence $y \in A^\tau$. Therefore A^τ is a D -filter of L . ■

110 The following lemma is a direct consequence of the above definition.

111 **Lemma 3.** *For any two non-empty subsets A and B of a lattice L , we have*

- 112 (1) $A \subseteq B$ implies $B^\tau \subseteq A^\tau$
- 113 (2) $A \subseteq A^{\tau\tau}$
- 114 (3) $A^{\tau\tau\tau} = A^\tau$
- 115 (4) $A^\tau = L$ if and only if $A = D$

116 **Proposition 4.** *For any two filters F, G of a lattice L , $(F \vee G)^\tau = F^\tau \cap G^\tau$.*

117 **Proof.** Clearly $(F \vee G)^\tau \subseteq F^\tau \cap G^\tau$. Conversely, let $x \in F^\tau \cap G^\tau$. Let $c \in F \vee G$
 118 be an arbitrary element. Then $c = i \vee j$ for some $i \in F$ and $j \in G$. Now
 119 $x^{**} \vee c^{**} = x^{**} \vee (i \vee j)^{**} = x^{**} \vee (i^{**} \wedge j^{**}) = (x^{**} \vee i^{**}) \wedge (x^{**} \vee j^{**}) = 1 \wedge 1 = 1$.
 120 Thus $x \in (F \vee G)^\tau$ and therefore $(F \vee G)^\tau = F^\tau \cap G^\tau$. ■

121 The following corollary is a direct consequence of the above results.

122 **Corollary 5.** *Let L be a lattice. For any $a, b \in L$, the following properties hold:*

- 123 (1) $a \leq b$ implies $(a)^\tau \subseteq (b)^\tau$,
- 124 (2) $(a \wedge b)^\tau = (a)^\tau \cap (b)^\tau$,
- 125 (3) $(a)^\tau = L$ if and only if a is dense,
- 126 (4) $a \in (b)^\tau$ implies $a \vee b \in D$,
- 127 (5) $a^* = b^*$ implies $(a)^\tau = (b)^\tau$.

128 Clearly $A^\tau \subseteq A^\circ$. We derive a set of equivalent conditions for a filter to
 129 satisfy the reverse inclusion which leads to a characterization of Stone lattices.

130 **Theorem 6.** *The following assertions are equivalent in a lattice L :*

- 131 (1) L is a Stone lattice;
- 132 (2) for any filter F of L , $F^\tau = F^\circ$;
- 133 (3) for any $a \in L$, $(a)^\tau = (a)^\circ$;
- 134 (4) for any two filters F, G of L , $F \cap G \subseteq D$ if and only if $F \subseteq G^\tau$;
- 135 (5) for $a, b \in L$, $a \vee b \in D$ implies $a^{**} \vee b^{**} = 1$;
- 136 (6) for $a \in L$, $(a)^{\tau\tau} = (a^*)^\tau$.

137 **Proof.** (1) \Rightarrow (2): Assume that L is a Stone lattice. Let F be a filter of L .
 138 Clearly $F^\tau \subseteq F^\circ$. Conversely, let $x \in F^\circ$. Then $x \vee y \in D$ for all $y \in F$. Since L
 139 is Stone, $x^{**} \vee y^{**} = (x \vee y)^{**} = 0^* = 1$ for all $y \in F$. Therefore $x \in F^\tau$.

140 (2) \Rightarrow (3): It is clear.

141 (3) \Rightarrow (4): Assume condition (3). Let F, G be two filters of L . Suppose $F \cap G \subseteq D$.
 142 Let $x \in F$. For any $y \in G$, we get $x \vee y \in F \cap G \subseteq D$. Hence $x \vee y \in D$. Now

$$\begin{aligned} x \vee y \in D \text{ for all } y \in G &\Rightarrow x \in (y)^\circ \text{ for all } y \in G \\ &\Rightarrow x \in (y)^\tau \text{ for all } y \in G \\ &\Rightarrow x^{**} \vee y^{**} = 1 \text{ for all } y \in G \end{aligned}$$

143 which yields that $x \in G^\tau$. Conversely, suppose that $F \subseteq G^\tau$. Let $x \in F \cap G$.
 144 Then $x \in F \subseteq G^\tau$ and $x \in G$. Hence $x \in G \cap G^\tau \subseteq D$. Therefore $F \cap G \subseteq D$.

145 (4) \Rightarrow (5): Assume condition (4). Let $a, b \in L$ be such that $a \vee b \in D$. Then

$$\begin{aligned} a \vee b \in D &\Rightarrow [a] \cap [b] \subseteq D \\ &\Rightarrow [a] \subseteq [b]^\tau \quad \text{by (4)} \\ &\Rightarrow a \in [b]^\tau \\ &\Rightarrow a^{**} \vee b^{**} = 1 \end{aligned}$$

146 (5) \Rightarrow (6): Assume condition (5). Let $a \in L$. Clearly, we have $a \vee a^* \in D$. By
 147 assumption (5), we get that $a^{**} \vee a^{***} = 1$. Hence $a^* \in (a)^\tau$. Thus $(a)^{\tau\tau} \subseteq (a^*)^\tau$.
 148 Conversely, let $x \in (a^*)^\tau$ and $t \in (a)^\tau$. Since $t \in (a)^\tau$, we get that $a^{**} \vee t^{**} = 1$.
 149 Hence $a^* \wedge t^* = 0$. Thus $t^* \leq a^{**}$. Now

$$\begin{aligned} x \in (a^*)^\tau &\Rightarrow a^* \vee x^{**} = 1 \\ &\Rightarrow a^{**} \wedge x^* = 0 \\ &\Rightarrow t^* \wedge x^* = 0 \quad \text{since } t^* \leq a^{**} \\ &\Rightarrow t \vee x \in D \\ &\Rightarrow t^{**} \vee x^{**} = 1 \quad \text{by (5)} \end{aligned}$$

150 which holds for all $t \in (a)^\tau$. Hence $x \in (a)^{\tau\tau}$. Therefore $(a^*)^\tau \subseteq (a)^{\tau\tau}$.

151 (6) \Rightarrow (1): Assume condition (6). Let $a \in L$. Since $a \in (a)^{\tau\tau} = (a^*)^\tau$, we get
 152 $a^* \vee a^{**} = a^{***} \vee a^{**} = 1$. Therefore L is a Stone lattice. \blacksquare

153 Now, we define coherent filters.

154 **Definition.** A filter F of a lattice L is called a *coherent filter* if for all $x, y \in$
 155 L , $(x)^\tau = (y)^\tau$ and $x \in F$ imply that $y \in F$.

156 Clearly each $(x)^\tau, x \in L$ is a coherent filter. It is evident that any filter F is
 157 a coherent filter if it satisfies $(x)^{\tau\tau} \subseteq F$ for all $x \in F$.

158 **Theorem 7.** *The following assertions are equivalent in a lattice L :*

- 159 (1) L is a Boolean algebra;
- 160 (2) every element is closed;
- 161 (3) for any filter F , $x^{**} \in F$ implies $x \in F$;
- 162 (4) every principal filter is a coherent filter;
- 163 (5) every filter is a coherent filter;
- 164 (6) every prime filter is a coherent filter;
- 165 (7) for $a, b \in L$, $(a)^\tau = (b)^\tau$ implies $a = b$;
- 166 (8) for $a, b \in L$, $a^* = b^*$ implies $a = b$.

167 **Proof.** (1) \Rightarrow (2): It is proved in [[7], Theorem 2.15].

168 (2) \Rightarrow (3): It is clear.

169 (3) \Rightarrow (4): Assume that every element of L is closed. Let $[x]$ be a principal filter
 170 of L . Since $x \vee x^* \in D$, we get $(x \vee x^*)^{**} = 1 \in [1]$. By (3), we get $x \vee x^* \in [1]$,
 171 which gives $x \vee x^* = 1$. Let $a, b \in L$ be such that $(a)^\tau = (b)^\tau$ and $a \in [x]$. Then

$$\begin{aligned}
 x \vee x^* = 1 &\Rightarrow a \vee x^* = 1 \quad \text{since } a \in [x] \\
 &\Rightarrow a^{**} \vee x^{***} = 1 \\
 &\Rightarrow x^* \in (a)^\tau = (b)^\tau \\
 &\Rightarrow b^{**} \vee x^* = 1 \\
 &\Rightarrow (b^{**} \vee x^*)^* = 0 \\
 &\Rightarrow b^* \wedge x^{**} = 0 \\
 &\Rightarrow b^* \wedge x = 0 \quad \text{since } x \leq x^{**} \\
 &\Rightarrow x \leq b^{**}
 \end{aligned}$$

172 which yields $b^{**} \in [x]$. By (3), we get $b \in [x]$. Hence $[x]$ is a coherent filter.

173 (4) \Rightarrow (5): Assume condition (4). Let F be a filter of L . Choose $a, b \in L$.
 174 Suppose $(a)^\tau = (b)^\tau$ and $a \in F$. Then clearly $[a] \subseteq F$. Since $(a)^\tau = (b)^\tau$ and $[a]$
 175 is a coherent filter, we get that $b \in [a] \subseteq F$. Therefore F is a coherent filter.

176 (5) \Rightarrow (6): It is clear.

177 (6) \Rightarrow (7): Assume that every prime filter of L is a coherent filter. Let $a, b \in L$
 178 such that $(a)^\tau = (b)^\tau$. Suppose $a \neq b$. Then there exists a prime filter P such
 179 that $a \in P$ and $b \notin P$. By the hypothesis, P is a coherent filter of L . Since
 180 $(a)^\tau = (b)^\tau$ and $a \in P$, we get $b \in P$, which is a contradiction. Therefore $a = b$.

181 (7) \Rightarrow (8): By Corollary 5(5), it is direct.

182 (8) \Rightarrow (1): Assume condition (8). Then L has a unique dense element. Therefore
 183 L is a Boolean algebra. ■

184 **Definition.** For any filter F of a lattice L , define $\pi(F)$ as follows:

$$\pi(F) = \{x \in L \mid (x)^\tau \vee F = L\}$$

The following lemma is an immediate consequence of the above definition.

Lemma 8. *For any two filters F, G of a lattice L , the following properties hold:*

- (1) $F \subseteq G$ implies $\pi(F) \subseteq \pi(G)$
- (2) $\pi(F \cap G) = \pi(F) \cap \pi(G)$.

Proof. Routine verification. ■

Proposition 9. *For any filter F of a lattice L , $\pi(F)$ is a D -filter of L .*

Proof. Clearly $D \subseteq \pi(F)$. Let $x, y \in \pi(F)$. Then $(x)^\tau \vee F = (y)^\tau \vee F = L$. Hence $(x \wedge y)^\tau \vee F = \{(x)^\tau \cap (y)^\tau\} \vee F = \{(x)^\tau \vee F\} \cap \{(y)^\tau \vee F\} = L$. Hence $x \wedge y \in \pi(F)$. Again let $x \in \pi(F)$ and $x \leq y$. Then $L = (x)^\tau \vee F \subseteq (y)^\tau \vee F$. Thus $y \in \pi(F)$. Therefore $\pi(F)$ is a D -filter in L . ■

Lemma 10. *Let F be a filter of a lattice L . Then F is a D -filter of L if and only if $\pi(F) \subseteq F$.*

Proof. Assume that F is a D -filter of L . Let $x \in \pi(F)$. Then $(x)^\tau \vee F = L$. Hence $x = a \wedge b$ for some $a \in (x)^\tau \subseteq (x)^\circ$ and $b \in F$. Then $x \vee a \in D \subseteq F$ and $x \vee b \in F$. Thus $x = x \vee x = x \vee (a \wedge b) = (x \vee a) \wedge (x \vee b) \in F$. Therefore $\pi(F) \subseteq F$. Converse is clear because of $D \subseteq \pi(F) \subseteq F$. ■

Definition. A filter F of a lattice L is called *strongly coherent* if $F = \pi(F)$.

Proposition 11. *Every strongly coherent filter of a lattice is a coherent filter.*

Proof. Let F be a strongly coherent filter of a lattice L . Clearly F is a D -filter of L . Let $x, y \in L$ be such that $(x)^\tau = (y)^\tau$ and $x \in F = \pi(F)$. Then $(x)^\tau \vee F = L$. Hence $(y)^\tau \vee F = L$ and so $y \in \pi(F) = F$. Thus F is a coherent filter of L . ■

For any filter F of a lattice L , it can be noted that $F \subseteq D$ if and only if $F^{\tau\tau} = D$. A D -filter F of a lattice L is called a τ -closed if $F = F^{\tau\tau}$. Clearly D is the smallest τ -closed filter and L is the largest τ -closed filter of the lattice L .

Proposition 12. *Every τ -closed filter of a lattice is a coherent filter.*

Proof. Let F be a τ -closed filter of a lattice L . Let $x, y \in L$ be such that $(x)^\tau = (y)^\tau$. Suppose $x \in F$. Then, we get that $y \in (y)^{\tau\tau} = (x)^{\tau\tau} \subseteq F^{\tau\tau} = F$. Therefore F is a coherent filter of L . ■


Definition. A lattice L is called a *semi Stone lattice* if $(x)^\tau \vee (x)^{\tau\tau} = L$ for all $x \in L$.

216 **Theorem 13.** *Every Stone lattice is a semi Stone lattice.*

217 **Proof.** Assume that L is a Stone lattice. Let $x \in L$. Suppose $(x)^\tau \vee (x)^{\tau\tau} \neq L$.
 218 Then there exists a maximal filter M such that $(x)^\tau \vee (x)^{\tau\tau} \subseteq M$. Then $(x)^\tau \subseteq M$
 219 and $x \in (x)^{\tau\tau} \subseteq M$. Since M is maximal, we get $x^* \notin M$. Since L is Stone, we
 220 get $x^* \vee x^{**} = 1$. Hence $x^* \in (x)^\tau$. Thus $(x)^\tau \not\subseteq M$, which is a contradiction.
 221 Hence $(x)^\tau \vee (x)^{\tau\tau} = L$. Therefore L is a semi Stone lattice. ■

222 The converse of the above theorem is not true. For, consider

223 **Example 14.** Consider the following bounded and finite distributive lattice
 224 $L = \{0, a, b, c, 1\}$ whose Hasse diagram is given by:



225
 226
 227 Clearly L is a pseudo-complemented lattice. It can be easily observed that $(a)^\tau =$
 228 $(b)^\tau = D$ and $(c)^\tau = (1)^\tau = L$. Hence $(a)^{\tau\tau} = (b)^{\tau\tau} = L$. Observe that L is a
 229 semi Stone lattice. But L is not a Stone lattice because of $a^* \vee a^{**} = b \vee a = c \neq 1$.

230 **Theorem 15.** *The following assertions are equivalent in a lattice L :*

- 231 (1) L is a semi Stone lattice;
- 232 (2) every τ -closed filter is strongly coherent;
- 233 (3) for each $x \in L$, $(x)^{\tau\tau}$ is strongly coherent.

234 **Proof.** (1) \Rightarrow (2): Assume that L is a semi Stone lattice. Let F be a τ -closed
 235 filter of L . Then F is a D -filter with $F^{\tau\tau} = F$. Clearly $\pi(F) \subseteq F$. Conversely,
 236 let $x \in F$. It can be easily verified that $(x)^{\tau\tau} \subseteq F^{\tau\tau}$. Hence $L = (x)^\tau \vee (x)^{\tau\tau} \subseteq$
 237 $(x)^\tau \vee F^{\tau\tau} = (x)^\tau \vee F$. Thus $x \in \pi(F)$. Therefore F is strongly coherent.

238 (2) \Rightarrow (3): Since each $(x)^{\tau\tau}$ is τ -closed, it is obvious.

239 (3) \Rightarrow (1): Assume condition (3). Let $x \in L$. Then we get $\pi((x)^{\tau\tau}) = (x)^{\tau\tau}$.
 240 Since $x \in (x)^{\tau\tau}$, we get $(x)^\tau \vee (x)^{\tau\tau} = L$. Therefore L is a semi Stone lattice. ■

241

4. MEDIAN FILTERS

242 In this section, the notion of a median filter is introduced in lattices. Char-
 243 acterization theorems of median filters are derived for every prime D -filter to
 244 become median and every maximal filter to become median.

245 **Proposition 16.** *Let P be a prime filter of a lattice L . Then the following*
 246 *assertions are equivalent:*

- 247 (1) $D \subseteq P$;
- 248 (2) for any $x \in L$, $x \in P$ if and only if $x^* \notin P$;
- 249 (3) for any $x \in L$, $x^{**} \in P$ if and only if $x \in P$;
- 250 (4) for any $x, y \in L$ with $x^* = y^*$, $x \in P$ implies that $y \in P$;
- 251 (5) $D \cap (L - P) = \emptyset$.

252 **Proof.** (1) \Rightarrow (2): Assume that $D \subseteq P$. Suppose $x \in P$. If $x^* \in P$, then
 253 $0 = x \wedge x^* \in P$, which is a contradiction. Hence $x^* \notin P$. Conversely, let $x^* \notin P$.
 254 Clearly $x \vee x^* \in D \subseteq P$. Since P is prime and $x^* \notin P$, we get $x \in P$.

255 (2) \Rightarrow (3): It is clear.

256 (3) \Rightarrow (4): It is clear.

257 (4) \Rightarrow (5): Assume condition (4). Suppose $x \in D \cap (L - P)$. Then, we get
 258 $x^* = 0 = 1^*$ and $x \notin P$. Since $1 \in P$, by (4), we get that $x \in P$ which is a
 259 contradiction. Therefore $D \cap (L - P) = \emptyset$.

260 (5) \Rightarrow (1): It is obvious. ■

261 **Theorem 17.** *Let M be a proper filter of a lattice L . The following assertions*
 262 *are equivalent:*

- 263 (1) M is maximal;
- 264 (2) M is a prime D -filter;
- 265 (3) $x \notin M$ implies $x^* \in M$.

266 **Proof.** (1) \Rightarrow (2): Assume that M is a maximal filter of L . Clearly M is a prime
 267 filter. Let $x \in D$. Then, we get $x^* = 0$. Suppose $x \notin M$. Then $M \vee [x] = L$.
 268 Hence $0 = m \wedge x$ for some $0 \neq m \in M$. Then $m \leq x^* = 0$, which is a contradiction.
 269 Hence $x \in M$. Thus $D \subseteq M$. Therefore M is a prime D -filter.

270 (2) \Rightarrow (3): Assume that M is a prime D -filter of L . Suppose $x \notin M$. Clearly
 271 $x \vee x^* \in D \subseteq M$. Since M is prime and $x \notin M$, we have $x^* \in M$.

272 (3) \Rightarrow (1): Assume condition (3). Suppose M is not maximal. Let Q be a proper
 273 filter such that $M \subset Q$. Choose $x \in Q - M$. Since $x \notin M$, by (3), we get that
 274 $x^* \in M \subset Q$. Therefore $0 = x \wedge x^* \in Q$, which is a contradiction. ■

275 From Theorem 17, one can notice that the class of all maximal filters and
 276 the class of all prime D -filters are the same. Since every prime D -filter is max-
 277 imal, we can conclude that every prime D -filter is minimal. Therefore maximal
 278 filters, prime D -filter, and minimal prime D -filters are the same in a pseudo-
 279 complemented distributive lattice. The notion of median filters is now introduced.

280 **Definition.** A maximal filter M of a lattice L is called *median* if to each $x \in M$,
 281 there exists $y \notin M$ such that $x^{**} \vee y^{**} = 1$.

From Example 14, we initially observe that a maximal filter of a lattice need not be median, For consider the maximal filter $M = \{1, a, c\}$ of L . Notice that for $a \in M$, there is no $x \notin M$ such that $a^{**} \vee x^{**} = 1$. Therefore M is not median.

Lemma 18. *Let M be a maximal filter of a lattice L . For any $x \in L$, it holds $x \notin M$ implies $(x)^\tau \subseteq M$*

Proof. Suppose $x \notin M$. Let $a \in (x)^\tau$. Then $a^{**} \vee x^{**} = 1$. Hence $(a \vee x)^{**} = 1$ and so $a \vee x \in D \subseteq M$. Since $x \notin M$, we get $a \in M$. Therefore $(x)^\tau \subseteq M$. ■

Lemma 19. *Let M be a median filter of a lattice L . For any $x \in L$, we have $x \in M$ if and only if $(x)^{\tau\tau} \subseteq M$*

Proof. Suppose that $x \in M$. Let $a \in (x)^{\tau\tau}$. Then, we get $(x)^\tau \subseteq (a)^\tau$. Since $x \in M$ and M is a median filter, there exists $y \notin M$ such that $x^{**} \vee y^{**} = 1$. Then, we get $y \in (x)^\tau \subseteq (a)^\tau$. Since $y \notin M$, we must have $(y)^\tau \subseteq M$. Hence $a \in (a)^{\tau\tau} \subseteq (y)^\tau \subseteq M$. Therefore $(x)^{\tau\tau} \subseteq M$. ■

In the following, we derive a characterization theorem of median filters.

Theorem 20. *Let M be a maximal filter of a lattice L . For each $x \in L$, the following assertions are equivalent:*

- (1) M is median;
- (2) $x \notin M$ if and only if $(x)^\tau \subseteq M$;
- (3) $x^{**} \in M$ implies $(x)^\tau \not\subseteq M$.

Proof. (1) \Rightarrow (2): Assume that M is a median filter of L and $x \in L$. Suppose $x \notin M$. By Lemma 18, we have $(x)^\tau \subseteq M$. Conversely, assume that $(x)^\tau \subseteq M$. Suppose $x \in M$. Since M is median, there exists $y \notin M$ such that $x^{**} \vee y^{**} = 1$. Hence $y \in (x)^\tau \subseteq M$, which is a contradiction. Therefore $x \notin M$.

(2) \Rightarrow (3): Assume condition (2). Let $x \in L$. Suppose $x^{**} \in M$. By Proposition 16, we get $x \in M$. By (2), we get $(x)^\tau \not\subseteq M$.

(3) \Rightarrow (1): Assume that condition (3) holds. Suppose $x \in M$. Clearly $x^{**} \in M$. By the assumed condition, we get that $(x)^\tau \not\subseteq M$. Then there exists $y \in (x)^\tau$ such that $y \notin M$. Hence $x^{**} \vee y^{**} = 1$ where $y \notin M$. Therefore M is median. ■

Theorem 21. *Every median filter of a lattice is a coherent filter.*

Proof. Let M be a median filter of a lattice L . Suppose $x, y \in L$ be such that $(x)^\tau = (y)^\tau$ and $x \in M$. Since M is median, there exists $a \notin M$ such that $x^{**} \vee a^{**} = 1$. Hence $a \in (x)^\tau = (y)^\tau$. Thus $1 = y^{**} \vee a^{**} \leq (y \vee a)^{**}$. Hence $(y \vee a)^{**} = 1$, which gives that $(y \vee a)^* = 0$. Thus $y \vee a \in D \subseteq M$. Since M is prime and $a \notin M$, it yields that $y \in M$. Therefore M is a coherent filter. ■

In the following theorem, we derive a set of equivalent conditions for a Stone lattice to become a Boolean algebra in terms of median filters and maximal filters.

Theorem 22. *Let L be a Stone lattice. Then the following are equivalent:*

- (1) L is a Boolean algebra;
- (2) every prime filter is maximal;
- (3) every prime filter is median;
- (4) every prime filter is a D -filter.

Proof. (1) \Rightarrow (2): It is well known.

(2) \Rightarrow (3): Since L is a Stone lattice, it is through.

(3) \Rightarrow (4): Since every median filter is a D -filter, it is clear.

(4) \Rightarrow (1): Assume condition (3). Then $D \subseteq \bigcap \{P \mid P \text{ is a prime filter}\} = \{1\}$. Hence $D = \{1\}$, which gives $x \vee x^* \in D = \{1\}$. Thus it is through. ■

Definition. For any maximal filter M of a lattice L , define

$$\Omega(M) = \{x \in L \mid (x)^\tau \not\subseteq M\}.$$

Lemma 23. *For any maximal filter M , $\Omega(M)$ is a D -filter contained in M .*

Proof. Clearly $D \subseteq \Omega(M)$. Let $x, y \in \Omega(M)$. Then $(x)^\tau \not\subseteq M$ and $(y)^\tau \not\subseteq M$. Since M is prime, we get $(x \wedge y)^\tau = (x)^\tau \cap (y)^\tau \not\subseteq M$. Hence $x \wedge y \in \Omega(M)$. Let $x \in \Omega(M)$ and $x \leq y$. Then $(x)^\tau \not\subseteq M$ and $(x)^\tau \subseteq (y)^\tau$. Since $(x)^\tau \not\subseteq M$, we get $(y)^\tau \not\subseteq M$. Thus $y \in \Omega(M)$. Therefore $\Omega(M)$ is a D -filter of L . Now, let $x \in \Omega(M)$. Then, we get $(x)^\tau \not\subseteq M$. Hence there exists $a \in (x)^\tau$ such that $a \notin M$. Since $a \in (x)^\tau$, we get $1 = a^{**} \vee x^{**} \leq (a \vee x)^{**}$. Thus $a \vee x \in D \subseteq M$. Since $a \notin M$, we must have $x \in M$. Therefore $\Omega(M) \subseteq M$. ■

Let us denote that \mathcal{M} is the set of all maximal filters of a lattice L . For any $a \in L$, we also denote $\mathcal{M}_{a^*} = \{M \in \mathcal{M} \mid a^* \in M\}$.

Theorem 24. *Let L be a lattice and $a \in L$. Then $(a)^\tau \subseteq \bigcap_{M \in \mathcal{M}_{a^*}} \Omega(M)$.*

Proof. Let $x \in (a)^\tau$ and $M \in \mathcal{M}_{a^*}$. Then $x^{**} \vee a^{**} = 1$ and $a^* \in M$. Suppose $a \in M$. Then $0 = a \wedge a^* \in M$, which is a contradiction. Hence $a \notin M$. Hence $a \in (x)^\tau$ such that $a \notin M$. Thus $(x)^\tau \not\subseteq M$. Hence $x \in \Omega(M)$. Thus $(a)^\tau \subseteq \Omega(M)$ which is true for all $M \in \mathcal{M}_{a^*}$. Therefore $(a)^\tau \subseteq \bigcap_{M \in \mathcal{M}_{a^*}} \Omega(M)$. ■

Corollary 25. *Let L be a lattice and $a \in L$. Then $a^* \in M$ implies $(a)^\tau \subseteq \Omega(M)$.*

In Example 14, consider $P = \{1, a, c\}$. Clearly $D = \{1, c\}$ and P is a prime D -filter. For any element $x \in P$, there exists no $y \notin P$ such that $x^{**} \vee y^{**} = 1$. Hence P is not median. However, in the following, some equivalent conditions are derived for every prime D -filter of a lattice to become a median filter.

350 **Theorem 26.** *The following conditions are equivalent in a lattice L :*

- 351 (1) L is a Stone lattice;
- 352 (2) every D -filter is strongly coherent;
- 353 (3) every maximal filter is strongly coherent;
- 354 (4) every maximal filter is median;
- 355 (5) for any $M \in \mathcal{M}$, $\Omega(M)$ is median;
- 356 (6) for any $a, b \in L$, $a \vee b \in D$ implies $(a)^\tau \vee (b)^\tau = L$;
- 357 (7) for any $a \in L$, $(a)^\tau \vee (a^*)^\tau = L$.

358 **Proof.** (1) \Rightarrow (2): Assume that L is a Stone lattice. Let F be a D -filter of L .
 359 Clearly $\pi(F) \subseteq F$. Conversely, let $x \in F$. Since L is a Stone lattice, we get
 360 $x^* \vee x^{**} = 1$. Suppose $(x)^\tau \vee F \neq L$. Then there exists a maximal filter M of
 361 L such that $(x)^\tau \vee F \subseteq M$. Hence $(x)^\tau \subseteq M$ and $x \in F \subseteq M$. Since M is a
 362 prime, we get $x^* \notin M$. Since $x^{**} \vee x^* = 1$, we get $x^* \in (x)^\tau \subseteq M$ which is a
 363 contradiction. Thus $(x)^\tau \vee F = L$. Therefore F is a strongly coherent filter.

364 (2) \Rightarrow (3): It is obvious.

365 (3) \Rightarrow (4): Assume that every maximal filter is strongly coherent. Let M be a
 366 maximal filter of L . Then by our assumption, $\pi(M) = M$. Let $x \in M$. Then
 367 $(x)^\tau \vee M = L$. Hence $a \wedge b = 0$ for some $a \in (x)^\tau$ and $b \in M$. Since $a \in (x)^\tau$, we
 368 get $a^{**} \vee x^* = 1$. Suppose $a \in M$. Then $0 = a \wedge b \in M$, which is a contradiction.
 369 Hence $a \notin M$. Therefore M is median.

370 (4) \Rightarrow (5): Assume condition (4). Let $M \in \mathcal{M}$. Clearly $\Omega(M) \subseteq M$. Conversely,
 371 let $x \in M$. Since M is median, there exists $y \notin M$ such that $x^{**} \vee y^* = 1$. Hence
 372 $(x)^\tau \not\subseteq M$. Thus $x \in \Omega(M)$. Therefore $\Omega(M) = M$ is a median filter.

373 (5) \Rightarrow (6) : Assume condition (5). Let $a, b \in L$ be such that $a \vee b \in D$. Suppose
 374 $(a)^\tau \vee (b)^\tau \neq L$. Then there exists a maximal filter M such that $(a)^\tau \vee (b)^\tau \subseteq M$.
 375 Since $\Omega(M)$ is median, by Theorem 20, we get

$$\begin{aligned}
 (a)^\tau \vee (b)^\tau \subseteq M &\Rightarrow (a)^\tau \subseteq M \text{ and } (b)^\tau \subseteq M \\
 &\Rightarrow (a)^\tau \subseteq \Omega(M) \text{ and } (b)^\tau \subseteq \Omega(M) \\
 &\Rightarrow a \notin \Omega(M) \text{ and } b \notin \Omega(M) \quad \text{since } \Omega(M) \text{ is median} \\
 &\Rightarrow a \vee b \notin M
 \end{aligned}$$

376 which is a contradiction to that $a \vee b \in D \subseteq M$. Therefore $(a)^\tau \vee (b)^\tau = L$.

377 (6) \Rightarrow (7): Let $a \in L$. Since $a \vee a^* \in D$, by (6), we are through.

378 (7) \Rightarrow (1): Assume condition (7). Let $x \in L$. Then by (7), we have $(x)^\tau \vee (x^*)^\tau =$
 379 L . Hence $0 \in (x)^\tau \vee (x^*)^\tau$. Then $0 = a \wedge b$ for some $a \in (x)^\tau$ and $b \in (x^*)^\tau$. Since
 380 $b \in (x^*)^\tau$, we get $b^{**} \vee x^* = 1$, and so $b^* \wedge x^{**} = 0$. Thus $b^* \leq x^*$. Now

$$1 = a^{**} \vee x^{**} \quad \text{since } a \in (x)^\tau$$

$$\begin{aligned} &\leq b^* \vee x^{**} && \text{since } a \wedge b = 0 \\ &\leq x^* \vee x^{**} && \text{since } b^* \leq x^* \end{aligned}$$

381 which gives that $x^* \vee x^{**} = 1$. Therefore L is a Stone lattice. \blacksquare

382 For any filter F of a lattice L , we denote $\mathcal{M}_F = \{M \in \mathcal{M} \mid F \subseteq M\}$.

383 **Theorem 27.** For any filter F of a lattice L , $\pi(F) = \bigcap_{M \in \mathcal{M}_F} \Omega(M)$.

384 **Proof.** Let $x \in \pi(F)$ and $F \subseteq M$ where $M \in \mathcal{M}$. Then $L = (x)^\tau \vee F \subseteq (x)^\tau \vee M$.
 385 Suppose $(x)^\tau \subseteq M$, then $M = L$, which is a contradiction. Hence $(x)^\tau \not\subseteq M$.
 386 Thus $x \in \Omega(M)$ for all $M \in \mathcal{M}_F$. Therefore $\pi(F) \subseteq \bigcap_{M \in \mathcal{M}_F} \Omega(M)$.

387 Conversely, let $x \in \bigcap_{M \in \mathcal{M}_F} \Omega(M)$. Then, we get $x \in \Omega(M)$ for all $M \in \mathcal{M}_F$.
 388 Suppose $(x)^\tau \vee F \neq L$. Then there exists a maximal filter M_0 such that $(x)^\tau \vee F \subseteq$
 389 M_0 . Hence $(x)^\tau \subseteq M_0$ and $F \subseteq M_0$. Since $F \subseteq M_0$, by hypothesis, we get
 390 $x \in \Omega(M_0)$. Thus $(x)^\tau \not\subseteq M_0$, which is a contradiction. Therefore $(x)^\tau \vee F = L$.
 391 Thus $x \in \pi(F)$. Therefore $\bigcap_{M \in \mathcal{M}_F} \Omega(M) \subseteq \pi(F)$. \blacksquare

392 **Theorem 28.** The following assertions are equivalent in a lattice L :

- 393 (1) L is a Stone lattice;
- 394 (2) for any $M \in \mathcal{M}$, $\Omega(M)$ is maximal;
- 395 (3) for any $F, G \in \mathcal{F}(L)$, $F \vee G = L$ implies $\pi(F) \vee \pi(G) = L$;
- 396 (4) for any $F, G \in \mathcal{F}(L)$, $\pi(F) \vee \pi(G) = \pi(F \vee G)$;
- 397 (5) for any two distinct maximal filters M, N , $\Omega(M) \vee \Omega(N) = L$;
- 398 (6) for any $M \in \mathcal{M}$, M is the unique member of μ such that $\Omega(M) \subseteq M$.

399 **Proof.** (1) \Rightarrow (2) : Assume that L is a Stone lattice. Let $M \in \mathcal{M}$. Clearly,
 400 we have $\Omega(M) \subseteq M$. Conversely, let $x \in M$. Since L is a Stone lattice, By
 401 Theorem 26, we get that M is a median filter. Then there exists $y \notin M$ such that
 402 $x^{**} \vee y^{**} = 1$. Hence $y \in (x)^\tau$ and $y \notin M$. Thus $(x)^\tau \not\subseteq \Omega(M)$. Hence $x \in \Omega(M)$.
 403 Therefore $\Omega(M) = M$ is a maximal filter.

404 (2) \Rightarrow (3) : Assume condition (2). Clearly $\Omega(M) = M$ for all $M \in \mathcal{M}$. Let
 405 $F, G \in \mathcal{F}(L)$ be such that $F \vee G = L$. Suppose $\pi(F) \vee \pi(G) \neq L$. Then there
 406 exists a maximal filter M of L such that $\pi(F) \vee \pi(G) \subseteq M$. Hence $\pi(F) \subseteq M$
 407 and $\pi(G) \subseteq M$. Now, we get

$$\begin{aligned} \pi(F) \subseteq M &\Rightarrow \bigcap_{M \in \mathcal{M}_F} \Omega(M) \subseteq M \\ &\Rightarrow \Omega(M_i) \subseteq M \text{ for some } M_i \in \mathcal{M}_F \text{ (since } M \text{ is prime)} \\ &\Rightarrow M_i \subseteq M \quad \text{By condition (2)} \\ &\Rightarrow F \subseteq M \end{aligned}$$

408 Similarly, we can get $G \subseteq M$. Hence $L = F \vee G \subseteq M$, which is a contradiction.
 409 Therefore $\pi(F) \vee \pi(G) = L$.

410 (3) \Rightarrow (4) : Assume condition (3). Let $F, G \in \mathcal{F}(L)$. Clearly, we have $\pi(F) \vee$
 411 $\pi(G) \subseteq \pi(F \vee G)$. Let $x \in \pi(F \vee G)$. Then $((x)^\tau \vee F) \vee ((x)^\tau \vee G) = (x)^\tau \vee F \vee G =$
 412 L . Hence by condition (3), we get that $\pi((x)^\tau \vee F) \vee \pi((x)^\tau \vee G) = L$. Thus
 413 $x \in \pi((x)^\tau \vee F) \vee \pi((x)^\tau \vee G)$. Hence $x = r \wedge s$ for some $r \in \pi((x)^\tau \vee F)$ and
 414 $s \in \pi((x)^\tau \vee G)$. Now, we have

$$\begin{aligned} r \in \pi((x)^\tau \vee F) &\Rightarrow (r)^\tau \vee (x)^\tau \vee F = L \\ &\Rightarrow L = ((r)^\tau \vee (x)^\tau) \vee F \subseteq (r \vee x)^\tau \vee F \\ &\Rightarrow (r \vee x)^\tau \vee F = L \\ &\Rightarrow r \vee x \in \pi(F) \end{aligned}$$

415 Similarly, we can get $s \vee x \in \pi(G)$. Hence

$$\begin{aligned} x &= x \vee x \\ &= x \vee (r \wedge s) \\ &= (x \vee r) \wedge (x \vee s) \in \pi(F) \vee \pi(G) \end{aligned}$$

416 Hence $\pi(F \vee G) \subseteq \pi(F) \vee \pi(G)$. Therefore $\pi(F) \vee \pi(G) = \pi(F \vee G)$.

417 (4) \Rightarrow (5) : Assume condition (4). Let M, N be two distinct maximal filters of
 418 L . Choose $x \in M - N$ and $y \in N - M$. Since $x \notin N$ and $y \notin M$, we get $x^* \in N$
 419 and $y^* \in M$. Hence $(x \wedge y^*) \wedge (y \wedge x^*) = (x \wedge x^*) \wedge (y \wedge y^*) = 0$. Then

$$\begin{aligned} L &= \pi(L) \\ &= \pi([0]) \\ &= \pi([(x \wedge y^*) \wedge (y \wedge x^*)]) \\ &= \pi([x \wedge y^*] \vee [y \wedge x^*]) \\ &= \pi([x \wedge y^*]) \vee \pi([y \wedge x^*]) \quad \text{By condition (4)} \\ &\subseteq \Omega(M) \vee \Omega(N) \quad \text{since } [x \wedge y^*] \subseteq M, [y \wedge x^*] \subseteq N \end{aligned}$$

420 Therefore $\Omega(M) \vee \Omega(N) = L$.

421 (5) \Rightarrow (6) : Assume condition (5). Let $M \in \mathcal{M}$. Clearly $\Omega(M) \subseteq M$. Suppose
 422 $N \in \mathcal{M}$ such that $N \neq M$ and $\Omega(M) \subseteq N$. Since $\Omega(N) \subseteq N$, by hypothesis, we
 423 get $L = \Omega(M) \vee \Omega(N) \subseteq N$, which is a contradiction. Therefore M is the unique
 424 maximal filter of L such that $\Omega(M)$ is contained in M .

425 (6) \Rightarrow (1) : Let M be a maximal filter of L . Suppose $\Omega(M) \neq M$. Then there
 426 exists a maximal filter M_0 such that $\Omega(M) \subseteq M_0$, which contradicts uniqueness
 427 of M . Hence $\Omega(M) = M$. Let $x \in M = \Omega(M)$. Then there exists $y \notin M$ such
 428 that $x^{**} \vee y^{**} = 1$. Hence M is median. By Theorem 26, L is a Stone lattice. ■

Acknowledgements: The author would like to thank the referee for his valuable suggestions and comments.

REFERENCES

- [1] R. Balbes and A. Horn, *Stone lattices*, Duke Math. Journal **37** (1970) 537–545.
- [2] G. Birkhoff, *Lattice Theory* (Amer. Math. Soc. Colloq. XXV, Providence, U.S.A, 1967).
- [3] I. Chajda, R. Halaš and J. Kühr, *Semilattice structures* (Heldermann Verlag, Germany, ISBN 978-3-88538-230-0, 2007).
- [4] W.H. Cornish, *Congruences on distributive pseudo-complemented lattices*, Bull. Austral. Math. Soc. **8** (1973) 167–179.
- [5] O. Frink, *Pseudo-complements in semi-lattices*, Duke Math. Journal **29** (1962) 505–514. doi: 10.1215/S0012-7094-62-02951-4.
- [6] G. Gratzner, *General lattice theory* (Academic press, New york, San Francisco, U.S.A., 1978).
- [7] M. Sambasiva Rao, *δ -ideals in pseudo-complemented distributive lattices*, Archivum Mathematicum **48(2)** (2012) 97–105. doi:10.5817/AM2012-2-97.
- [8] A.P. Paneendra Kumar, M. Sambasiva Rao, and K. Sobhan Babu, *Generalized prime D -filters of distributive lattices*, Archivum Mathematicum **57(3)** (2021) 157–174. doi: 10.5817/AM2021-3-157.
- [9] T.P. Speed, *On Stone lattices*, Jour. Aust. Math. Soc. **9(3-4)** (1969) 297–307. doi: <https://doi.org/10.1017/S1446788700007217>
- [10] M.H. Stone, *A theory of representations for Boolean algebras*, Tran. Amer. Math. Soc. **40** (1936) 37–111. doi:<https://doi.org/10.2307/1989664>.