

A NOTE ON INTRA REGULARITY ON SEMIGROUPS OF PARTIAL TRANSFORMATIONS WITH INVARIANT SET

THAPAKORN PANTARAK AND YANISA CHAIYA

Department of Mathematics and Statistics
Faculty of Science and Technology
Thammasat University, Pathum Thani, 12120, Thailand

e-mail: thapakorn.pan@dome.tu.ac.th
yanisa@mathstat.sci.tu.ac.th

Abstract

Let X be any non-empty set and $P(X)$ denote the semigroup (under the composition of functions) of partial transformations on a set X . Let Y be a fixed non-empty subset of X and

$$\overline{PT}(X, Y) = \{\alpha \in P(X) : (\text{dom } \alpha \cap Y)\alpha \subseteq Y\}.$$

Then $\overline{PT}(X, Y)$ is a semigroup consisting of all mappings in $P(X)$ that leave $Y \subseteq X$ invariant. In this paper, we present criteria for checking the intra-regularity of elements in $\overline{PT}(X, Y)$ and apply these results to quantify intra-regular elements in $\overline{PT}(X, Y)$, when X is finite.

Keywords: partial transformation semigroup, intra regularity, invariant set.

2020 Mathematics Subject Classification: 20M20.

1. INTRODUCTION AND PRELIMINARIES

Let S be a semigroup. An element a of S is said to be *intra-regular* if there exist $x, y \in S$, such that $a = xa^2y$. The notion of intra regularity was introduced in Croisot's theory of decompositions [2, Section 4.1].

Let X be a non-empty set and $T(X)$ denote the semigroup (under the composition of mappings) of all transformations from X into itself. It is known as *full transformation semigroup*. The study of algebraic properties on semigroups in such types was started by Doss [4] in 1955. The author completely described its Green's relations. Particularly, a characterization of \mathcal{J} -relation can be directly

used to identify intra regularity, because $\alpha \in T(X)$ is intra-regular if and only if $\alpha \mathcal{J} \alpha^2$. Several other properties of $T(X)$ have been researched extensively by the fact that any semigroup can be embedded in $T(X)$ for some an appropriate set X .

Let Y be a non-empty subset of X . In 1966, Magill [7] introduced a subsemigroup of $T(X)$, defined by

$$\overline{T}(X, Y) = \{\alpha \in T(X) \mid Y\alpha \subseteq Y\}.$$

In addition, $\overline{T}(X, Y)$ is a generalization of $T(X)$, since $\overline{T}(X, X) = T(X)$. This fact inspired Homyam and Sanwong [5] to give a complete description of Green's relations on $\overline{T}(X, Y)$. Later in 2013, Choomanee *et al.* [1] used these results to provide characterization and number of intra-regular elements on $\overline{T}(X, Y)$.

For any non-empty set X , the super semigroup of all transformation semigroups on X is a *partial transformation semigroup*, which is defined by

$$P(X) = \{\alpha : A \rightarrow X \mid A \subseteq X\}.$$

Its Green's relations was shown in [6]. Similarly, the characterization of intra-regular was explored immediately.

For a fixed non-empty subset Y of X , in analogy with $\overline{T}(X, Y)$, consider

$$\overline{PT}(X, Y) = \{\alpha \in P(X) \mid (\text{dom } \alpha \cap Y)\alpha \subseteq Y\},$$

where $\text{dom } \alpha$ and $Y\alpha$ denote the domain of α and $(\text{dom } \alpha \cap Y)\alpha$, respectively. Since $\overline{PT}(X, X) = P(X)$, we may regard $\overline{PT}(X, Y)$ as a generalization of $P(X)$. Note that id_X , the identity map on X , belongs to $\overline{PT}(X, Y)$.

We now provide important preliminaries for this paper. Some basic mathematical terminologies and relevant notations used in what follows on semigroups are prescribed. Further, we refer to [2, 3, 6] for more information. Indeed, throughout this paper, the functions are written on the right, i.e., in the composition $\alpha\beta$, α is applied first. For any $\alpha \in P(X)$, the notations $\text{dom } \alpha$ and $\text{im } \alpha$ denote the *domain* of α and the *range* of α , respectively. Additionally, for any $x \in \text{im } \alpha$, $x\alpha^{-1}$ denotes the set of inverse images of x under α , i.e., $x\alpha^{-1} = \{z \in \text{dom } \alpha : z\alpha = x\}$. In addition, the following notation is applied

$$\alpha = \begin{pmatrix} X_i \\ a_i \end{pmatrix}.$$

Here, the script i belongs to some (unmentioned) index set I , the abbreviation $\{a_i\}$ denotes $\{a_i : i \in I\}$ and that $\text{im } \alpha = \{a_i\}$ and $a_i\alpha^{-1} = X_i \subseteq \text{dom } \alpha$ where $\bigcup_{i \in I} X_i = \text{dom } \alpha$. More specifically, when $\alpha \in \overline{PT}(X, Y)$, we have $Y\alpha \subseteq Y$. Thus, the domain of α can be divided into three parts as follows

$$\alpha = \begin{pmatrix} A_i & B_j & C_k \\ a_i & b_j & c_k \end{pmatrix},$$

where $A_i \cap Y \neq \emptyset, B_j, C_k \subseteq X \setminus Y; a_i, b_j \in Y$ and $c_k \in X \setminus Y$. Here, I, J and K can be empty.

In this paper, we describe the necessary and sufficient conditions for elements of $\overline{PT}(X, Y)$ to be intra-regular. The results recapture the known results on $P(X)$ when we focus on $Y = X$. Moreover, they are used to deduce the results for $T(X, Y)$, when elements with X as their domain are considered. We also apply the results to quantify the intra-regular elements in the $\overline{PT}(X, Y)$ when X is a finite set.

2. MAIN RESULTS

This section provides criteria for checking intra regularity of elements in $\overline{PT}(X, Y)$. By somewhat abusing the notation, we use $A\alpha$ to denote $(\text{dom } \alpha \cap A)\alpha$ for any $A \subseteq X$. Note for $\alpha, \beta \in \overline{PT}(X, Y)$, we have $\text{dom } (\alpha\beta) \subseteq \text{dom } \alpha, \text{im } (\alpha\beta) \subseteq \text{im } \beta$, and $|\text{im } \alpha| \leq |\text{dom } \alpha|$.

Lemma 1. *Let $\alpha, \beta \in \overline{PT}(X, Y)$. Then, $\alpha = \gamma\beta\mu$ for some $\gamma, \mu \in \overline{PT}(X, Y)$ if and only if $|\text{im } \alpha| \leq |\text{im } \beta|, |Y\alpha| \leq |Y\beta|$, and $|\text{im } \alpha \setminus Y| \leq |\text{im } \beta \setminus Y|$.*

Proof. Assume that $\alpha = \gamma\beta\mu$ for some $\gamma, \mu \in \overline{PT}(X, Y)$. Then $|\text{im } \alpha| = |(\text{dom } \alpha)\alpha| = |(\text{dom } (\gamma\beta\mu))\gamma\beta\mu| \leq |(\text{dom } (\gamma\beta))\gamma\beta\mu| = |(\text{im } (\gamma\beta))\mu| \leq |\text{im } (\gamma\beta)| \leq |\text{im } \beta|, |Y\alpha| = |(\text{dom } \alpha \cap Y)\alpha| = |(\text{dom } (\gamma\beta\mu) \cap Y)\gamma\beta\mu| \leq |(\text{dom } (\gamma\beta) \cap Y)\gamma\beta\mu| = |(Y\gamma\beta)\mu| \leq |Y\gamma\beta| \leq |Y\beta|$, and $|\text{im } \alpha \setminus Y| = |\text{im } (\gamma\beta\mu) \setminus Y| \leq |\text{im } (\beta\mu) \setminus Y| = |(\text{dom } (\beta\mu))\beta\mu \setminus Y| \leq |(\text{dom } (\beta\mu))\beta \setminus Y| \leq |(\text{dom } \beta)\beta \setminus Y| = |\text{im } \beta \setminus Y|$.

Conversely, assume that $|\text{im } \alpha| \leq |\text{im } \beta|, |Y\alpha| \leq |Y\beta|$, and $|\text{im } \alpha \setminus Y| \leq |\text{im } \beta \setminus Y|$. Write

$$\alpha = \begin{pmatrix} A_i & B_j & C_k \\ a_i & b_j & c_k \end{pmatrix},$$

where $A_i \cap Y \neq \emptyset; B_j, C_k \subseteq X \setminus Y; \{a_i\}, \{b_j\} \subseteq Y$; and $\{c_k\} \subseteq X \setminus Y$. By our assumptions, we can write

$$\beta = \begin{pmatrix} U_i & U_l & V_m & W_n & W_k \\ u_i & u_l & v_m & w_n & w_k \end{pmatrix},$$

where $U_i \cap Y \neq \emptyset \neq U_l \cap Y; V_m, W_n, W_k \subseteq X \setminus Y; \{u_i\}, \{u_l\}, \{v_m\} \subseteq Y; \{w_n\}, \{w_k\} \subseteq X \setminus Y$; and $|I| + |J| + |K| \leq |I| + |L| + |M| + |N| + |K|$.

Case 1. $|J| \leq |L| + |M| + |N|$. Let $L \cup M \cup N = P \dot{\cup} Q$, such that $|P| = |J|$. Then, we can express $\{U_l\} \cup \{V_m\} \cup \{W_n\} = \{R_p\} \cup \{S_q\}$ and rewrite β as

$$\beta = \begin{pmatrix} U_i & R_p & S_q & W_k \\ u_i & r_p & s_q & w_k \end{pmatrix}.$$

Since $|J| = |P|$, there exists a bijective function $\varphi : J \rightarrow P$. For each i, j , and k , choose $y_i \in U_i \cap Y$, $x_{j\varphi} \in R_{j\varphi}$, and $z_k \in W_k$, respectively. Now, define

$$\gamma = \begin{pmatrix} A_i & B_j & C_k \\ y_i & x_{j\varphi} & z_k \end{pmatrix} \text{ and } \mu = \begin{pmatrix} u_i & r_{j\varphi} & w_k \\ a_i & b_j & c_k \end{pmatrix}.$$

Hence, $\gamma, \mu \in \overline{PT}(X, Y)$ and $\alpha = \gamma\beta\mu$.

Case 2. $|J| > |L| + |M| + |N|$. Then, $\text{im } \beta$ is an infinite set. This implies $|J| \leq |I|$ or $|J| \leq |K|$ are infinite cardinals.

Subcase 2.1. $|J| \leq |I|$. Let $I = P \dot{\cup} Q$, such that $|P| = |I|$ and $|Q| = |J|$. Then, we can express $\{U_i\} = \{R_p\} \cup \{S_q\}$ in which $R_p \cap Y \neq \emptyset$ and rewrite β as

$$\beta = \begin{pmatrix} R_p & S_q & U_l & V_m & W_n & W_k \\ r_p & s_q & u_l & v_m & w_n & w_k \end{pmatrix}.$$

Since $|P| = |I|$ and $|Q| = |J|$, there exist bijective functions $\varphi : I \rightarrow P$ and $\psi : J \rightarrow Q$. For each i, j , and k , choose $y_{i\varphi} \in R_{i\varphi} \cap Y$, $x_{j\psi} \in S_{j\psi}$ and $z_k \in W_k$, respectively. Define

$$\gamma = \begin{pmatrix} A_i & B_j & C_k \\ y_{i\varphi} & x_{j\psi} & z_k \end{pmatrix} \text{ and } \mu = \begin{pmatrix} r_{i\varphi} & s_{j\psi} & w_k \\ a_i & b_j & c_k \end{pmatrix}.$$

Hence, $\gamma, \mu \in \overline{PT}(X, Y)$ and $\alpha = \gamma\beta\mu$.

Subcase 2.2. $|J| \leq |K|$. Let $K = G \dot{\cup} H$, such that $|G| = |J|$ and $|H| = |K|$. Then, we can express $\{W_k\} = \{D_g\} \cup \{E_h\}$ and rewrite β as

$$\beta = \begin{pmatrix} U_i & U_l & V_m & W_n & D_g & E_h \\ u_i & u_l & v_m & w_n & d_g & e_h \end{pmatrix}.$$

Since $|G| = |J|$ and $|H| = |K|$, there exist bijective functions $\sigma : J \rightarrow G$ and $\theta : K \rightarrow H$. For each i, j , and k , choose $y_i \in U_i \cap Y$, $x_{j\sigma} \in D_{g\sigma}$ and $z_{k\theta} \in E_{k\theta}$, respectively. Define

$$\gamma = \begin{pmatrix} A_i & B_j & C_k \\ y_i & x_{j\sigma} & z_{k\theta} \end{pmatrix} \text{ and } \mu = \begin{pmatrix} u_i & d_{j\sigma} & e_{k\theta} \\ a_i & b_j & c_k \end{pmatrix}.$$

Hence, $\gamma, \mu \in \overline{PT}(X, Y)$ and $\alpha = \gamma\beta\mu$. ■

Since $\text{im } \alpha^2 \subseteq \text{im } \alpha, Y\alpha^2 \subseteq Y\alpha$ and $\text{im } \alpha^2 \setminus Y \subseteq \text{im } \alpha \setminus Y$, we obtain the following criterion.

Theorem 2. *Let $\alpha \in \overline{PT}(X, Y)$. Then, α is intra-regular if and only if $|\text{im } \alpha| = |\text{im } \alpha^2|, |Y\alpha| = |Y\alpha^2|$ and $|\text{im } \alpha \setminus Y| = |\text{im } \alpha^2 \setminus Y|$.*

In order to re-write the above criterion in terms of α , where $\text{im } \alpha$ is finite, the following three lemmas are needed.

Lemma 3. *Let $\alpha \in \overline{PT}(X, Y)$ be such that $\alpha^2 \neq \emptyset$ and $\text{im } \alpha$ is finite. Then, $|\text{im } \alpha| = |\text{im } \alpha^2|$ if and only if $\text{im } \alpha \subseteq \text{dom } \alpha$ and $\alpha|_{\text{im } \alpha}$ is injective.*

Proof. Assume $|\text{im } \alpha| = |\text{im } \alpha^2|$. To show $\text{im } \alpha \subseteq \text{dom } \alpha$, we suppose, to the contrary, that there exists $x \in \text{im } \alpha \setminus \text{dom } \alpha$. Then, $|\text{im } \alpha^2| = |(\text{dom } \alpha^2)\alpha^2| \leq |(\text{dom } \alpha)\alpha^2| = |(\text{im } \alpha)\alpha| < |\text{im } \alpha|$ which is a contradiction. Thus, $\text{im } \alpha \subseteq \text{dom } \alpha$. To show $\alpha|_{\text{im } \alpha}$ is injective, assume the contrary that there exist distinct $x_1, x_2 \in \text{im } \alpha$, such that $x_1\alpha = x_2\alpha$. Since $x_1, x_2 \in \text{im } \alpha$, there exist $x'_1, x'_2 \in \text{dom } \alpha$ such that $x'_1\alpha = x_1$ and $x'_2\alpha = x_2$. Since α is a function and $x_1 \neq x_2$, we obtain $x'_1 \neq x'_2$. However, $x'_1\alpha^2 = x_1\alpha = x_2\alpha = x'_2\alpha^2$, which leads $|\text{im } \alpha^2| < |\text{im } \alpha|$, a contradiction.

Conversely, assume that $\text{im } \alpha \subseteq \text{dom } \alpha$ and $\alpha|_{\text{im } \alpha}$ is injective. Let $\text{im } \alpha = \{x_1, \dots, x_n\} \subseteq \text{dom } \alpha$. Since $\alpha|_{\text{im } \alpha}$ is injective,

$$|\text{im } \alpha| = |\{x_1, \dots, x_n\}| = |\{x_1\alpha, \dots, x_n\alpha\}| = |\text{im } \alpha^2|. \quad \blacksquare$$

Lemma 4. *Let $\alpha \in \overline{PT}(X, Y)$ be such that $\alpha^2 \neq \emptyset$ and $\text{im } \alpha$ is finite. Then, $|Y\alpha| = |Y\alpha^2|$ if and only if $Y\alpha \subseteq \text{dom } \alpha$ and $\alpha|_{Y\alpha}$ is injective.*

Proof. Assume that $|Y\alpha| = |Y\alpha^2|$. To show $Y\alpha \subseteq \text{dom } \alpha$, we suppose, to the contrary, that there exists $x \in Y\alpha \setminus \text{dom } \alpha$. Then, $|Y\alpha^2| = |(Y\alpha)\alpha| < |Y\alpha|$ which is a contradiction. To show $\alpha|_{Y\alpha}$ is injective, we assume the contrary that there exist distinct $y_1, y_2 \in Y\alpha$, such that $y_1\alpha = y_2\alpha$. Since $y_1, y_2 \in Y\alpha$, there exist $y'_1, y'_2 \in \text{dom } \alpha \cap Y$, such that $y'_1\alpha = y_1$ and $y'_2\alpha = y_2$. Since α is a function and $y_1 \neq y_2$, we obtain that $y'_1 \neq y'_2$. Thus, $y'_1\alpha^2 = y_1\alpha = y_2\alpha = y'_2\alpha^2$, which leads $|Y\alpha^2| < |Y\alpha|$, a contradiction.

Conversely, assume that $Y\alpha \subseteq \text{dom } \alpha$ and $\alpha|_{Y\alpha}$ is injective. Let $Y\alpha = \{y_1, \dots, y_n\} \subseteq \text{dom } \alpha$. Since $\alpha|_{Y\alpha}$ is injective,

$$|Y\alpha| = |\{y_1, \dots, y_n\}| = |\{y_1\alpha, \dots, y_n\alpha\}| = |Y\alpha^2|. \quad \blacksquare$$

Lemma 5. *Let $\alpha \in \overline{PT}(X, Y)$ be such that $\alpha^2 \neq \emptyset$ and $\text{im } \alpha$ is finite. Then, $|\text{im } \alpha \setminus Y| = |\text{im } \alpha^2 \setminus Y|$ if and only if $\text{im } \alpha \setminus Y \subseteq \text{dom } \alpha$, $\alpha|_{\text{im } \alpha \setminus Y}$ is injective, and $\text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \subseteq X \setminus Y$.*

Proof. Assume that $|\text{im } \alpha \setminus Y| = |\text{im } \alpha^2 \setminus Y|$. To show $\text{im } \alpha \setminus Y \subseteq \text{dom } \alpha$, we suppose, to the contrary, that there exists $x \in (\text{im } \alpha \setminus Y) \setminus \text{dom } \alpha$. Then, $|\text{im } \alpha^2 \setminus Y| \leq |(\text{im } \alpha \setminus Y)\alpha| < |\text{im } \alpha \setminus Y|$, which is a contradiction. To show $\alpha|_{\text{im } \alpha \setminus Y}$ is injective, assume the contrary that there exist distinct $x_1, x_2 \in \text{im } \alpha \setminus Y$ such that $x_1\alpha = x_2\alpha$. Since $x_1, x_2 \in \text{im } \alpha \setminus Y$, there exist $x'_1, x'_2 \in \text{dom } \alpha \setminus Y$ such that $x'_1\alpha = x_1$ and $x'_2\alpha = x_2$. Since α is a function and $x_1 \neq x_2$, then $x'_1 \neq x'_2$. Thus, $x'_1\alpha^2 = x_1\alpha = x_2\alpha = x'_2\alpha^2$, which leads $|\text{im } \alpha^2 \setminus Y| \leq |(\text{im } \alpha \setminus Y)\alpha| < |\text{im } \alpha \setminus Y|$, a contradiction. To show $\text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \subseteq X \setminus Y$, we letting $z \in \text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \cap Y$.

Hence, there exists $x \in (\text{im } \alpha \setminus Y) \cap \text{dom } \alpha$ such that $x\alpha = z$. This yields $|\text{im } \alpha^2 \setminus Y| < |(\text{im } \alpha \setminus Y)\alpha| \leq |\text{im } \alpha \setminus Y|$, which is a contradiction.

Conversely, assume that all three aforementioned conditions hold. Let $\text{im } \alpha \setminus Y = \{x_1, \dots, x_n\} \subseteq \text{dom } \alpha$. Since $\alpha|_{\text{im } \alpha \setminus Y}$ is injective and $\text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \subseteq X \setminus Y$, we obtain $|\text{im } \alpha \setminus Y| = |\{x_1, \dots, x_n\}| = |\{x_1\alpha, \dots, x_n\alpha\}| = |\text{im } \alpha^2 \setminus Y|$. ■

Combining Theorem 2 with Lemmas 3–5, we obtain the following theorem.

Theorem 6. *Let $\alpha \in \overline{PT}(X, Y)$ be such that $\text{im } \alpha$ is finite. Then, α is intra-regular if and only if all of the following statements hold:*

1. $\text{im } \alpha \subseteq \text{dom } \alpha$;
2. $\alpha|_{\text{im } \alpha}$ is injective;
3. $\text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \subseteq X \setminus Y$.

We now provide properties of intra-regular elements with respect to the finiteness of their images.

Lemma 7. *Let $\alpha \in \overline{PT}(X, Y)$ be such that $\text{im } \alpha$ is finite. If α is intra-regular, then the following statements hold:*

1. $(X \setminus Y)\alpha \subseteq (X \setminus Y) \cup Y\alpha$.
2. $|y\alpha^{-1} \cap Y\alpha| = 1$ for all $y \in Y\alpha$.
3. $|x\alpha^{-1} \cap (\text{im } \alpha \setminus Y\alpha)| = 1$ for all $x \in \text{im } \alpha \setminus Y\alpha$.

Proof. Assume that α is intra-regular. According to Theorem 6, we can express this α as

$$\alpha = \begin{pmatrix} A_1 & \cdots & A_r & B_1 & \cdots & B_s & C_1 & \cdots & C_t \\ a_1 & \cdots & a_r & b_1 & \cdots & b_s & c_1 & \cdots & c_t \end{pmatrix},$$

where $A_i \cap Y \neq \emptyset, B_j, C_k \subseteq X \setminus Y$ and $a_i, b_j \in Y, c_k \in X \setminus Y$ for all $i \in I = \{1, \dots, r\}, j \in J = \{1, \dots, s\}$ and $k \in K = \{1, \dots, t\}$. To show $J = \emptyset$, we suppose, to the contrary, that there exists $j_0 \in J$. Then $b_{j_0} \in \text{im } \alpha \subseteq \text{dom } \alpha$. Since $b_{j_0} \in Y$, we obtain $b_{j_0} \in A_{i_0}$ for some $i_0 \in I$. Hence, $B_{j_0}\alpha^2 = a_{i_0}$. Since $\text{im } (\alpha|_{\text{im } \alpha \setminus Y}) \subseteq X \setminus Y$, we obtain

$$|\text{im } \alpha^2| = |\text{im } \alpha^2 \cap Y| + |\text{im } \alpha^2 \setminus Y| \leq r + (s - 1) + t < r + s + t = |\text{im } \alpha|$$

which is a contradiction by Theorem 2. Consequently,

$$(*) \quad \alpha = \begin{pmatrix} A_1 & \cdots & A_r & C_1 & \cdots & C_t \\ a_1 & \cdots & a_r & c_1 & \cdots & c_t \end{pmatrix}.$$

1. According to (*), we have $(X \setminus Y)\alpha \subseteq \text{im } \alpha \subseteq Y\alpha \cup X \setminus Y$ which yields $(X \setminus Y)\alpha \subseteq (X \setminus Y) \cup Y\alpha$.

2. Since $\text{im } \alpha \subseteq \text{dom } \alpha$, we obtain a_1, \dots, a_r belong to A_i for some $i \in I$. Since $\alpha|_{\text{im } \alpha}$ is injective, we obtain $|A_i \cap \{a_1, \dots, a_r\}| = 1$ for all $i \in I$, that is, $|y\alpha^{-1} \cap Y\alpha| = 1$ for all $y \in Y\alpha$.

3. Since $\text{im } \alpha \subseteq \text{dom } \alpha$ and $\alpha|_{\text{im } \alpha}$ is injective, we obtain c_1, \dots, c_t belong to C_k for some $k \in K$ and $|C_k \cap \{c_1, \dots, c_t\}| = 1$ for all $k \in K$, that is, $|x\alpha^{-1} \cap (\text{im } \alpha \setminus Y\alpha)| = 1$ for all $x \in \text{im } \alpha \setminus Y\alpha$. ■

For the rest of this section, X is assumed to be a finite set with n elements and $\emptyset \neq Y \subseteq X$ has m elements. The following lemmas will serve as our starting point as we count the intra-regular elements in $\overline{PT}(X, Y)$.

Lemma 8. *Let $|X| = n, |Y| = m$ and $\alpha \in \overline{PT}(X, Y)$ be intra-regular. If $\text{im } \alpha \cap Y \neq \emptyset$, then $\alpha|_Y : (\text{dom } \alpha \cap Y) \rightarrow Y$ has $\sum_{r=1}^m \sum_{s=0}^{m-r} \binom{m}{r} \binom{m-r}{s} r!r^s$ different forms.*

Proof. Let $\emptyset \neq \text{im } \alpha \cap Y = Y' = \{a_1, \dots, a_r\}$, where $1 \leq r \leq m$. From (*), we obtain $Y\alpha = \text{im } \alpha \cap Y = Y'$. We can write

$$\alpha|_Y = \begin{pmatrix} A_1 & \cdots & A_r \\ a_1 & \cdots & a_r \end{pmatrix},$$

where $\bigcup_{i=1}^r A_i = \text{dom } \alpha \cap Y$. By Theorem 6 and Lemma 7, we have $\text{im } \alpha \subseteq \text{dom } \alpha$, $\alpha|_{\text{im } \alpha}$ is injective and $|A_i \cap \{a_1, \dots, a_r\}| = 1$ for all $i = 1, \dots, r$. Since the number of permutations on $\{a_1, \dots, a_r\}$ is $r!$ and the number of ways of choosing Y' with $|Y'| = r$ is equal to $\binom{m}{r}$, we obtain that $\alpha|_{Y'} : Y' \rightarrow Y'$ has $\binom{m}{r}r!$ different forms. Let $Y^* = (\text{dom } \alpha \cap Y) \setminus Y'$ be such that $|Y^*| = s$. Then, the number of ways of choosing Y^* is equal to $\binom{m-r}{s}$ and there are r options for where to place each elements of Y^* in A_1, \dots, A_r . Moreover, since $1 \leq r \leq m$, we conclude that $\alpha|_Y$ has $\sum_{r=1}^m \sum_{s=0}^{m-r} \binom{m}{r} \binom{m-r}{s} r!r^s$ different forms. ■

Lemma 9. *Let $|X| = n, |Y| = m$, $\alpha \in \overline{PT}(X, Y)$ be intra-regular and $X' = \text{im } \alpha \setminus Y\alpha$. Then $\alpha|_{X'} : X' \rightarrow X'$ has $\sum_{t=0}^{n-m} \binom{n-m}{t} t!$ different forms.*

Proof. Suppose that $|X'| = t$ where $0 \leq t \leq n - m$. By Lemma 7(3), $\alpha|_{X'}$ is a permutation on X' . Hence, $\alpha|_{X'}$ has $t!$ different forms. Since the number of ways of choosing X' is equal to $\binom{n-m}{t}$, we obtain that $\alpha|_{X'} : X' \rightarrow X'$ has $\sum_{t=0}^{n-m} \binom{n-m}{t} t!$ different forms. ■

Theorem 10. *The number of intra-regular elements in $\overline{PT}(X, Y)$ is*

$$\begin{aligned} & \sum_{r=1}^m \sum_{s=0}^{m-r} \sum_{t=0}^{n-m} \sum_{j=0}^{n-m-t} \binom{m}{r} \binom{m-r}{s} \binom{n-m}{t} \binom{n-m-t}{j} r!t!r^s(r+t)^j \\ & + \sum_{t=1}^{n-m} \sum_{j=0}^{n-m-t} \binom{n-m}{t} \binom{n-m-t}{j} t!t^j + 1, \end{aligned}$$

where $|X| = n$ and $|Y| = m$.

Proof. Let $\alpha \in \overline{PT}(X, Y)$ be an intra-regular element. Suppose that $Y' = Y\alpha = \{a_1, \dots, a_r\}$ and $X' = \text{im } \alpha \setminus Y\alpha = \{c_1, \dots, c_t\}$.

Case 1. $Y' \neq \emptyset$. Lemma 7 allow us to express this α as

$$\alpha = \begin{pmatrix} A_1 & \cdots & A_r & C_1 & \cdots & C_t \\ a_1 & \cdots & a_r & c_1 & \cdots & c_t \end{pmatrix},$$

where $A_i \cap Y \neq \emptyset, C_k \subseteq X \setminus Y$ and $|A_i \cap \{a_1, \dots, a_r\}| = 1, |C_k \cap \{c_1, \dots, c_t\}| = 1$ for all $i = 1, \dots, r$ and $k = 1, \dots, t$. Let $Y^* = (\text{dom } \alpha \cap Y) \setminus Y'$ be such that $|Y^*| = s$. By Lemmas 8 and 9, there are $\sum_{r=1}^m \sum_{s=0}^{m-r} \sum_{t=0}^{n-m} \binom{m}{r} \binom{m-r}{s} \binom{n-m}{t} r!t!r^s$ different forms of $\alpha|_{Y \cup X'}$. Let $X^* = \text{dom } \alpha \setminus (Y \cup X')$ be such that $|X^*| = j$. Then, there are $r + t$ different ways to arrange each element of X^* in $A_1, \dots, A_r, C_1, \dots, C_t$ and X^* has $\binom{n-m-t}{j}$ forms, where $0 \leq j \leq n - m - t$. Therefore, the number of intra-regular elements in this case is

$$\sum_{r=1}^m \sum_{s=0}^{m-r} \sum_{t=0}^{n-m} \sum_{j=0}^{n-m-t} \binom{m}{r} \binom{m-r}{s} \binom{n-m}{t} \binom{n-m-t}{j} r!t!r^s(r+t)^j.$$

Case 2. $Y' = \emptyset$. If $X' = \emptyset$, then $\alpha = \emptyset$, which is intra-regular. If $X' \neq \emptyset$, then $\alpha|_{X'} : X' \rightarrow X'$ has $\sum_{t=1}^{n-m} \binom{n-m}{t} t!$ different forms by Lemma 9. Let $X^* = \text{dom } \alpha \setminus X' \subseteq (X \setminus Y) \setminus X'$ be such that $|X^*| = j$. Then, there are t different ways to arrange each element of X^* in C_1, \dots, C_t and X^* has $\binom{n-m-t}{j}$ forms, where $0 \leq j \leq n - m - t$. Therefore, the number of intra-regular elements in this case is

$$\sum_{t=1}^{n-m} \sum_{j=0}^{n-m-t} \binom{n-m}{t} \binom{n-m-t}{j} t!t^j.$$

As a result, $\overline{PT}(X, Y)$ has the following number of intra-regular elements

$$\begin{aligned} & \sum_{r=1}^m \sum_{s=0}^{m-r} \sum_{t=0}^{n-m} \sum_{j=0}^{n-m-t} \binom{m}{r} \binom{m-r}{s} \binom{n-m}{t} \binom{n-m-t}{j} r!t!r^s(r+t)^j \\ & + \sum_{t=1}^{n-m} \sum_{j=0}^{n-m-t} \binom{n-m}{t} \binom{n-m-t}{j} t!t^j + 1. \end{aligned}$$

This completes the proof. ■

Acknowledgement

This work was supported by Thammasat University Research Unit in Algebra and Its Applications.

REFERENCES

- [1] W. Choomanee, P. Honyam and J. Sanwong, *Regularity in semigroups of transformations with invariant sets*, Int. J. Pure Appl. Math. **87(1)** (2013) 151–164.
<https://doi.org/10.12732/ijpam.v87i1.9>
- [2] A.H. Clifford and G.B. Preston, *The Algebraic Theory of Semigroups, Vol. I*, Mathematical Surveys, 7 (American Mathematical Society, Providence, R.I., 1961).
<https://doi.org/10.1090/surv/007.1>
- [3] A.H. Clifford and G.B. Preston, *The Algebraic Theory of Semigroups, Vol. II*, Mathematical Surveys, 7 (American Mathematical Society, Providence, R.I., 1967).
<https://doi.org/10.1090/surv/007.2>
- [4] C. Doss, *Certain Equivalence Relations in Transformation Semigroups*, Master's thesis, directed by D.D. Miller (University of Tennessee, 1955).
- [5] P. Honyam and J. Sanwong, *Semigroups of transformations with invariant set*, J. Korean Math. Soc. **48(2)** (2011) 289–300.
<https://doi.org/10.4134/JKMS.2011.48.2.289>
- [6] J.M. Howie, *Fundamentals of Semigroup Theory* (London Mathematical Society Monographs, New Series, 12, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995).
- [7] K.D. Jr. Magill, *Subsemigroups of $S(X)$* , Math. Japonica **11** (1966) 109–115.

Received 12 February 2023

Revised 15 May 2023

Accepted 16 May 2023