

4 **ON 1-ABSORBING PRIME AND WEAKLY 1-ABSORBING**  
5 **PRIME IDEALS OF SEMIRINGS**

6 SAMPAD DAS<sup>1</sup>

7 *Department of Mathematics*  
8 *Jadavpur University, Kolkata-700032, India*  
9 **e-mail:** jumathsampad@gmail.com

10 AND

11 MANASI MANDAL

12 *Department of Mathematics*  
13 *Jadavpur University, Kolkata-700032, India*  
14 **e-mail:** manasi\_ju@yahoo.in

15 **Abstract**

16 In this study, we introduce the concept of 1-absorbing prime ideals and  
17 weakly 1-absorbing prime ideals of commutative semirings with non-zero  
18 identity. A proper ideal  $I$  of a commutative semiring  $S$  is said to be a  
19 1-absorbing (resp. a weakly 1-absorbing) prime ideal if whenever  $abc \in I$   
20 (resp.  $0 \neq abc \in I$ ) for some non-units  $a, b, c \in S$ , then either  $ab \in I$   
21 or  $c \in I$ . The relationships among 1-absorbing prime ideals, prime ideals,  
22 2-prime ideals, and 2-absorbing ideals are investigated. 1-absorbing prime  
23 ideals of subtractive valuation domain are studied. Many properties, results,  
24 and characterizations of 1-absorbing prime (resp. weakly 1-absorbing prime)  
25 ideals are given.

26 **Keywords:** semiring, 1-absorbing prime ideals, weakly 1-absorbing prime  
27 ideals, 2-prime ideals.

28 **2020 Mathematics Subject Classification:** 16Y60.

---

<sup>1</sup>Corresponding author.

## 1. INTRODUCTION

The concept of semiring was introduced by H.S. Vandiver in 1935, as a generalization of both rings and distributive lattices. Several authors have applied this concept in various disciplines in many ways. Semirings have wide applications in different branches of mathematics and computer science such as Optimization Theory, Automata Theory, Cryptography, Graph Theory, Topology, etc. A non-empty set  $S$  together with two binary operations addition  $+$  and multiplication  $\cdot$  is called semiring if:

- (1)  $(S, +)$  is a commutative monoid with identity element 0.
- (2)  $(S, \cdot)$  is a monoid with identity element  $1 \neq 0$ .
- (3) The multiplication both from left and right is distributes over addition.
- (4)  $0 \cdot a = a \cdot 0 = 0$  for every  $a \in S$ .

A semiring  $S$  is commutative if  $ab = ba$  for all  $a, b \in S$ . In this paper, all semirings are assumed to be commutative and with non-zero identity.

A non-empty subset  $I$  of a semiring  $S$  is called an ideal of  $S$  if for  $a, b \in I$  and  $r \in S$ ,  $a + b \in I$  and  $ra \in I$ . An ideal  $I$  is proper if  $I \neq S$ . An ideal  $I$  of a semiring  $S$  is called subtractive ideal (or  $k$ -ideal) if  $a, a + b \in I$  and  $b \in S$ , implies  $b \in I$ . A semiring  $S$  is called subtractive if every ideal of  $S$  is a subtractive ideal. For an ideal  $I$  of  $S$  and  $a \in S$  define  $(I : a) = \{x \in S : ax \in I\}$ . It is easy to see that if  $I$  is a subtractive ideal of  $S$ , then  $(I : a)$  is a subtractive ideal of  $S$ . A semiring  $S$  is said to be a regular semiring if for every element  $r \in S$  there exists an element  $x \in S$  such that  $rxr = r$ . A proper ideal  $I$  of a semiring  $S$  is said to be a strong ideal if for each  $a \in I$  there exists  $b \in I$  such that  $a + b = 0$  [4]. Radical of an ideal  $I$  is defined as  $\sqrt{I} = \{a \in S : a^n \in I \text{ for some positive integer } n\}$ . An ideal  $I$  is irreducible, if and only if  $A \cap B = I$  for two ideals  $A, B$  implies  $A = I$  or  $B = I$ . An element  $u \in S \setminus \{0\}$  is called a unit if there exists an element  $u'$  such that  $uu' = 1$ . We denote the set of all units of  $S$  by  $U(S)$ . A non-zero element  $a \in S$  is said to be a semi-unit in  $S$  if there exists  $r, s \in S$  such that  $1 + ra = sa$ . Let  $(S_1, +, \cdot)$  and  $(S_2, +, \cdot)$  be two semirings with zero elements  $0_{S_1}, 0_{S_2}$  and identity elements  $1_{S_1}, 1_{S_2}$  respectively, a mapping  $f : S_1 \rightarrow S_2$  is said to be a semiring homomorphism if  $f(a + b) = f(a) + f(b)$ ,  $f(ab) = f(a)f(b)$ ,  $f(0_{S_1}) = 0_{S_2}$  and  $f(1_{S_1}) = 1_{S_2}$  for all  $a, b \in S_1$ . A one-one homomorphism is called monomorphism. Let  $U$  be a multiplicatively closed subset of a semiring  $S$ . The relation is defined on the set  $S \times U$  by  $(s, a) \sim (t, b) \iff xsb = xat$  for some  $x \in U$  is an equivalence relation and the equivalence class of  $(s, a) \in S \times U$  denoted by  $s/a$ . The set of all equivalence classes of  $S \times A$  under " $\sim$ " is denoted by  $S_U$ .  $S_U$  forms a semiring, where addition and multiplication are defined by  $s/a + t/b = (sb + ta)/ab$  and  $(s/a)(t/b) = st/ab$ . Suppose that  $S$  is a commutative semiring,  $U$  is a multiplicatively closed subset and  $I$  is an ideal of  $S$ , then the set

68  $I_U = \{a/b : a \in I, b \in U\}$  is an ideal of  $S_U$ . The set  $I_U$  is called the localization  
69 of the ideal  $I$  at  $U$  [15].

70 A semifield is a commutative semiring in which the non-zero elements form  
71 a group under multiplication. Semiring  $S$  is referred to as "reduced semiring"  
72 if  $S$  contains no non-zero nilpotent elements. A semiring  $S$  is a semidomain if  
73 it is multiplicatively cancellative, i.e., if  $ab = ac$  and  $a \neq 0$ , then  $b = c$  for any  
74  $a, b, c \in S$ . A semiring  $S$  is a valuation semiring if it is a semidomain and the  
75 set of ideals are totally ordered by inclusion [13]. A semiring  $S$  is said to be an  
76 entire semiring if  $ab = 0$  implies either  $a = 0$  or  $b = 0$  for any  $a, b \in S$  [13].

77 A proper ideal  $I$  of  $S$  is called a prime (resp., weakly prime) if for any  $a, b \in S$ ,  
78  $ab \in I$  (resp.,  $0 \neq ab \in I$ ) implies either  $a \in I$  or  $b \in I$  [8]. A proper ideal  $I$  of a  
79 semiring  $S$  is primary if for all  $x, y \in S$ , we have that  $xy \in I$  and  $x \notin I$  implies  
80  $y^n \in I$  for some positive integer  $n$ . If  $I$  is primary, then  $\sqrt{I} = P$  is a prime ideal  
81 of  $S$  [[2], Theorem 38]. In this case, we also say that  $I$  is a  $P$ -primary ideal of  $S$ .  
82 A prime ideal  $P$  of a semiring  $S$  is called to be a minimal prime ideal of  $S$ , if for  
83 every prime ideal  $I$ ,  $I \subseteq P$  implies either  $I = \langle 0 \rangle$  or  $I = P$ . Let  $P \supseteq I$  be ideals  
84 of a semiring  $S$ , where  $P$  is prime, then  $P$  is a minimal prime ideal of  $I$  if and  
85 only if for each  $x \in P$ , there is some  $y \in S \setminus P$  and a nonnegative integer  $i$  such  
86 that  $yx^i \in I$  [14].

87 A proper ideal  $M$  is maximal (resp.  $k$ -maximal) if  $M \subseteq M' \subseteq S$  implies  
88 either  $M = M'$  or  $M' = S$  for any ideal (resp.  $k$ -ideal)  $M'$  of  $S$ . A semiring  
89 is said to be a local semiring if it has only one maximal ideal. A semiring  $S$  is  
90 a local semiring if and only if  $S \setminus U(S)$  is an ideal of  $S$ . The Jacobson radical  
91  $Jac(S)$  of a semiring  $S$  is defined as the intersection of all maximal  $k$ -ideals of  $S$ .  
92 A prime ideal  $P$  of  $S$  is said to be a divided prime ideal of  $S$  if  $P \subseteq \langle x \rangle$  for every  
93  $x \in S \setminus P$ . If each prime ideal of  $S$  is divided, then  $S$  is called divided semiring  
94 [17]. For general references on semiring theory, one may refer to [9, 12] and [15].

95 A. Badawi introduced 2-absorbing (resp., weakly 2-absorbing) ideals in com-  
96 mutative rings as a generalization of prime ideals. The concept of 2-absorbing  
97 ideals in a commutative semiring was introduced by Darani [7]. A proper ideal  
98  $I$  of a semiring  $S$  is called a 2-absorbing ideal if for any  $a, b, c \in S$ ,  $abc \in I$   
99 implies  $ab \in I$  or  $bc \in I$  or  $ca \in I$ . The concept of 2-prime ideals of a ring  
100 was introduced by Beddani and Messirdi [5]. A proper ideal  $I$  of a semiring  $S$  is  
101 called 2-prime if for any  $a, b \in S$  such that  $ab \in I$ , either  $a^2 \in I$  or  $b^2 \in I$ , which  
102 was introduced by Khanra *et al.* [10]. The concept of 1-absorbing prime ideals  
103 which is another extension of prime ideals was introduced in [18]. The notions  
104 of 1-absorbing primary and weakly 1-absorbing primary ideals were investigated  
105 in [16]. A proper ideal  $I$  of a ring  $R$  is called a 1-absorbing prime (resp. weakly  
106 1-absorbing prime) ideal if for all non-unit elements  $a, b, c \in R$  such that  $abc \in I$   
107 (resp.  $0 \neq abc \in I$ ), then either  $ab \in I$  or  $c \in I$  [11, 18]. In this paper, we general-  
108 ize this notion of 1-absorbing prime ideals in commutative semiring with non-zero

identity. Throughout the paper, unless otherwise stated,  $S$  stands for commutative semiring with non-zero identity and  $\mathbb{Z}_0^+$  for the semiring of non-negative integers with usual binary addition and multiplication.

We shortly summarize the content of the paper. In the first section, we recall some essential preliminaries. In Section 2, we introduce 1-absorbing prime ideals as a generalization of prime ideals of a semiring. Some properties of 1-absorbing prime ideals are studied. We give a characterization of 1-absorbing prime ideals (cf. Theorem 7). We show (cf. Theorem 12) that in a subtractive valuation semiring  $S$ , an ideal  $I$  is a 1-absorbing prime ideal of  $S$  if and only if either  $I = P$  or  $I = P^2$ , where  $P = \sqrt{I}$  is a prime ideal of  $S$ . In Section 3, we define weakly 1-absorbing prime ideal, as a generalization of weakly prime ideals. Some properties of 1-absorbing prime ideals are studied. We characterize a strong local semiring whose all proper ideals are 1-absorbing prime (cf. Theorem 24). At the end, we characterize 1-absorbing prime ideals in a decomposable semiring (cf. Theorem 26).

## 2. 1-ABSORBING PRIME IDEALS

**Definition 1.** A proper ideal  $I$  of a commutative semiring  $S$  is said to be a 1-absorbing prime ideal if whenever  $abc \in I$  for some non-units  $a, b, c \in S$ , then either  $ab \in I$  or  $c \in I$ .

It is clear that every prime ideal is 1-absorbing prime and every 1-absorbing prime ideal is 2-absorbing ideal, but the converse may not be true in general which can be seen from the following examples:

**Example 1.** In semiring  $S = \mathbb{Z}_0^+$ , the ideal  $I = 3\mathbb{Z}_0^+ \setminus \{3\}$  is a 1-absorbing prime ideal which is not a prime ideal.

**Example 2.** Let  $S = \mathbb{Z}_0^+ \times \mathbb{Z}_0^+$  and  $I = 2\mathbb{Z}_0^+ \times 3\mathbb{Z}_0^+$ . Then  $I$  is a 2-absorbing ideal, however  $I$  is not a 1-absorbing prime ideal of  $S$ . Indeed,  $(2, 5)(4, 2)(7, 3) \in I$  but neither  $(2, 5)(4, 2) \in I$  nor  $(7, 3) \in I$ .

**Theorem 1.** Every 1-absorbing prime ideal of a semiring  $S$  is a 2-prime ideal of  $S$ .

**Proof.** Suppose that  $I$  is a 1-absorbing prime ideal of  $S$  and  $xy \in I$  for some elements  $x, y \in S$ . If either  $x$  or  $y$  is a unit, then  $I$  is a 2-prime ideal of  $S$ . So assume that  $x, y$  are non-unit elements of  $S$ . Then  $x^2y \in I$ , this implies either  $x^2 \in I$  or  $y \in I$ . Thus either  $x^2 \in I$  or  $y^2 \in I$  and hence  $I$  is a 2-prime ideal of  $S$ . ■

The converse of Theorem 1 may not be true, as shown in the following example.

**Example 3.** In semiring  $S = \mathbb{Z}_0^+$ , the ideal  $I = 4\mathbb{Z}_0^+$  is a 2-prime ideal but  $I$  is not a 1-absorbing prime ideal, since  $2 \cdot 3 \cdot 2 \in I$  but neither  $2 \cdot 3 \in I$  nor  $2 \in I$ .

So the class of all 1-absorbing prime ideals of a semiring  $S$  is an intermediate class between the class of all prime ideals and the classes of 2-prime ideals, 2-absorbing ideals of  $S$ .

**Theorem 2.** Let  $I$  be a 1-absorbing prime ideal of a semiring  $S$ . Then  $x^2 \in I$  for every  $x \in \sqrt{I}$  and  $\sqrt{I}$  is a prime ideal of  $S$ .

**Proof.** Suppose  $x \in \sqrt{I}$ . Then there exists a smallest positive integer  $n$  such that  $x^n \in I$ . For  $n = 1$  or  $n = 2$ , it is clear. For  $n \geq 3$ ,  $x^n = xxx^{n-2} \in I$ . That implies either  $x^2 \in I$  or  $x^{n-2} \in I$ . Continuing this process, it is easy to conclude that  $x^2 \in I$ . Now consider  $ab \in \sqrt{I}$  for some  $a, b \in S$ . If one of  $a$  or  $b$  is a unit, then there is nothing to prove. Assume that both  $a$  and  $b$  are non-units and  $ab \in \sqrt{I}$ . Then  $(ab)^2 = aab^2 \in I$ , which implies that either  $a^2 \in I$  or  $b^2 \in I$ . That is either  $a \in \sqrt{I}$  or  $b \in \sqrt{I}$ . Therefore,  $\sqrt{I}$  is a prime ideal of  $S$ . ■

**Theorem 3.** Let  $I$  be a 2-prime ideal of  $S$ . If  $(P^2 : x) \subseteq I$  for any  $x \in P \setminus I$ , where  $\sqrt{I} = P$ . Then  $I$  is a 1-absorbing prime ideal of  $S$ .

**Proof.** Suppose that  $xyz \in I$  and  $xy \notin I$  for some non-units  $x, y, z \in S$ . Since  $I$  is a 2-prime ideal and  $(xy)z \in I$ , we obtain either  $(xy)^2 \in I$  or  $z^2 \in I$ . If  $(xy)^2 \in I$ , then  $xy \in P \setminus I$ . But then  $xy \in (P^2 : xy) \subseteq I$ , which is a contradiction. So  $z^2 \in I$ , that is  $z \in P$ . If  $z \in I$ , then we are done. If  $z \in P \setminus I$ , then  $z \in (P^2 : z) \subseteq I$ , a contradiction. Therefore,  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 4.** Suppose that a semiring  $S$  has a 1-absorbing prime ideal which is not a prime ideal. Then,  $S$  is a local semiring.

**Proof.** Assume that  $I$  is a 1-absorbing prime ideal which is not a prime ideal of  $S$ . So there exist non-unit elements  $a, b \in S$  such that  $ab \in I$  but  $a \notin I$ ,  $b \notin I$ . Now consider the set of all non-units  $S \setminus U(S)$ . Let  $x, y \in S \setminus U(S)$ . Then  $xab \in I$  and  $yab \in I$ . Since  $I$  is a 1-absorbing prime ideal of  $S$  and  $b \notin I$ , we have  $xa \in I$  and  $ya \in I$ . Hence  $(x + y)a = xa + ya \in I$ . If  $(x + y)$  is a unit, then  $a \in I$ , which is a contradiction. Therefore,  $x + y \in S \setminus U(S)$ . Again for any  $s \in S$  and  $x \in S \setminus U(S)$ ,  $sxab \in I$ . Since  $I$  is a 1-absorbing prime ideal of  $S$  and  $a, b \notin I$ ,  $sx \in S \setminus U(S)$  and hence  $S \setminus U(S)$  is an ideal of  $S$ . Therefore,  $S$  is a local semiring. ■

**Corollary 1.** In a commutative non-local semiring, a proper ideal is a 1-absorbing prime if and only if it is prime.

**Theorem 5.** Let  $S$  be a local semiring with maximal ideal  $M$ . A proper ideal  $I$  of  $S$  is 1-absorbing prime ideal if and only if either  $I$  is a prime ideal or  $M^2 \subseteq I$ .

**Proof.** Suppose that  $I$  is a 1-absorbing prime ideal of  $S$ . If  $I$  is not a prime ideal, then there exist non-units  $a, b \in S$  such that  $ab \in I$  but  $a \notin I$  and  $b \notin I$ . Let  $x, y \in M$ . Here  $xyab \in I$ . Since  $I$  is a 1-absorbing prime ideal and  $b \notin I$ , we have  $xya \in I$ . Again  $xy \in I$ , since  $a \notin I$ . Therefore  $xy \in I$  for any  $x, y \in M$ . Hence  $M^2 \subseteq I$ .

Conversely, if  $I$  is a prime ideal of  $S$ , then  $I$  is a 1-absorbing prime ideal of  $S$ . Consider  $M^2 \subseteq I$  and  $abc \in I$  for some non-units  $a, b, c \in S$ . Since  $a, b, c$  are non-units, so  $a, b, c \in M$ . Thus  $ab \in M^2 \subseteq I$  and hence  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Corollary 2.** *In a local semiring  $S$  with unique maximal ideal  $M$ , if  $M^2 = \langle 0 \rangle$ , then every proper ideal of  $S$  is a 1-absorbing prime ideal.*

**Theorem 6.** *Let  $S$  be a semiring. Then,  $\langle 0 \rangle$  is a 1-absorbing ideal of  $S$  if and only if  $S$  is an entire semiring or  $(S, M)$  is a local semiring such that  $M^2 = \langle 0 \rangle$ .*

**Proof.** Suppose that  $\langle 0 \rangle$  is a 1-absorbing ideal of  $S$  and  $S$  is not an entire semiring. So  $\langle 0 \rangle$  is a 1-absorbing prime ideal of  $S$  which is not a prime ideal. Thus, by Theorem 4 and Theorem 5,  $S$  is a local semiring with maximal ideal  $M$  such that  $M^2 = 0$ .

Conversely, if  $S$  is an entire semiring, then  $\langle 0 \rangle$  is a prime ideal of  $S$  and so  $\langle 0 \rangle$  is 1-absorbing prime ideal of  $S$ . If  $S$  is a local semiring with maximal ideal  $M$  such that  $M^2 = 0$ , then by Corollary 2,  $\langle 0 \rangle$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 7.** *Let  $I$  be a proper ideal of a semiring  $S$ , then the following are equivalent:*

- (1)  $I$  is a 1-absorbing prime ideal of  $S$ .
- (2) If  $abJ \subseteq I$  for non-units  $a, b \in S$ , then either  $ab \in I$  or  $J \subseteq I$ .
- (3) For any proper ideals  $K, L$  and non-unit element  $a$  of  $S$ , if  $aKL \subseteq I$ , then either  $aK \subseteq I$  or  $L \subseteq I$ .
- (4) If for any ideals  $J, K, L$  of  $S$ ,  $JKL \subseteq I$ , then either  $JK \subseteq I$  or  $L \subseteq I$ .

**Proof.** (1)  $\implies$  (2) Suppose that  $abJ \subseteq I$  and  $ab \notin I$ . Consider  $j \in J$ . Then  $abj \in abJ \subseteq I$ . Since  $I$  is a 1-absorbing prime ideal of  $S$ , we have  $j \in I$ . This implies that  $J \subseteq I$ .

(2)  $\implies$  (3) Assume that  $aKL \subseteq I$  and  $aK \not\subseteq I$ . Then there exists  $k \in K$  such that  $ak \notin I$ . As  $akL \subseteq I$ , by (2),  $L \subseteq I$ . Therefore either  $aK \subseteq I$  or  $L \subseteq I$ .

(3)  $\implies$  (4) Suppose  $JKL \subseteq I$  and  $JK \not\subseteq I$  for some ideals  $J, K, L$  of  $S$ . Then there exists an element  $j \in J$  such that  $jK \not\subseteq I$ . By (3), we have  $L \subseteq I$ . Therefore either  $JK \subseteq I$  or  $L \subseteq I$ .

(4)  $\implies$  (1) Assume that  $abc \in I$  for some non-units  $a, b, c \in S$ . Consider  $J = \langle a \rangle$ ,  $K = \langle b \rangle$ ,  $L = \langle c \rangle$ . Then  $JKL = abcS \subseteq I$ . By (4), either  $JK \subseteq I$  or  $L \subseteq I$ . So either  $ab \in I$  or  $c \in I$ . Hence  $I$  is a 1-absorbing prime ideal of  $S$ . ■

219 **Proposition 1.** *Let  $I$  be a 1-absorbing prime ideal of a semiring  $S$ . Then for*  
 220 *every non-unit  $c \in S \setminus I$ ,  $(I : c)$  is a prime ideal of  $S$ .*

221 **Proof.** Suppose  $ab \in (I : c)$  for some  $a, b \in S$  and non-unit  $c \in S \setminus I$ . Without  
 222 loss of generality, we may assume that  $a$  and  $b$  are non-units. Since  $I$  is a 1-  
 223 absorbing prime ideal of  $S$  and  $acb \in I$ , we have either  $ac \in I$  or  $b \in I$ . This  
 224 implies either  $a \in (I : c)$  or  $b \in I \subseteq (I : c)$ . Therefore,  $(I : c)$  is a prime ideal of  
 225  $S$  for every non-unit  $c \in S \setminus I$ . ■

226 **Theorem 8.** *If  $I$  is a  $P$ -primary ideal of  $S$  such that  $(P^2 : x) \subseteq I$  for all*  
 227  *$x \in P \setminus I$ , then  $I$  is a 1-absorbing prime ideal of  $S$ .*

228 **Proof.** Let  $abc \in I$  for some non-units  $a, b, c \in S$  and  $ab \notin I$ . If possible, let  
 229  $c \notin I$ . Since  $I$  is a  $P$ -primary ideal of  $S$  and  $ab \notin I$ , we have  $c \in P$ . Also  
 230  $c \notin I$  implies  $ab \in P$ . So  $abc \in P^2$ . By the given hypothesis,  $(P^2 : c) \subseteq I$ .  
 231 Thus  $ab \in (P^2 : c) \subseteq I$ , which is a contradiction. Hence  $c \in I$  and thus  $I$  is a  
 232 1-absorbing prime ideal of  $S$ . ■

233 **Theorem 9.** *Let  $P$  be a nonzero divided prime ideal of a semidomain  $S$ . Then,*  
 234  *$P^2$  is a 1-absorbing prime ideal of  $S$ .*

235 **Proof.** First, we show that  $P^2$  is a  $P$ -primary ideal of  $S$ . Let  $xy \in P^2$  and  $y \notin P$ .  
 236 Suppose  $xy = \sum_{i=1}^n x_i y_i$  for some  $x_i, y_i \in P$ ,  $i = 1, 2, \dots, n$ . Since  $P$  is a divided  
 237 prime ideal and  $y \notin P$ , we have  $P \subseteq Sy$ . Thus  $x_i = s_i y$ , where  $s_i \in S$ , for  
 238  $i = 1, 2, \dots, n$ . Since  $P$  is a prime ideal and  $y \notin P$ , we have  $s_i \in P$ . Also since  
 239  $S$  is a semidomain,  $xy = \sum_{i=1}^n s_i y y_i = (\sum_{i=1}^n s_i y_i) y$  implies  $x = \sum_{i=1}^n s_i y_i$ , that  
 240 is  $x \in P^2$ . Thus  $P^2$  is a  $P$ -primary ideal of  $S$ . Also,  $(P^2 : x) \subseteq P^2$  for every  
 241  $x \in P \setminus P^2$ . So by Theorem 8,  $P^2$  is a 1-absorbing prime ideal of  $S$ . ■

242 **Theorem 10.** *In a semiring  $S$ , for any 1-absorbing prime ideal  $I$  of  $S$ , there*  
 243 *exists exactly one minimal prime ideal containing  $I$ .*

244 **Proof.** If possible, let  $P_1$  and  $P_2$  be two minimal prime ideals containing  $I$ . Then  
 245 there exist elements  $a, b \in S$  such that  $a \in P_1 \setminus P_2$  and  $b \in P_2 \setminus P_1$ . So there  
 246 exist  $p_2 \notin P_2$  and  $p_1 \notin P_1$  such that  $ap_2^n \in I$ ,  $bp_1^n \in I$ . Since  $p_1, p_2 \notin I$  and  $I$  is  
 247 a 1-absorbing prime ideal,  $ap_2^n \in I$  implies  $ap_2 \in I$  and  $bp_1^n \in I$  implies  $bp_1 \in I$ .  
 248 Thus  $a^2 \in I$  and  $b^2 \in I$ , whence  $a^2 \in I \subseteq P_2$  implies  $a \in P_2$  and  $b^2 \in I \subseteq P_1$   
 249 implies  $b \in P_1$ , which is a contradiction. Therefore there exists one minimal prime  
 250 ideal of  $S$  containing  $I$ . ■

251 **Theorem 11.** *Let  $P$  be a non-zero divided prime ideal of a subtractive semiring*  
 252  *$S$  and  $I$  be an ideal of  $S$  such that  $\sqrt{I} = P$ . If  $I$  is a 1-absorbing prime ideal of*  
 253  *$S$ , then  $I$  is a  $P$ -primary ideal of  $S$  such that  $P^2 \subseteq I$ .*

**Proof.** Let  $ab \in I$  for some  $a, b \in S$  and  $b \notin P$ . Since  $P$  is a divided prime ideal of  $S$ , we have  $P \subseteq Sb$ . Thus  $a = sb$  for some  $s \in S$ . So, we get  $ab = b^2s \in I$ . Since  $b^2 \notin I$  and  $I$  is a 1-absorbing prime ideal of  $S$ , we conclude that  $s \in I$ . Therefore  $a = sb \in I$  and so  $I$  is a  $P$ -primary ideal of  $S$ . Now, let  $x, y \in P$ . Then by Theorem 2,  $x^2, y^2 \in I$ . Thus  $x(x+y)y = x^2y + xy^2 \in I$ . Since  $I$  is a 1-absorbing prime ideal of  $S$ , we have either  $x(x+y) = x^2 + xy \in I$  or  $y \in I$ . Since  $S$  is a subtractive semiring,  $xy \in I$  and hence  $P^2 \subseteq I$ . ■

**Theorem 12.** Suppose that  $S$  is a subtractive valuation semiring and  $I$  is an ideal of  $S$ . Then  $I$  is a 1-absorbing prime ideal of  $S$  if and only if either  $I = P$  or  $I = P^2$ , where  $P = \sqrt{I}$  is a prime ideal of  $S$ .

**Proof.** Suppose  $I$  is a 1-absorbing prime ideal of  $S$ . Since every subtractive valuation semiring is a divided semidomain [6], Proposition 1.4], so  $I$  is a  $P$ -primary ideal of  $S$  such that  $P^2 \subseteq I$ , by Theorem 11. Therefore, by [[6], Theorem 1.2], either  $I = P$  or  $I = P^2$ . Conversely, suppose that either  $I = P$  or  $I = P^2$ , where  $P = \sqrt{I}$  is a prime ideal of  $S$ . If  $I = P$ , then  $I$  is a 1-absorbing prime ideal of  $S$ . If  $I = P^2$ , then by Theorem 9,  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 13.** Let  $I$  be an irreducible subtractive ideal of  $S$  and  $(I : x) = (I : x^2)$  for every  $x \in S \setminus I$ , then  $I$  is a 1-absorbing prime ideal of  $S$ .

**Proof.** Let  $abc \in I$  and  $c \notin I$  for some non-units  $a, b, c \in S$ . By the assumption,  $(I : c) = (I : c^2)$ . If possible, let  $ab \notin I$ . Consider  $r \in (I + (ab)) \cap (I + (c))$ . Then  $r = i_1 + abs_1 = i_2 + cs_2$  for some  $i_1, i_2 \in I$  and  $s_1, s_2 \in S$ . Thus  $rc = i_1c + abcs_1 = i_2c + c^2s_2 \in I$ . Since  $I$  is a subtractive ideal of  $S$ , we get  $c^2s_2 \in I$ . So  $cs_2 \in I$ . Therefore,  $r = i_2 + cs_2 \in I$ . This shows that  $(I + (ab)) \cap (I + (c)) \subseteq I$  and hence  $(I + (ab)) \cap (I + (c)) = I$ , a contradiction. Thus  $ab \in I$  and so  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 14.** Let  $I$  be a 1-absorbing prime subtractive ideal of  $S$  with  $\sqrt{I} = P$ . If  $I \neq P$ , then  $\Omega = \{(I : x) : x \in P \setminus I\}$  under set inclusion is a totally ordered set.

**Proof.** If possible, let there exist  $a, b \in P \setminus I$  such that neither  $(I : a) \subseteq (I : b)$  nor  $(I : b) \subseteq (I : a)$ . Then there exists  $x, y \in S \setminus I$  so that  $x \in (I : a) \setminus (I : b)$  and  $y \in (I : b) \setminus (I : a)$ . Since  $P \subseteq (I : c)$  for any  $c \in S \setminus I$ ,  $x \in (I : a) \setminus P$  and  $y \in (I : b) \setminus P$ . Thus  $xy \notin P$ , since  $P$  is a prime ideal of  $S$ . Consider  $x(a+b)y = xay + xby \in I$ . This implies  $x(a+b) = xa + xb \in I$ . Since  $I$  is a subtractive ideal and  $xa \in I$ , we get  $xb \in I$ , that is  $x \in (I : b)$ , which is a contradiction. Hence  $\Omega = \{(I : x) : x \in P \setminus I\}$  under set inclusion is a totally ordered set. ■



**Theorem 15.** *In a regular semiring  $S$ , every irreducible ideal of  $S$  is a 1-absorbing prime ideal of  $S$ .*

**Proof.** Let  $S$  be a regular semiring and  $I$  be an irreducible ideal of  $S$ . Let  $abc \in I$  and  $ab \notin I$  for some non-units  $a, b, c \in S$ . If possible, let  $c \notin I$ . Now consider the ideals  $J = (I + (ab))$  and  $K = (I + (c))$ , properly containing  $I$ . Since  $I$  is an irreducible ideal of  $S$ ,  $J \cap K \not\subseteq I$ . Thus, there exists  $p \in S$  such that  $p \in (I + (ab)) \cap (I + (c)) \setminus I$ . So  $p \in (I + (ab)(I + (c))) \setminus I$ , by regularity of  $S$  [[9], Proposition 6.35]. Then, there are  $p_1, p_2 \in I$  and  $s_1, s_2 \in S$  such that  $p = (p_1 + s_1ab)(p_2 + s_2c) = p_1p_2 + p_1s_2c + s_1abp_2 + s_1s_2abc$ . This implies that  $p \in I$ , which is a contradiction. Hence  $c \in I$ , and so  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 16.** *Let  $S$  be a commutative semiring and  $U$  be a multiplicative closed subset of  $S$ . If  $I$  is a 1-absorbing prime ideal of  $S$  with  $U \cap I = \emptyset$ , then  $I_U$  is a 1-absorbing prime ideal of  $S_U$ .*

**Proof.** Let  $(a/s)(b/t)(c/r) \in I_U$  for some non-units  $(a/s), (b/t), (c/r) \in S_U$ , where  $a, b, c \in S$  and  $s, t, r \in U$ . Then  $(ua)bc \in I$  for some  $u \in U$ . So either  $uab \in I$  or  $c \in I$ , as  $I$  is a 1-absorbing prime ideal of  $S$ . Thus either  $(a/s)(b/t) \in I_U$  or  $(c/r) \in I_U$ . Therefore,  $I_U$  is a 1-absorbing prime ideal of  $S_U$ . ■

### 3. WEAKLY 1-ABSORBING PRIME IDEALS

**Definition 2.** A proper ideal  $I$  of a commutative semiring  $S$  is said to be a weakly 1-absorbing prime ideal if whenever  $0 \neq abc \in I$  for some non-units  $a, b, c \in S$ , either  $ab \in I$  or  $c \in I$ .

Clearly, every 1-absorbing prime ideal is also a weakly 1-absorbing prime ideal but the converse may be true in general which can be seen from the following examples.

**Example 4.** In the semiring  $\mathbb{Z}_{15}$ , the ideal  $I = \{0\}$  is clearly a weakly 1-absorbing prime ideal of  $S$  but not 1-absorbing prime, as  $0 = \bar{3}\bar{3}\bar{5} \in I$  but neither  $\bar{3}\bar{3} \in I$  nor  $\bar{5} \in I$ .

**Theorem 17.** *If  $I$  is a weakly 1-absorbing prime of a reduced semiring  $S$ , then  $\sqrt{I}$  is a weakly prime ideal of  $S$ .*

**Proof.** Suppose that  $0 \neq ab \in \sqrt{I}$  for some non-units  $a, b \in S$ . Then  $(ab)^n \in I$  for some positive integer  $n$ . Since  $S$  is a reduced semiring, so  $0 \neq (ab)^n = aa^{n-1}b^n \in I$ . Thus either  $aa^{n-1} = a^n \in I$  or  $b^n \in I$  which implies either  $a \in \sqrt{I}$  or  $b \in \sqrt{I}$ . So  $\sqrt{I}$  is a weakly prime ideal of  $S$ . ■

**Theorem 18.** *If the zero ideal of a semiring  $S$  is a 1-absorbing prime, then every weakly 1-absorbing prime ideal is a 1-absorbing prime of  $S$ .*

**Proof.** Suppose that  $I$  is a weakly 1-absorbing prime ideal of  $S$  and  $abc \in I$  for some non-units  $a, b, c \in S$ . If  $0 \neq abc$ , then either  $ab \in I$  or  $c \in I$ , that is,  $I$  is a 1-absorbing prime ideal. If  $abc = 0$  and since zero ideal is 1-absorbing prime, either  $ab = 0 \in I$  or  $c = 0 \in I$ . Therefore,  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 19.** *Let  $S_1$  and  $S_2$  be two semirings and  $f : S_1 \longrightarrow S_2$  be a monomorphism such that  $f(a)$  is a non-unit in  $S_2$  for every non-unit element  $a$  in  $S_1$ . Then the following statements hold:*

- (1) *If  $I$  is a weakly 1-absorbing prime ideal of  $S_2$ , then  $f^{-1}(I)$  is a weakly 1-absorbing prime ideal of  $S_1$ .*
- (2) *If  $J$  is a weakly 1-absorbing prime  $k$ -ideal of  $S_1$  with  $\text{Ker}(f) \subseteq J$  and  $f$  is onto steady homomorphism, then  $f(J)$  is a weakly 1-absorbing prime  $k$ -ideal of  $S_2$ .*

**Proof.** (1) Assume that  $0 \neq abc \in f^{-1}(I)$  for some non-units  $a, b, c \in S_1$ . Since  $f$  is monomorphism, we have  $0 \neq f(abc) = f(a)f(b)f(c) \in I$  for non-units  $f(a), f(b), f(c) \in S_2$ . As  $I$  is a weakly 1-absorbing prime ideal of  $S_2$ , so either  $f(a)f(b) = f(ab) \in I$  or  $f(c) \in I$ . That is either  $ab \in f^{-1}(I)$  or  $c \in f^{-1}(I)$ . Therefore,  $f^{-1}(I)$  is a weakly 1-absorbing prime ideal of  $S_1$ .

(2) Clearly,  $f(J)$  is a  $k$ -ideal of  $S_2$  as  $J$  is a  $k$ -ideal of  $S_1$ . Now consider,  $0 \neq abc \in f(J)$  for some non-units  $a, b, c \in S_2$ . There exist non-units  $x, y, z \in S_1$  such that  $f(x) = a, f(y) = b, f(z) = c$ . So  $0 \neq abc = f(x)f(y)f(z) = f(xyz) \in f(J)$ . Thus  $f(xyz) = f(j)$  for some  $j \in J$ . Since  $f$  is steady, we have  $xyz + k_1 = j + k_2$  for some  $k_1, k_2 \in \text{Ker}(f)$ . As  $J$  is a  $k$ -ideal of  $S_1$  and  $\text{Ker}(f) \subseteq J$ , we obtain  $0 \neq xyz \in J$ . Since  $J$  is a weakly 1-absorbing prime ideal of  $S_1$ , either  $xy \in J$  or  $z \in J$ . This implies either  $ab = f(xy) \in f(J)$  or  $c = f(z) \in f(J)$ . Hence,  $f(J)$  is a weakly 1-absorbing prime  $k$ -ideal of  $S_2$ . ■

**Definition 3.** Let  $I$  be a weakly 1-absorbing prime ideal of  $S$  and  $a, b, c \in S$  are non-units of  $S$ . We call  $(a, b, c)$  is a 1-triple zero of  $I$  if  $abc = 0, ab \notin I$  and  $c \notin I$ .

**Theorem 20.** *Let  $I$  be a subtractive weakly 1-absorbing prime ideal of  $S$  and  $(a, b, c)$  be a 1-triple zero of  $I$ . Then,  $abI = 0$  and if  $a, b \notin (I : c)$ , then  $I^3 = 0$ .*

**Proof.** Suppose that  $I$  is a subtractive weakly prime ideal of  $S$  and  $(a, b, c)$  is a 1-triple zero of  $I$ . So  $abc = 0$  and  $ab \notin I, c \notin I$ . If possible, let  $abI \neq 0$ . Then there exists  $i \in I$  such that  $0 \neq abi$ . This implies that  $0 \neq ab(c + i) = abc + abi \in I$ . Also  $c + i$  is a non-unit, because  $(c + i)$  is a unit implies  $ab \in I$ . Since  $I$  is weakly 1-absorbing prime and  $ab \notin I$ , so  $c + i \in I$ . As  $I$  is a subtractive ideal of  $S$ , we get  $c \in I$ , which is a contradiction. Therefore,  $abI = 0$ .

361 We first show that if  $a, b \notin (I : c)$ , then  $bcI = caI = aI^2 = bI^2 = cI^2 = 0$ .  
 362 Since  $a, b \notin (I : c)$ ,  $ac \notin I$  and  $bc \notin I$ . Suppose  $bcI \neq 0$ . Then there exists  
 363  $i \in I$  such that  $bci \neq 0$ . Since  $abc = 0$ , we have  $0 \neq bci = (a + i)bc \in I$ .  
 364 Here  $(a + i)$  is a non-unit, because if  $a + i$  is a unit, then  $bc \in I$ , which is a  
 365 contradiction. Since  $I$  is weakly 1-absorbing prime, either  $(a + i)b \in I$  or  $c \in I$ ,  
 366 that is either  $ab \in I$  or  $c \in I$ , a contradiction, as  $I$  is a subtractive ideal of  
 367  $S$ . Therefore  $bcI = 0$ . Similarly,  $caI = 0$ . Now, we will show that  $aI^2 = 0$ .  
 368 Suppose that  $ai_1i_2 \neq 0$  for some  $i_1, i_2 \in I$ . Since  $abc = 0$  and  $abI = acI = 0$ ,  
 369 we have  $a(b + i_1)(c + i_2) = abc + abi_2 + aci_1 + ai_1i_2 = ai_1i_2 \neq 0$ . If  $c + i_2$  is  
 370 a unit, then  $a(b + i_1) \in I$ , and so  $ab \in I$ . Similarly, if  $b + i_1$  is a unit, then  
 371  $ac \in I$ , both contradict the hypothesis. Since  $I$  is a weakly 1-absorbing prime  
 372 ideal, either  $a(b + i_1) \in I$  or  $c + i_2 \in I$ . Thus  $ab \in I$  or  $c \in I$  as  $I$  is a subtractive  
 373 ideal of  $S$ , which is a contradiction. So  $aI^2 = 0$ . Similarly, one can show that  
 374  $bI^2 = 0$  and  $cI^2 = 0$ . Now, assume that  $I^3 \neq 0$ . So there exist  $i_1, i_2, i_3 \in I$  such  
 375 that  $0 \neq i_1i_2i_3 \in I$ . Since  $abI = bcI = caI = aI^2 = bI^2 = cI^2 = 0$ , we have  
 376  $0 \neq i_1i_2i_3 = (a + i_1)(b + i_2)(c + i_3) \in I$  and  $(a + i_1), (b + i_2), (c + i_3)$  are non-units.  
 377 So either  $(a + i_1)(b + i_2) \in I$  or  $c + i_3 \in I$ . Since  $I$  is a subtractive ideal, we have  
 378 either  $ab \in I$  or  $c \in I$ , which is a contradiction. Therefore,  $I^3 = 0$ . ■

379 **Definition 4** ([1], Proposition 8.21). An ideal  $I$  of a semiring  $S$  is called a  $Q$ -  
 380 ideal if there exists a subset  $Q$  of  $S$  such that: 1)  $S = \cup\{q + I : q \in Q\}$ , 2) If  
 381  $q_1, q_2 \in Q$ , then  $(q_1 + I) \cap (q_2 + I) \neq \emptyset \iff q_1 = q_2$ . Let  $I$  be a  $Q$ -ideal of a  
 382 semiring  $S$ . Then  $S/I_Q = \{q + I : q \in Q\}$  forms a semiring under the following  
 383 addition “ $\oplus$ ” and multiplication “ $\odot$ ”,  $(q_1 + I) \odot (q_2 + I) = q_3 + I$  where  $q_3 \in Q$   
 384 is unique such that  $q_1 + q_2 + I \subseteq q_3 + I$ , and  $(q_1 + I) \odot (q_2 + I) = q_4 + I$  where  
 385  $q_4 \in Q$  is unique such that  $q_1q_2 + I \subseteq q_4 + I$ . This semiring  $S/I_Q$  is called the  
 386 quotient semiring of  $S$  by  $I$  and denoted by  $(S/I_Q, \oplus, \odot)$  or just  $S/I_Q$ .

387 **Theorem 21.** Let  $I$  be a  $Q$ -ideal and  $J$  be a  $k$ -ideal of a semiring  $S$ , so that  
 388  $I \subseteq J$ . If  $J$  is a weakly 1-absorbing prime ideal of  $S$ , then  $J/I_{(Q \cap J)}$  is a weakly  
 389 1-absorbing prime ideal of  $S/I_Q$ .

390 Let  $I$  be a weakly 1-absorbing prime ideal of  $S$  such that  $u(S/I) = \{x + I : x \in u(S) \cap Q\}$ . If  $J/I_{(Q \cap J)}$  is a weakly 1-absorbing prime ideal of  $S/I_Q$ , then  $J$  is a weakly 1-absorbing prime ideal of  $S$ .

393 **Proof.** Suppose that  $J$  is a weakly 1-absorbing prime ideal of  $S$  and  $0 \neq (q_1 + I) \odot (q_2 + I) \odot (q_3 + I) \in J/I_{(Q \cap J)}$  for some non-units  $q_1 + I, q_2 + I, q_3 + I \in S/I_Q$   
 394 where  $q_1, q_2, q_3 \in Q$ . So there exists a unique element  $q_4 \in J \cap Q$  such that  
 395  $q_1q_2q_3 + I \subseteq q_4 + I \in J/I_{(Q \cap J)}$ . It follows that  $q_1, q_2, q_3$  are non-units in  $S$  and  
 396  $0 \neq q_1q_2q_3 \in J$ . Since  $J$  is a weakly 1-absorbing prime ideal of  $S$ , we get  $q_1q_2 \in J$   
 397 or  $q_3 \in J$ . Therefore if  $q_3 \in J$ , then  $q_3 + I \in J/I_{(Q \cap J)}$ . Otherwise if  $q_1q_2 \in J$ , then  
 398  $(q_1 + I) \odot (q_2 + I) = q_5 + I$  where  $q_5$  is the unique element in  $Q$  and  $q_1q_2 + r = q_5 + s$

for some  $r, s \in I$ . Since  $J$  is a  $k$ -ideal,  $q_5 \in J$ . So  $(q_1 + I) \odot (q_2 + I) \in J/I_{(Q \cap J)}$ . Hence  $J/I_{(Q \cap J)}$  is a weakly 1-absorbing prime ideal of  $S/I_Q$ .

Let  $0 \neq abc \in J$  for some non-units  $a, b, c \in S$ . If  $abc \in I$ , then either  $ab \in I \subseteq J$  or  $c \in I \subseteq J$ . Assume that  $abc \notin I$ . Since  $I$  is a  $Q$ -ideal, there exist  $q_1, q_2, q_3 \in Q$  such that  $a \in q_1 + I$ ,  $b \in q_2 + I$  and  $c \in q_3 + I$ . Thus  $a = q_1 + i_1$ ,  $b = q_2 + i_2$ ,  $c = q_3 + i_3$  for some  $i_1, i_2, i_3 \in I$ . So  $abc = (q_1 + i_1)(q_2 + i_2)(q_3 + i_3) = q_1q_2q_3 + i_1q_2q_3 + i_2q_1q_3 + i_3q_1q_2 + i_1i_2q_3 + i_2i_3q_1 + i_1i_2i_3$ . Since  $J$  is a  $k$ -ideal of  $S$ , we have  $q_1q_2q_3 \in J$ . Also  $(q_1 + I) \odot (q_2 + I) \odot (q_3 + I) = q_4 + I$  for some unique element  $q_4 \in Q$ , where  $q_1q_2q_3 + I \subseteq q_4 + I$ . Thus  $q_1q_2q_3 + i_4 = q_4 + i_5 \in J$ . Since  $J$  is a  $k$ -ideal,  $q_4 \in J \cap Q$  and  $q_4 + I \in J/I_{(Q \cap J)}$ . By hypothesis,  $q_1 + I, q_2 + I, q_3 + I$  are non-units elements of  $S/I_Q$  and  $(q_1 + I) \odot (q_2 + I) \odot (q_3 + I) \in J/I_{(Q \cap J)}$ . If there exists a unique element  $q \in Q$  such that  $q + I$  is a zero element of  $S/I_Q$  and  $(q_1 + I) \odot (q_2 + I) \odot (q_3 + I) = q + I$ . Thus  $q_1q_2q_3 + r = q + s$  for some  $r, s \in I$ , as  $J$  is a  $k$ -ideal,  $q_1q_2q_3 \in I$ . So  $abc \in I$ , which is a contradiction. Hence  $0 \neq (q_1 + I) \odot (q_2 + I) \odot (q_3 + I) \in J/I_{(Q \cap J)}$ . Since  $J/I_{(Q \cap J)}$  is a weakly 1-absorbing prime ideal, we conclude that either  $q_1q_2 + I \in J/I_{(Q \cap J)}$  or  $q_3 + I \in J/I_{(Q \cap J)}$ .  $(q_1 + I) \odot (q_2 + I) \in J/I_{(Q \cap J)}$  implies  $ab \in J$  or  $q_3 + I \in J/I$  implies  $c \in J$ . Therefore,  $J$  is a weakly 1-absorbing prime ideal of  $S$ . ■

**Theorem 22.** Let  $S$  be a local semiring and  $I$  be a subtractive ideal of  $S$ . If  $I$  is a weakly 1-absorbing prime ideal of  $S$ , then either  $I$  is a 1-absorbing prime ideal of  $S$  or  $I^3 = 0$ .

**Proof.** Suppose that  $I^3 \neq 0$ . Then there exists  $x, y, z \in I$  such that  $xyz \neq 0$ . If possible, let  $I$  is not a weakly 1-absorbing prime ideal of  $S$ . So there exist non-units  $a, b, c \in S$  such that  $abc = 0$  and  $ab \notin I, c \notin I$ .

Since  $S$  is a local semiring, set of non-units forms an ideal. So  $(a + x), (b + y), (c + z)$  are non-units. Consider  $(a + x)(b + y)(c + z) = abc + bcx + acy + abz + ayz + bxz + cxy + xyz \in I$ .

We claim that  $abz = bcx = acy = 0$ . If  $abz \neq 0$ , then  $0 \neq abz = ab(c + z) \in I$ . Since  $I$  is a weakly 1-absorbing prime ideal, either  $ab \in I$  or  $c + z \in I$ . As  $I$  is a subtractive ideal of  $S$ , either  $ab \in I$  or  $c \in I$ , which is a contradiction. Hence we may assume that  $abz = 0$ . In a similar way, one can show  $bcx = acy = 0$ . Also  $ayz = bxz = cxy = 0$ . If suppose  $ayz \neq 0$ , then we have  $0 \neq ayz = ayz + abz + acy + abc = a(y + b)(z + c)$ . Hence either  $a(b + y) \in I$  or  $c + z \in I$ . Since  $I$  is a subtractive ideal, so in both cases we arrive at a contradiction. Thus  $ayz = 0$ . In a similar way,  $bxz = cxy = 0$ .

Therefore,  $0 \neq xyz = (a + x)(b + y)(c + z) \in I$  but  $ab \notin I, c \notin I$  and  $I$  is a subtractive ideal of  $S$ , implies neither  $(a + x)(b + y) \in I$  nor  $c + z \in I$ , which contradicts our hypothesis that  $I$  is a weakly 1-absorbing prime ideal of  $S$ . Therefore,  $I$  is a 1-absorbing prime ideal of  $S$ . ■

**Theorem 23.** *Let  $S$  be a semiring and  $Jac(S)$  be the Jacobson radical of  $S$ . If  $Jac(S)$  is a strong ideal of  $S$ , then for any  $a, b, c \in Jac(S)$ , the ideal  $I = \langle abc \rangle$  is a weakly 1-absorbing prime ideal of  $S$  if and only if  $abc = 0$ .*

**Proof.** If  $abc = 0$  for some  $a, b, c \in Jac(S)$ , then clearly  $I = \langle abc \rangle$  is a weakly 1-absorbing prime ideal of  $S$ . Conversely, assume that  $I$  is a weakly 1-absorbing prime ideal of  $S$ . If possible, let  $abc \neq 0$ . Since  $abc \in I$ , we have either  $ab \in I$  or  $c \in I$ , that is, either  $ab = abcx$  or  $c = abcy$  for some  $x, y \in S$ . Suppose  $ab = abcx$ . As  $c \in Jac(S)$  and  $Jac(S)$  is a strong ideal, so there exists  $c' \in Jac(S)$  such that  $c + c' = 0$ . This implies  $ab(1 + c'x) = 0$  and by [[3], Lemma 3.4],  $(1 + c'x)$  is a semiunit of  $S$ . So there exist  $s, t \in S$  such that  $1 + s(1 + c'x) = t(1 + c'x)$ . Thus  $ab = 0$  and so  $abc = 0$ , which is a contradiction. Also by similar arguments, if  $c \in I$ , we get  $c = 0$ , a contradiction. Therefore  $abc = 0$ . ■

**Theorem 24.** *Let  $S$  be a strong local semiring with maximal ideal  $M$ . Then every proper ideal of  $S$  is a weakly 1-absorbing prime ideal if and only if  $M^3 = \{0\}$ .*

**Proof.** Suppose that every proper ideal of  $S$  is weakly 1-absorbing prime and  $a, b, c \in M$ . Then the ideal  $I = \langle abc \rangle$  is a weakly 1-absorbing prime. By Theorem 23,  $abc = 0$  and so  $M^3 = \{0\}$ . Conversely, suppose that  $M^3 = \{0\}$  and  $I$  is a nonzero proper ideal of  $S$ . So there do not exist any non-units  $a, b, c \in S$  such that  $0 \neq abc \in I$ . Hence,  $I$  is a weakly 1-absorbing prime ideal of  $S$ . ■

**Theorem 25.** *Let  $S = S_1 \times S_2$ , where  $S_1$  and  $S_2$  be commutative semirings. If  $I_1$  is a proper ideal of  $S_1$ . Then the following are equivalent:*

- (1)  $I_1$  is a 1-absorbing prime ideal of  $S_1$ .
- (2)  $I_1 \times S_2$  is a 1-absorbing prime ideal of  $S$ .
- (3)  $I_1 \times S_2$  is a weakly 1-absorbing prime ideal of  $S$ .

**Proof.** (1)  $\implies$  (2) and (2)  $\implies$  (3) are trivial.

(3)  $\implies$  (1) Suppose that  $I_1 \times S_2$  is a weakly 1-absorbing prime ideal of  $S$  and  $abc \in I_1$  for some non-units  $a, b, c \in S_1$ . Then  $0 \neq (abc, 1) = (a, 1)(b, 1)(c, 1) \in I_1 \times S_2$  and  $(a, 1), (b, 1), (c, 1)$  are non-units of  $S$ . Thus either  $(a, 1)(b, 1) = (ab, 1) \in I_1 \times S_2$  or  $(c, 1) \in I_1 \times S_2$ , that is either  $ab \in I_1$  or  $c \in I_1$ . Therefore,  $I_1$  is a 1-absorbing prime ideal of  $S_1$ . ■

**Theorem 26.** *Let  $S = S_1 \times S_2$ , where  $S_1$  and  $S_2$  be commutative semirings but not semifields. If  $I_1$  and  $I_2$  are nonzero ideals of  $S_1$  and  $S_2$  respectively, then the following are equivalent:*

- (1)  $I_1 \times I_2$  is a weakly 1-absorbing prime ideal of  $S$ .
- (2) Either  $I_1$  is a prime ideal of  $S_1$  and  $I_2 = S_2$  or  $I_2$  is a prime ideal of  $S_2$  and  $I_1 = S_1$ .

(3)  $I_1 \times I_2$  is a 1-absorbing prime ideal of  $S$ .

(4)  $I_1 \times I_2$  is a prime ideal of  $S$ .

**Proof.** (1)  $\implies$  (2) Suppose that  $I_1 \times I_2$  is a weakly 1-absorbing prime ideal of  $S$  and  $0 \neq (a, 0) \in I_1 \times I_2$  for some non zero  $a \in I_1$ . Then  $0 \neq (a, 0) = (1, 0)(1, 0)(a, 1) \in I_1 \times I_2$ . Therefore, either  $(1, 0)(1, 0) = (1, 0) \in I_1 \times I_2$  or  $(a, 1) \in I_1 \times I_2$ . So either  $1 \in I_1$  or  $1 \in I_2$ , that is, either  $I_1 = S_1$  or  $I_2 = S_2$ .

Consider  $I_2 = S_2$  and  $ab \in I_1$  for some non-units  $a, b \in S_1$ . Since  $S_2$  is not a semifield, there exists a non-unit  $0 \neq z \in S_2$ . Consider  $0 \neq (ab, z) = (a, 1)(1, z)(b, 1) \in I_1 \times I_2$ , then either  $(a, z) = (a, 1)(1, z) \in I_1 \times I_2$  or  $(b, 1) \in I_1 \times I_2$ . Thus either  $a \in I_1$  or  $b \in I_2$ . Therefore,  $I_1$  is a prime ideal of  $S_1$ . Similarly,  $I_2$  is a prime ideal of  $S_2$  when  $I_1 = S_1$ .

(2)  $\implies$  (3) follows from Theorem 25.

(3)  $\implies$  (4) is clear from the Corollary 1.

(4)  $\implies$  (1) is trivial. ■

#### REFERENCES

- [1] P.J. Allen, *A fundamental theorem of homomorphism for semirings*, Proc. Amer. Math. Soc. **21** (1969) 412–416.  
<https://doi.org/10.1090/S0002-9939-1969-0237575-4>
- [2] P.J. Allen, J. Neggers and H.S. Kim, *Ideal theory in commutative A-semirings*, Kyungpook Math. J. **46** (2006) 261–271.
- [3] S.E. Atani, *The zero-divisor graph with respect to ideals of a commutative semiring*, Glas. Math. **43(63)** (2008) 309–320.  
<https://doi.org/10.3336/gm.43.2.06>
- [4] R.E. Atani, S.E. Atani, *Spectra of semimodules*, Bul. Acad. De Stiinte A Republicii Moldova Math. **67(3)** (2011) 15–28.
- [5] C. Beddani and W. Messirdi, *2-Prime ideals and their applications*, J. Algebra and Its Appl. **15(3)** (2016) 1650051.  
<https://doi.org/10.1142/S0219498816500511>
- [6] H. Behzadipour and P. Nasehpour, *On 2-absorbing ideals of commutative semirings*, J. Algebra and Its Appl. **19(2)** (2020) 2050034.  
<https://doi.org/10.1142/S0219498820500346>
- [7] A.Y. Darani, *On 2-absorbing and weakly 2-absorbing ideals of commutative Semirings*, Kyungpook Math. J. **52(1)** (2021) 91–97.  
<https://doi.org/10.5666/KMJ.2012.52.1.91>
- [8] M.K. Dubey, *Prime and weakly prime ideals in semirings*, Quasigroups and Related Systems **20** (2012) 151–156.
- [9] J.S. Golan, *Semirings and Their Applications*, Kluwer Academic Publishers (Dordrecht, 1999).  
<https://doi.org/10.1007/978-94-015-9333-5>

- [10] B. Khanra, M. Mandal and S. Das, *A note on 2-prime and  $n$ -weakly 2-prime ideals of semirings*, Quasigroups and Related Systems **30(2)** (2022) 241–256.  
<https://doi.org/10.56415/qrs.v30.20>
- [11] S. Koc, Ü. Tekir and E. Yildiz, *On weakly 1-absorbing prime ideals*, Ricerche Mat. **72** (2023) 723–738.  
<https://doi.org/10.1007/s11587-020-00550-4>
- [12] P. Nasehpour, *Some remarks on semirings and their ideals*, Asian-Eur. J. Math. **12(7)** (2019) 2050002.  
<https://doi.org/10.1142/S1793557120500023>
- [13] P. Nasehpour, *Valuation semirings*, J. Algebra Appl. **17(4)** (2018) 1850073.  
<https://doi.org/10.1142/S0219498818500731>
- [14] P. Nasehpour, *Pseudocomplementation and minimal prime ideals in semirings*, Algebra Univ. **79(11)** (2018).  
<https://doi.org/10.1007/s00012-018-0496-x>
- [15] P. Nasehpour, *Some remarks on ideals of commutative semirings*, Quasigroups and Related Systems **26** (2018) 281–298.
- [16] R. Nikandish, M.J. Nikmehr and A. Yassine, *Some results on 1-absorbing primary and weakly 1-absorbing primary ideals of commutative rings*, Bull. Korean Math. Soc. **58(5)** (2021) 1069–1078.  
<https://doi.org/10.4134/BKMS.b190868>
- [17] L. Sawalmeh and M. Saleh, *On 2-absorbing ideals of commutative semirings*, J. Algebra and Its Appl. **22(3)** (2023) 2350063.  
<https://doi.org/10.1142/S0219498823500639>
- [18] A. Yassine, M.J. Nikmehr and R. Nikandish, *On 1-absorbing prime ideals of commutative rings*, J. Algebra and Its Appl. **20(10)** (2021) 2150175.  
<https://doi.org/10.1142/S0219498821501759>